

SPATIAL AND TEMPORAL DISTRIBUTION PATTERNS OF LARVAL SCIAENIDS IN THE ESTUARINE SYSTEM AND ADJACENT CONTINENTAL SHELF OFF SANTOS, SOUTHEASTERN BRAZIL

Renata Roque Porcaro, Maria de Lourdes Zani-Teixeira*, Mario Katsuragawa, Cláudia Namiki, Márcio Hidekazu Ohkawara and Jana Menegassi del Favero

Instituto Oceanográfico da Universidade de São Paulo
(Praça do Oceanográfico, 191, 05508-120 São Paulo, SP, Brasil)

*Corresponding author: zanit@usp.br

<http://dx.doi.org/10.1590/S1679-87592014067606202>

ABSTRACT

Based on data collected during the project "The influence of the Santos-São Vicente estuarine system on the ecosystem of the adjacent continental shelf" (ECOSAN), from Nov/2004 to Mar/2006, the space-temporal distribution pattern of Sciaenidae larvae in relation to biotic and abiotic factors was investigated. Bongo net was used for sampling, aboard the R/V "Prof. W. Besnard" on the shelf, and the boats "Veliger II" and "Albacora" in the estuarine region. Twelve taxa were identified: *Bairdiella ronchus*, *Ctenosciaena gracilicirrhus*, *Cynoscion* spp., *Isopisthus parvipinnis*, *Macrodon atricauda*, *Micropogonias furnieri*, *Menticirrhus* spp., *Nebris microps*, *Ophioscion punctatissimus*, *Paralichthys brasiliensis*, *Stellifer rastrifer* and *Stellifer* spp. Sciaenidae larvae were widespread on the continental shelf in both campaigns (frequency of occurrence >80%). They were most abundant in Mar/2006 (mean = 6.47 larvae.m⁻²; sd = 8.36) and least abundant in Sep/2005 (mean = 2.37 larvae.m⁻²; sd = 3.33). The average abundance in the estuarine region ranged from 0.12 larvae.m⁻² (sd = 0.11) in Jun/2005 to 4.28 larvae.m⁻² (sd = 1.99) in Nov/2004. Frequency of occurrence minimum occurred in Aug/2005 (50%) and the maximum in Nov/2004 (100%). The presence of Sciaenidae larvae in most locations and periods suggests that the reproductive process of this family in the region is continuous, spawning being more intense in the warmer months.

RESUMO

Baseando-se em dados coletados durante o projeto "A Influência do Complexo Estuarino da Baixada Santista sobre o Ecossistema da Plataforma Continental Adjacente" (ECOSAN), nov/2004 - mar/2006, foram investigados os padrões de distribuição espaço-temporal das larvas de Sciaenidae relacionando-os com fatores bióticos e abióticos. A rede bongô foi utilizada como amostrador do icteoplâncton, a bordo de três embarcações: o N/Oc "Prof. W. Besnard" na plataforma, e os barcos "Veliger II" e "Albacora" na baía e canais. Doze táxons foram identificados: *Bairdiella ronchus*, *Ctenosciaena gracilicirrhus*, *Cynoscion* spp., *Isopisthus parvipinnis*, *Macrodon atricauda*, *Micropogonias furnieri*, *Menticirrhus* spp., *Nebris microps*, *Ophioscion punctatissimus*, *Paralichthys brasiliensis*, *Stellifer rastrifer* and *Stellifer* spp. Na plataforma continental as larvas de cieniídeos tiveram ampla distribuição nas duas campanhas (frequência de ocorrência > 80%), sendo mais abundantes em mar/2006 (6,47±8,36 larvas.m⁻²) e menos abundantes em set/2005 (2,37±3,33 larvas.m⁻²). No estuário, a abundância média das larvas de cieniídeos variou entre 0,12±0,11 larvas.m⁻² (jun/2005) e 4,28±1,99 larvas.m⁻² (nov/2004); a frequência de ocorrência foi mínima em ago/2005 (50%) e máxima em nov/2004 (100%). A presença das larvas na maioria das estações de coleta do sistema estuarino de Santos e plataforma continental adjacente sugere uma atividade reprodutiva contínua, com desova mais intensa nos meses mais quentes.

Descriptors: Sciaenidae, Ichthyoplankton, Abundance, Environmental variables.
Descritores: Sciaenidae, Ictioplâncton, Abundância, Variáveis ambientais.

INTRODUCTION

Sciaenidae is a large family of about 70 genera and 270 species. They inhabit marine, brackish, and freshwater areas of the Atlantic, Indian, and

Pacific oceans (NELSON, 2006). Most of the species of this family are classified as demersal fish, but some may feed in the pelagic environment. At least 54 species of Sciaenidae have been described on the Brazilian coast (CASATTI; MENEZES, 2003), some

of them being important fishery resources. On the inner shelf of South and Southeastern Brazil the species *Micropogonias furnieri* (Desmarest, 1823), *Macrodon atricauda* (Günther, 1880), *Cynoscion guatucupa* (Cuvier, 1830), *Cynoscion jamaicensis* (Vaillant & Bocourt, 1883), and *Menticirrhus americanus* (Linnaeus, 1758) are among those most exploited by commercial fisheries (VALENTINI; PEZZUTO, 2006).

Many fishery resources, including some sciaenids, spawn in the ocean and the early stages are transported to estuaries, where the larvae may find a suitable environment for their survival and growth (GODEFROID et al., 1999; COSTA; ARAÚJO, 2003).

Larval fish usually are planktonic and they are in general morphologically very distinct from the corresponding adults. The transformation takes place gradually until they settle down in the adult environment. The knowledge of this process is important in enabling us to understand the population dynamics of fishes, as the size of adult stocks depends directly on their success in surviving during the early developmental stages (HANNA, 1998). The distribution of larval fish is heterogeneous and can vary in time and space, depending mainly on the fishes' reproductive strategies, including the place, time and type of spawning, adult abundance, the duration of the early stages, larval behavior and the presence of potential predators and prey, among other factors (NORCROSS; SHAW, 1984; LEIS, 1991; SASSA et al., 2004). The larval distribution is also influenced by climate and hydrographic events such as currents, winds, vortexes and the upwelling and stratification of the water column (DENMAN; POWELL, 1984; NORCROSS; SHAW, 1984).

Although the coastal area of the present study is known to be heavily impacted by human activities such as industrialization, port activities and urbanization (CETESB, 2001), the bay and the estuary in the Santos region interact with each other and form an environment recognized as an important nursery area for fish, crustaceans and molluscs (SCHAEFFER-NOVELLI; CINTRON, 1986). High diversity and abundance of fish have been observed in this region (PAIVA FILHO et al., 1987) and the sciaenids have been described as one of most important groups (PAIVA FILHO; SCHMIEGELOW, 1986). In the Santos Bay, Schmidt and Dias (2012) sampled 21 species of Sciaenidae, with *Stellifer rastrifer* (Jordan, 1889) representing 70.4% of the total composition.

Previous studies on the early life history of the sciaenids off Southern and Southeastern Brazil include those by Matsuura and Nakatani (1979) who

described the development of larval *Umbrina coroides* Cuvier, 1830 near Anchieta island; Sinque (1980) who studied the larval and juvenile development of *Cynoscion leiarchus* (Cuvier, 1830), *Menticirrhus americanus*, *Micropogonias furnieri*, *Stellifer rastrifer*, *Macrodon ancylodon* (Bloch & Schneider, 1801) and *Isopisthus parvipinnis* (Cuvier, 1830) in the Cananéia-Iguape estuary; Katsuragawa et al. (1993) who described the larval distribution of *Menticirrhus americanus* at Ubatuba; Godefroid et al. (2001) who described the larval and juvenile occurrence of *Menticirrhus americanus*, *Menticirrhus littoralis* (Holbrook, 1847), *Umbrina coroides* and *Micropogonias furnieri* in the Pontal do Sul region; Bruno and Muelbert (2002; 2009) who investigated the distribution and abundance patterns of eggs and larvae of *Micropogonias furnieri* in the Patos lagoon, and Itagaki et al. (2007) who described the larval and juvenile development of *Bairdiella ronchus* (Cuvier, 1830) in the Cananéia-Iguape estuary. However, no studies on early sciaenids have been undertaken in the estuarine and coastal area around Santos.

Therefore, the present work analyzed the spatial and temporal distribution patterns of the Sciaenidae larvae and evaluated the influence of abiotic and biotic factors in the Santos estuarine system and on the adjacent continental shelf. We also discuss the importance of the study area to early life stages of some Sciaenidae species.

MATERIAL AND METHODS

Study Area and Sampling Techniques

The Baixada Santista, located on the central coast of São Paulo state - Brazil, covers the São Vicente and Santos estuarine system, the Santos Bay and the adjacent Continental Shelf (CETESB, 2001). The Santos Bay has a fundamental role in the dilution of materials from the channels of Santos and São Vicente, including those from the submarine emissary (PIRES-VANIN et al., 2008). The Bertioga channel is an environment whose margins are better preserved than those of the Santos and São Vicente channels, even though the first is subject to habitat destruction as a result of urban occupation and the pollution that reaches its shores (PIRES-VANIN et al., 2008). The adjacent continental shelf is complex, due its proximity to the coastal region and a greater number of variables involved in the oceanographic structure of the region. Variations in water temperature and salinity in this environment demonstrate the influence of different water masses throughout the year: Tropical Water (TW), Coastal Water (CW) and South

Atlantic Central Water (SACW) (MIRANDA; KATSURAGAWA, 1991; CASTRO et al., 2006).

Samples were collected during research cruises of the project "The influence of the Santos-São Vicente estuarine system on the ecosystem of the adjacent continental shelf" (ECOSAN), by the Oceanographic Institute of São Paulo University (IOUSP). In three localities in the Santos - São Vicente estuarine system (Santos Bay, Port Channel and Bertioga Channel) the cruises were conducted monthly at 12 oceanographic stations from November

2004 to March 2006; on the adjacent continental shelf (North and South areas) out to the 50 m isobath sampling plan included 37 and 36 oceanographic stations, respectively in September/2005 and March/2006 (Fig. 1). Oceanographic cruises were carried out by the R/V "Prof. W. Besnard" on the continental shelf, and the boats "Veliger II" and "Albacora" in the bay and channels. Summary information of each sampling station is presented in Tables 1 and 2.




Fig. 1. Location of the sampling stations in the Santos estuarine system (Santos Bay, Port and Bertioga Channel) from November 2004 to December 2005 (A) and on the continental shelf between Peruíbe and south of São Sebastião island in September 2005 and March 2006 (B).

Table 1. Data of sampling stations carried out in Santos bay, Bertioga channel and Port channel in Santos-São Vicente estuarine system, Brazil, from November 2004 to December 2005 (LD - Local Depth (m); SD - Sampler Depth (m); T - Temperature; S - Salinity obtained at maximum depth of 5 m).

Santos bay

| Date | Hour | Long. | Lat. | LD | SD | T* | S* | Date | Hour | Long. | Lat. | LD | SD | T* | S* |
|------------|-------|---------|---------|------|------|-------|-------|------------|-------|---------|---------|------|------|-------|-------|
| 11/24/2004 | 11:58 | -46.376 | -24.011 | 10.4 | 9.0 | 22.96 | 35.59 | 5/21/2005 | 07:50 | -46.330 | -24.018 | 12.8 | 9.0 | 25.3 | 34.56 |
| 11/24/2004 | 11:23 | -46.362 | -23.986 | 7.6 | 6.0 | 22.24 | 35.72 | 5/21/2005 | 11:06 | -46.325 | -23.988 | 7.6 | 3.5 | 25.16 | 34.86 |
| 11/24/2004 | 13:14 | -46.330 | -24.018 | 13.4 | 12.0 | 22.26 | 35.25 | 6/24/2005 | 10:09 | -46.376 | -24.011 | 8.5 | 7.0 | 23.41 | 34.08 |
| 11/24/2004 | 10:14 | -46.325 | -23.988 | 6.9 | 5.0 | 22.04 | 35.72 | 6/24/2005 | 10:53 | -46.362 | -23.986 | 6.0 | 5.0 | 23.04 | 32.63 |
| 12/17/2004 | 13:19 | -46.375 | -24.011 | 11.0 | 9.0 | 24.03 | 35.43 | 6/24/2005 | 07:22 | -46.330 | -24.018 | 12.2 | 11.0 | 23.05 | 32.44 |
| 12/17/2004 | 10:07 | -46.363 | -23.988 | 7.9 | 6.0 | 24.74 | 34.84 | 6/24/2005 | 12:52 | -46.325 | -23.988 | 5.8 | 4.5 | 23.55 | 34.04 |
| 12/17/2004 | 14:45 | -46.329 | -24.016 | 13.0 | 11.5 | 24.79 | 34.20 | 7/23/2005 | 10:17 | -46.376 | -24.011 | 8.5 | 7.0 | 21.75 | 34.51 |
| 12/17/2004 | 09:27 | -46.325 | -23.989 | 7.0 | 4.5 | 24.84 | 33.16 | 7/23/2005 | 11:27 | -46.362 | -23.986 | 6.0 | 4.5 | 21.7 | 34.46 |
| 1/25/2005 | 11:03 | -46.375 | -24.011 | 10.5 | 8.0 | 25.43 | 34.00 | 7/23/2005 | 07:32 | -46.330 | -24.018 | 12.5 | 11.5 | 21.63 | 34.45 |
| 1/25/2005 | 12:15 | -46.363 | -23.988 | 7.6 | 5.0 | 24.77 | 34.94 | 7/23/2005 | 12:35 | -46.325 | -23.988 | 5.5 | 4.5 | 21.64 | 34.56 |
| 1/25/2005 | 09:29 | -46.329 | -24.016 | 12.0 | 9.0 | 25.33 | 34.34 | 8/20/2005 | 09:35 | -46.376 | -24.011 | 8.5 | 7.5 | 21.81 | 33.86 |
| 1/25/2005 | 12:48 | -46.325 | -23.989 | 7.3 | 4.5 | 25.04 | 34.51 | 8/20/2005 | 08:30 | -46.362 | -23.986 | 6.0 | 5.0 | 21.8 | 33.91 |
| 2/24/2005 | 13:04 | -46.376 | -24.011 | 11.0 | 9.0 | 25.25 | 34.31 | 8/20/2005 | 11:30 | -46.330 | -24.018 | 12.0 | 11.0 | 21.77 | 33.79 |
| 2/24/2005 | 12:09 | -46.362 | -23.986 | 7.6 | 5.5 | 25.61 | 34.03 | 8/20/2005 | 12:55 | -46.325 | -23.988 | 6.0 | 5.0 | 21.69 | 33.84 |
| 2/24/2005 | 14:25 | -46.330 | -24.018 | 13.7 | 11.0 | 26.27 | 34.17 | 10/1/2005 | 09:22 | -46.376 | -24.011 | 10.0 | 10.0 | 20.38 | 32.59 |
| 2/24/2005 | 07:44 | -46.325 | -23.988 | 6.4 | 4.5 | 25.59 | 33.54 | 10/1/2005 | 10:40 | -46.362 | -23.986 | 7.0 | 7.0 | 20.28 | 32.73 |
| 3/22/2005 | 13:20 | -46.376 | -24.011 | 11.0 | 8.5 | 26.57 | 34.80 | 10/1/2005 | 07:55 | -46.330 | -24.018 | 12.8 | 12.0 | 20.26 | 32.86 |
| 3/22/2005 | 11:18 | -46.362 | -23.986 | 7.6 | 5.0 | 27.51 | 33.82 | 10/1/2005 | 11:23 | -46.325 | -23.988 | 7.0 | 7.0 | 20.28 | 32.96 |
| 3/22/2005 | 14:34 | -46.330 | -24.018 | 12.8 | 10.0 | 26.78 | 34.25 | 12/13/2005 | 09:45 | -46.376 | -24.011 | 9.0 | 6.5 | 24.37 | 33.37 |
| 3/22/2005 | 12:00 | -46.325 | -23.988 | 6.7 | 3.5 | 26.82 | 34.38 | 12/13/2005 | 10:40 | -46.362 | -23.986 | 6.5 | 5.5 | 23.95 | 32.02 |
| 5/21/2005 | 09:50 | -46.376 | -24.011 | 10.7 | 8.0 | 25.5 | 35.01 | 12/13/2005 | 07:55 | -46.330 | -24.018 | 12.0 | 7.5 | 24.12 | 32.69 |
| 5/21/2005 | 10:25 | -46.362 | -23.986 | 7.6 | 5.0 | 25.4 | 34.72 | 12/13/2005 | 11:31 | -46.325 | -23.988 | 6.0 | 5.0 | 24.24 | 33.44 |

Bertioga channel

| Date | Hour | Long. | Lat. | LD | SD | T* | S* | Date | Hour | Long. | Lat. | LD | SD | T* | S* |
|------------|-------|---------|---------|------|------|-------|-------|------------|-------|---------|---------|------|------|-------|-------|
| 11/25/2004 | 08:33 | -46.133 | -23.857 | 11.9 | 8.0 | 24.78 | 30.89 | 5/22/2005 | 09:19 | -46.133 | -23.857 | 8.8 | 4.0 | 25.24 | 33.18 |
| 11/25/2004 | 09:03 | -46.148 | -23.860 | 3.0 | 2.0 | 24.84 | 30.80 | 5/22/2005 | 10:32 | -46.148 | -23.860 | 6.1 | 4.0 | 25.11 | 32.49 |
| 11/25/2004 | 09:30 | -46.156 | -23.866 | 5.0 | 4.0 | 24.82 | 30.45 | 5/22/2005 | 10:55 | -46.156 | -23.866 | 6.4 | 3.5 | 25.01 | 31.49 |
| 11/25/2004 | 11:35 | -46.165 | -23.878 | 6.7 | 5.5 | 24.84 | 30.91 | 5/22/2005 | 11:15 | -46.165 | -23.878 | 7.3 | 4.5 | 26.27 | 27.76 |
| 11/25/2004 | 12:10 | -46.186 | -23.891 | 4.3 | 3.0 | 24.90 | 29.19 | 5/22/2005 | 11:42 | -46.186 | -23.891 | 7.9 | 4.5 | 24.74 | 25.64 |
| 11/25/2004 | 12:34 | -46.193 | -23.902 | 7.0 | 6.0 | 25.40 | 27.36 | 5/22/2005 | 12:08 | -46.193 | -23.902 | 11.0 | 8.0 | 24.89 | 25.94 |
| 12/18/2004 | 07:05 | -46.133 | -23.857 | 12.5 | 12.0 | 26.07 | 34.16 | 6/25/2005 | 09:40 | -46.133 | -23.857 | 12.8 | 11.0 | 22.95 | 29.00 |
| 12/18/2004 | 08:45 | -46.151 | -23.859 | 5.2 | 4.0 | 26.10 | 33.99 | 6/25/2005 | 10:45 | -46.148 | -23.860 | 12.7 | 1.5 | -- | -- |
| 12/18/2004 | 09:25 | -46.157 | -23.867 | 5.5 | 5.0 | 26.11 | 33.82 | 6/25/2005 | 11:18 | -46.156 | -23.866 | 4.3 | 3.0 | 23.08 | 29.18 |
| 12/18/2004 | 09:55 | -46.165 | -23.878 | 6.4 | 6.0 | 26.08 | 32.92 | 6/25/2005 | 11:53 | -46.165 | -23.878 | 5.2 | 4.0 | 23.41 | -- |
| 12/18/2004 | 13:05 | -46.188 | -23.892 | 6.7 | 4.0 | 26.08 | 31.44 | 6/25/2005 | 12:25 | -46.186 | -23.891 | 3.4 | 2.5 | -- | 31.25 |
| 12/18/2004 | 13:36 | -46.193 | -23.902 | 7.0 | 4.0 | 26.32 | 26.48 | 6/25/2005 | 12:45 | -46.193 | -23.902 | 5.2 | 4.0 | 22.45 | 24.97 |
| 1/26/2005 | 09:15 | -46.133 | -23.857 | 12.8 | 10.0 | 25.69 | 21.56 | 7/24/2005 | 09:46 | -46.133 | -23.857 | 11.5 | 10.0 | 21.96 | 31.39 |
| 1/26/2005 | 10:29 | -46.151 | -23.859 | 5.8 | 4.0 | 25.56 | 21.17 | 7/24/2005 | 11:04 | -46.148 | -23.860 | 3.0 | 1.0 | -- | -- |
| 1/26/2005 | 12:00 | -46.157 | -23.867 | 5.8 | 4.0 | 25.69 | 21.56 | 7/24/2005 | 11:35 | -46.156 | -23.866 | 4.0 | 2.0 | 21.78 | 29.41 |
| 1/26/2005 | 12:14 | -46.165 | -23.878 | 7.3 | 5.0 | 25.76 | 19.90 | 7/24/2005 | 12:00 | -46.165 | -23.878 | 5.2 | 3.0 | 21.61 | 27.48 |
| 1/26/2005 | 12:42 | -46.188 | -23.892 | 5.5 | 4.0 | 25.78 | 18.99 | 7/24/2005 | 12:32 | -46.186 | -23.891 | 3.0 | 1.0 | -- | -- |
| 1/26/2005 | 13:01 | -46.193 | -23.902 | 6.4 | 5.0 | 25.78 | 18.99 | 7/24/2005 | 14:50 | -46.193 | -23.902 | 6.0 | 4.5 | 21.74 | 26.90 |
| 2/25/2005 | 08:22 | -46.133 | -23.857 | 13.8 | 8.5 | 27.29 | 30.76 | 8/21/2005 | 08:00 | -46.133 | -23.857 | 11.0 | 10.0 | 22.42 | 32.81 |
| 2/25/2005 | 09:37 | -46.148 | -23.860 | 5.0 | 3.0 | 27.39 | 28.15 | 8/21/2005 | 09:55 | -46.148 | -23.860 | 2.7 | 1.5 | -- | -- |
| 2/25/2005 | 10:02 | -46.156 | -23.866 | 4.6 | 3.5 | 27.58 | 27.02 | 8/21/2005 | 10:40 | -46.156 | -23.866 | 4.3 | 3.0 | -- | -- |
| 2/25/2005 | 10:26 | -46.165 | -23.878 | 5.8 | 4.0 | 27.42 | 28.08 | 8/21/2005 | 11:25 | -46.165 | -23.878 | 5.0 | 4.0 | 22.72 | 28.21 |
| 2/25/2005 | 12:40 | -46.186 | -23.891 | 7.3 | 5.0 | 27.63 | 26.37 | 8/21/2005 | 14:25 | -46.186 | -23.891 | 4.6 | 3.5 | -- | -- |
| 2/25/2005 | 13:10 | -46.193 | -23.902 | 10.0 | 7.0 | 27.63 | 26.37 | 8/21/2005 | 14:58 | -46.193 | -23.902 | 6.7 | 5.5 | 22.99 | 28.53 |
| 3/23/2005 | 08:05 | -46.133 | -23.857 | 11.3 | 8.0 | 27.52 | 31.18 | 10/2/2005 | 08:30 | -46.133 | -23.857 | 12.0 | 12.0 | 21.81 | 28.43 |
| 3/23/2005 | -- | -46.148 | -23.860 | 12.2 | -- | 27.62 | 30.42 | 10/2/2005 | 09:24 | -46.148 | -23.860 | 5.0 | 5.0 | 21.77 | 26.43 |
| 3/23/2005 | 09:35 | -46.156 | -23.866 | 6.4 | 3.0 | 27.52 | 30.33 | 10/2/2005 | 10:46 | -46.156 | -23.866 | 5.5 | 5.0 | 21.71 | 26.41 |
| 3/23/2005 | 09:59 | -46.165 | -23.878 | 5.8 | 2.5 | 27.52 | 29.36 | 10/2/2005 | 10:10 | -46.165 | -23.878 | 6.7 | 6.0 | 21.99 | 25.12 |
| 3/23/2005 | 10:28 | -46.186 | -23.891 | 6.4 | 4.0 | 27.89 | 27.15 | 10/2/2005 | 12:25 | -46.186 | -23.891 | 8.0 | 7.0 | 21.78 | 24.68 |
| 3/23/2005 | 10:57 | -46.193 | -23.902 | 5.8 | 5.0 | 28.05 | 26.32 | 10/2/2005 | 12:55 | -46.193 | -23.902 | 10.0 | 10.0 | 22.01 | 23.70 |
| 4/27/2005 | 09:03 | -46.133 | -23.857 | 13.4 | 12.0 | 24.07 | 18.42 | 12/14/2005 | 09:10 | -46.133 | -23.857 | 11.9 | 9.0 | 24.01 | 32.94 |
| 4/27/2005 | 10:36 | -46.148 | -23.860 | 6.4 | 6.4 | -- | -- | 12/14/2005 | 11:13 | -46.148 | -23.860 | 3.0 | 1.0 | -- | -- |
| 4/27/2005 | 10:59 | -46.156 | -23.866 | 5.8 | 5.0 | 24.74 | 21.61 | 12/14/2005 | 11:40 | -46.156 | -23.866 | 4.5 | 2.5 | 23.88 | 31.50 |
| 4/27/2005 | 11:22 | -46.165 | -23.878 | 6.7 | 6.7 | 24.33 | 20.06 | 12/14/2005 | 12:05 | -46.165 | -23.878 | 5.5 | 3.5 | 23.81 | 26.72 |
| 4/27/2005 | 11:49 | -46.186 | -23.891 | 6.1 | 6.4 | 24.21 | 19.76 | 12/14/2005 | 15:13 | -46.186 | -23.891 | 4.0 | 2.0 | -- | -- |
| 4/27/2005 | 12:10 | -46.193 | -23.902 | 7.9 | 9.5 | 24.15 | 19.44 | 12/14/2005 | 15:40 | -46.193 | -23.902 | 5.0 | 3.0 | 23.99 | 25.08 |

Port channel

| Date | Hour | Long. | Lat. | LD | SD | T* | S* | Date | Hour | Long. | Lat. | LD | SD | T* | S* |
|------------|-------|---------|---------|------|------|-------|-------|-----------|-------|---------|---------|------|------|-------|-------|
| 11/24/2004 | 07:43 | -46.300 | -23.993 | 20.0 | 10.0 | 23.73 | 31.71 | 5/21/2005 | 12:22 | -46.300 | -23.993 | 21.0 | 18.0 | 25.39 | 34.30 |
| 11/24/2004 | 08:41 | -46.303 | -23.959 | 14.0 | 10.0 | 23.52 | 32.42 | 5/21/2005 | 13:42 | -46.303 | -23.959 | 14.0 | 10.0 | 25.32 | 34.34 |
| 12/17/2004 | 07:56 | -46.299 | -23.993 | 18.0 | 14.5 | 24.97 | 32.79 | 6/24/2005 | 13:50 | -46.300 | -23.993 | 16.0 | 14.0 | 23.45 | 32.62 |
| 12/17/2004 | 06:31 | -46.303 | -23.959 | 14.0 | 12.0 | 25.24 | 30.76 | 6/24/2005 | 14:59 | -46.303 | -23.959 | 13.0 | 12.0 | 23.41 | 32.49 |
| 1/25/2005 | 08:18 | -46.299 | -23.993 | 12.5 | 13.0 | 25.63 | 29.47 | 7/23/2005 | 13:30 | -46.300 | -23.993 | 15.0 | 13.5 | 21.83 | 34.31 |
| 1/25/2005 | 06:43 | -46.303 | -23.959 | 13.1 | 11.0 | 25.55 | 31.67 | 7/23/2005 | 15:09 | -46.303 | -23.959 | 13.0 | 11.5 | 21.81 | 32.97 |
| 2/24/2005 | 08:45 | -46.300 | -23.993 | 19.5 | 18.0 | 26.00 | 32.39 | 8/20/2005 | 14:04 | -46.300 | -23.993 | 15.0 | 14.0 | 22.09 | 33.54 |
| 2/24/2005 | 09:40 | -46.303 | -23.959 | 13.1 | 11.0 | 26.47 | 30.53 | 8/20/2005 | 15:50 | -46.303 | -23.959 | 13.0 | 15.0 | 22.01 | 33.41 |
| 3/22/2005 | 09:44 | -46.300 | -23.993 | 18.0 | 15.0 | 26.83 | 31.57 | 10/1/2005 | 13:10 | -46.300 | -23.993 | 14.0 | 14.0 | 20.72 | 31.64 |
| 3/22/2005 | 08:00 | -46.303 | -23.959 | 13.1 | 10.0 | 26.63 | 30.64 | 10/1/2005 | 14:40 | -46.303 | -23.959 | 14.0 | 14.0 | 20.83 | 30.95 |
| 4/26/2005 | 07:38 | -46.300 | -23.993 | 18.5 | 16.5 | 26.61 | | | | | | | | | |

Table 2. Data of sampling stations of catches carried out on the continental shelf adjacent to the Santos-São Vicente estuarine system, Brazil, in September 2005 and March 2006 (LD - Local Depth (m); SD - Sampler Depth (m); T - Temperature; S - Salinity obtained at 5 m depth).

| Date | Hour | Long. | Lat. | LD | SD | T* | S* | Date | Hour | Long. | Lat. | LD | SD | T* | S* |
|-----------|-------|---------|---------|----|----|-------|-------|-----------|-------|---------|---------|----|----|-------|-------|
| 9/21/2005 | 20:30 | -45.560 | -24.226 | 59 | 40 | 20.47 | 34.22 | 3/12/2006 | 06:45 | -45.592 | -23.820 | 15 | 10 | 26.82 | 33.92 |
| 9/22/2005 | 01:23 | -45.707 | -23.971 | 30 | 25 | 20.78 | 34.14 | 3/12/2006 | 09:22 | -45.505 | -23.966 | 28 | 20 | 27.04 | 34.21 |
| 9/22/2005 | 03:20 | -45.790 | -23.829 | 20 | 20 | 20.47 | 33.06 | 3/12/2006 | 11:30 | -45.560 | -24.226 | 41 | 35 | 27.11 | 34.84 |
| 9/22/2005 | 05:10 | -45.991 | -23.836 | 15 | 10 | 20.61 | 32.68 | 3/12/2006 | 13:55 | -45.707 | -23.971 | 50 | 45 | 27.18 | 34.95 |
| 9/22/2005 | 06:43 | -46.039 | -23.918 | 18 | 15 | 20.43 | 32.11 | 3/13/2006 | 14:57 | -45.790 | -23.829 | 53 | 45 | 26.81 | 34.99 |
| 9/22/2005 | 09:00 | -45.899 | -23.992 | 28 | 20 | 20.62 | 33.77 | 3/13/2006 | 17:50 | -45.991 | -23.836 | 43 | 35 | 27.35 | 35.06 |
| 9/22/2005 | 11:19 | -45.820 | -24.119 | 41 | 35 | 20.62 | 33.97 | 3/13/2006 | 21:50 | -46.039 | -23.918 | 29 | 24 | 27.31 | 34.17 |
| 9/22/2005 | 15:27 | -45.701 | -24.328 | 60 | 50 | 20.27 | 33.52 | 3/13/2006 | 23:25 | -45.899 | -23.992 | 15 | 10 | 27.54 | 33.90 |
| 9/22/2005 | 19:27 | -45.880 | -24.373 | 57 | 40 | 19.66 | 33.38 | 3/14/2006 | 01:23 | -45.820 | -24.119 | 15 | 10 | 27.55 | 33.94 |
| 9/22/2005 | 22:00 | -45.965 | -24.226 | 46 | 30 | 20.31 | 33.88 | 3/14/2006 | 03:29 | -45.701 | -24.328 | 30 | 25 | 26.76 | 34.78 |
| 9/23/2005 | 01:25 | -46.053 | -24.077 | 30 | 25 | 20.53 | 33.61 | 3/14/2006 | 05:50 | -45.880 | -24.373 | 45 | 35 | 26.76 | 35.18 |
| 9/23/2005 | 03:02 | -46.101 | -24.000 | 20 | 15 | 20.41 | 31.77 | 3/14/2006 | 08:26 | -45.965 | -24.226 | 55 | 50 | 26.08 | 34.91 |
| 9/23/2005 | 04:42 | -46.178 | -24.042 | 20 | 15 | 20.49 | 31.75 | 3/14/2006 | 11:02 | -46.053 | -24.077 | 55 | 50 | 26.12 | 35.00 |
| 9/23/2005 | 06:18 | -46.257 | -24.084 | 25 | 25 | 20.34 | 31.17 | 3/14/2006 | 13:30 | -46.101 | -24.000 | 40 | 35 | 26.74 | 35.12 |
| 9/23/2005 | 09:10 | -46.209 | -24.164 | 32 | 25 | 20.29 | 32.50 | 3/14/2006 | 16:20 | -46.178 | -24.042 | 33 | 28 | 27.08 | 34.79 |
| 9/23/2005 | 14:16 | -46.120 | -24.314 | 41 | 25 | 20.19 | 33.74 | 3/14/2006 | 19:05 | -46.257 | -24.084 | 16 | 11 | 28.06 | 33.95 |
| 9/23/2005 | 16:44 | -46.034 | -24.460 | 57 | 40 | 20.06 | 33.57 | 3/14/2006 | 20:27 | -46.209 | -24.164 | 20 | 15 | 27.66 | 33.98 |
| 9/23/2005 | 19:24 | -46.188 | -24.548 | 55 | 40 | 19.60 | 33.42 | 3/14/2006 | 21:48 | -46.120 | -24.314 | 24 | 15 | 27.51 | 34.40 |
| 9/23/2005 | 21:30 | -46.274 | -24.400 | 40 | 25 | -- | -- | 3/14/2006 | 23:05 | -46.034 | -24.460 | 31 | 25 | 27.53 | 34.21 |
| 9/23/2005 | 23:40 | -46.362 | -24.252 | 31 | 15 | 20.65 | 32.20 | 3/15/2006 | 01:38 | -46.188 | -24.548 | 42 | 35 | 25.71 | 35.12 |
| 9/24/2005 | 01:31 | -46.357 | -24.088 | 20 | 20 | 20.60 | 31.57 | 3/15/2006 | 04:15 | -46.274 | -24.400 | 57 | 50 | 25.96 | 34.96 |
| 9/24/2005 | 02:46 | -46.458 | -24.091 | 16 | 10 | 20.51 | 31.26 | 3/15/2006 | 08:15 | -46.362 | -24.252 | 57 | 52 | 25.17 | 35.22 |
| 9/24/2005 | 04:20 | -46.617 | -24.170 | 14 | 10 | 20.60 | 31.31 | 3/15/2006 | 10:28 | -46.357 | -24.088 | 46 | 40 | 26.14 | 35.12 |
| 9/24/2005 | 06:23 | -46.517 | -24.338 | 30 | 25 | 20.51 | 33.43 | 3/15/2006 | 13:10 | -46.458 | -24.091 | 28 | 20 | 27.44 | 34.45 |
| 9/24/2005 | 08:19 | -46.430 | -24.486 | 45 | 40 | 19.67 | 33.35 | 3/15/2006 | 15:20 | -46.617 | -24.170 | 21 | 15 | 27.47 | 34.51 |
| 9/24/2005 | 10:32 | -46.342 | -24.635 | 55 | 50 | 19.65 | 33.26 | 3/15/2006 | 16:38 | -46.517 | -24.338 | 21 | 15 | 27.56 | 34.38 |
| 9/24/2005 | 12:45 | -46.497 | -24.721 | 53 | 35 | 20.04 | 33.29 | 3/15/2006 | 17:58 | -46.430 | -24.486 | 18 | 10 | 27.60 | 34.39 |
| 9/24/2005 | 14:43 | -46.584 | -24.576 | 43 | 25 | 19.88 | 33.36 | 3/15/2006 | 19:02 | -46.342 | -24.635 | 15 | 10 | 27.35 | 34.56 |
| 9/24/2005 | 16:35 | -46.670 | -24.426 | 30 | 15 | 20.36 | 33.43 | 3/15/2006 | 20:50 | -46.497 | -24.721 | 27 | 22 | 26.49 | 35.03 |
| 9/24/2005 | 18:19 | -46.768 | -24.262 | 14 | 5 | 20.14 | 32.63 | 3/15/2006 | 22:45 | -46.584 | -24.576 | 40 | 30 | 26.34 | 34.98 |
| 9/24/2005 | 19:44 | -46.914 | -24.368 | 15 | 5 | 20.11 | 31.74 | 3/16/2006 | 05:10 | -46.670 | -24.426 | 60 | 55 | 25.78 | 34.99 |
| 9/24/2005 | 21:29 | -46.825 | -24.514 | 28 | 15 | 20.38 | 32.72 | 3/16/2006 | 07:25 | -46.768 | -24.262 | 60 | 55 | 26.07 | 34.88 |
| 9/24/2005 | 23:28 | -46.740 | -24.663 | 41 | 25 | 20.09 | 33.19 | 3/16/2006 | 10:20 | -46.914 | -24.368 | 31 | 25 | 25.62 | 35.03 |
| 9/25/2005 | 01:31 | -46.653 | -24.809 | 49 | 45 | 19.99 | 33.23 | 3/16/2006 | 13:20 | -46.825 | -24.514 | 19 | 15 | 27.23 | 34.65 |
| 9/20/2005 | 08:53 | -45.592 | -23.820 | 18 | 17 | 20.82 | 34.26 | 3/16/2006 | 15:00 | -46.740 | -24.663 | 18 | 14 | 27.58 | 34.41 |
| 9/20/2005 | 15:12 | -45.505 | -23.966 | 42 | 40 | 20.54 | 34.39 | 3/16/2006 | 17:00 | -46.653 | -24.809 | 43 | 38 | 26.75 | 34.67 |
| | | | | | | | | 3/16/2006 | 19:20 | -45.405 | -24.105 | 54 | 50 | 26.68 | 34.80 |

According to Smith and Richardson (1977), ichthyoplankton was collected by oblique tows with a paired 60 cm bongo net with 333 mm and 505 mm mesh size. The net was towed in double oblique hauls at approximately 2 knots. Water volumes filtered were determined from flowmeters mounted at the centers of the nets. All samples were fixed in buffered 4% formaldehyde-seawater solution.

Temperature and salinity horizontal profiles, that were used to recognize the influence of the water masses, were taken at each station using CTD (Sea-Bird Electronics, Inc., mod. SBE19). These data were processed and provided by the Physical Oceanography team of IOUSP.

Larvae were examined under a stereomicroscope and the sciaenid identification was based on: Hildebrand and Cable (1934), Sinque (1980), Able and Fahay (1998), Chao (2002), Ditty et al. (2006), Fahay (2007) and Itagaki et al. (2007). Photographs taken with a digital camera connected to a stereomicroscope and the software ImageJ were used to measure the fish larvae.

Data Analysis

Analysis of variance (one-way ANOVA) was used to test the significant differences between the

abiotic data of water temperature and water salinity, both at 5 m, which were calculated monthly and by estuarine area. We chose to study the abiotic data at 5 m depth to avoid any influence of the air temperature at the sea surface.

Guille (1970) was used as the basis for the calculation of the frequency of larval occurrence (FO). For the abundance estimative, the number of larvae collected at each station was transformed into the number of larvae per m² of sea surface, in accordance with the following equation: $Y = (D \cdot X) / V$, where Y = number of larvae per square meter, D = maximum sampling depth (m), X = number of larvae collected, and V = volume of water filtered (m³). The mean abundance was calculated for all the sampling stations, not only for the positive ones. Filtered volume was calculated as $V = a \cdot n \cdot c$, where a = area of net mouth (0.2827 m²), n = number of rotations of the flowmeter and c = calibration factor of the flowmeter.

The influence of each abiotic variable on the Sciaenidae species was assessed by Canonic Correspondence Analysis (CCA) (LEGENDRE; LEGENDRE, 1998) and the distribution of the species in relation to the significant abiotic variables was determined by CANOCO. The Monte Carlo Permutation test was carried out to judge the significance of the selected variables. Rare species,

those with less than 5 % of frequency of occurrence, were eliminated from the biotic matrix. The abiotic data, after a first analysis using a single matrix, were divided into three matrices: environmental (temperature and salinity, both at 5 m depth), spatial (north shelf, south shelf, bay, port, channel and local depth) and temporal (months sampled). In all analyses the biological data (number of individuals per m^2) were transformed by $\log(x+1)$, and low weight was given to rare species. The percentage of explanation of each abiotic variable was calculated in accordance with Borcard et al. (1992).

One-way analysis of similarity (ANOSIM) was performed to determine the significance of spatial trends in the structure of the Sciaenidae assemblage. The contribution of each species to the dissimilarity between different groups' species was estimated using analysis of similarity percentages (SIMPER). Both analyses were performed using the software package PRIMER 6 (Plymouth Routines Multivariate Ecological Research) (CLARKE, 1993). Rare taxa, with an occurrence frequency of less than 5%, were excluded from the analyses. Numerical density data expressed as number of individuals per m^2 were transformed using the $\log(x+1)$ function to enhance the contribution of the less abundant species. The analyses were undertaken on the basis of the Bray–Curtis similarity index (BRAY; CURTIS, 1957). R values approximating to 1 indicate great differences between the groups analyzed, while R values approximating to zero indicate low differences.

RESULTS

Hydrographic Conditions

Santos estuarine system. Water temperature at 5 m varied over the different estuarine areas (ANOVA, $p < 0.05$) and over the sampling period (ANOVA, $p < 0.05$), following a seasonal pattern. The lowest values occurred between July and October, and the highest ones from November 2004 to March 2005, and in December 2005 (Fig. 2). The maximum temperature was 28.05°C in March 2005 in the Channel and the minimum was 20.26°C in October 2005 in the Bay.

Water salinity at 5 m also varied over the different estuarine areas (ANOVA, $p < 0.05$) and over the sampling period (ANOVA, $p < 0.05$). Differently from water temperature, water salinity showed no seasonal pattern. Highest salinity values were measured in the Bay, followed by the Port and then by the Channel (Fig. 3). The maximum salinity was 35.72 in November 2004 in the Bay and the minimum was 18.42 in April 2005 in the Channel.




Fig. 2. Monthly variation of mean water temperature at 5 m depth measured in the Santos estuarine system from November 2004 to December 2005.




Fig. 3. Monthly variation of mean water salinity at 5 m depth measured in the Santos estuarine system from November 2004 to December 2005.

Adjacent continental shelf. In September 2005 the water temperature was almost homogeneous throughout the continental shelf (Fig. 4). At the surface a range from 19.59°C to 20.91°C was observed, averaging 20.38°C ($sd = 0.37$). At the deeper layer (25 m) the temperature range was slightly lower, from 19.00°C to 20.95°C (mean = 19.95°C , $sd = 0.51$). In March 2006 the thermal structure was quite different from that observed in September 2005, with

warmer surface seawater ranging from 25.17°C to 28.01°C (mean = 26.96°C, sd = 0.68), but lower temperatures were observed for the 25 m layer, with a minimum of 16.98°C and maximum of 23.60°C (mean = 20.34°C, sd = 1.79). This minimum temperature value, associated with salinity values around 35.7, indicates the presence of South Atlantic Central Water (SACW) in the deeper layer.

In September 2005 the surface water salinity ranged from 30.92 to 34.38 (mean = 32.91, sd = 1.03). At the 25 m layer higher salinity values were observed, ranging from 33.37 to 34.53 (mean = 33.82, sd = 0.33). In March 2006 the range of salinity was from 33.90 to 35.22 (mean = 34.63, sd = 0.41) at the surface, and from 35.63 to 36.10 (mean = 35.88, sd = 0.12) at the 25 m layer (Fig. 5).




Fig. 3. Horizontal profiles of water temperature on the surface, at 5 m and at 25 m depth, measured in September 2005 and March 2006 between Peruíbe and the São Sebastião island, from the coast to the 50 m isobath. The dotted line indicates the boundary of the South Atlantic Central Water (SACW), according to Castro et al. (2006).




Fig. 4. Horizontal profiles of water salinity on the surface, at 5 m and at 25 m depth, measured in September 2005 and March 2006 between Peruibe and the São Sebastião island, from the coast to the 50 m isobath.

Fish Larvae Distribution and Abundance

Santos estuarine system. A total of 11,811 fish larvae were collected in the estuarine system, and the sciaenids accounted for 12.71% of the total number. Most of the sciaenids were caught during November and December 2004. The mean abundance of sciaenid larvae ranged from 0.12 larvae.m⁻² (sd = 0.11) in June 2005 to 4.28 larvae.m⁻² (sd = 1.99) in November 2004. The minimum FO was 50% in August 2005, and the maximum 100% in November 2004.

The following sciaenids were identified in the estuarine system: *Bairdiella ronchus*, *Cynoscion spp.*, *Isopisthus parvipinnis*, *Macrodon atricauda*, *Menticirrhus spp.*, *Micropogonias furnieri*, *Nebris microps* Cuvier, 1830, *Ophioscion punctatissimus* Meek & Hildebrand, 1925, *Paralichthys brasiliensis* (Steindachner, 1875), *Stellifer rastrifer* and *Stellifer spp.* Of these, *S. rastrifer* was the most abundant (mean = 0.36 larvae.m⁻²; sd = 0.35) and with higher FO (91.7%) (Table 3).

Adjacent continental shelf. A total of 17,572 fish larvae were collected on the continental shelf, 6.72% of them being sciaenids. Larvae of this family were widely distributed during both cruises, with FO greater than 85%, but were at their most abundant in March 2006. In this month the abundance varied from 0.27 larvae.m⁻² to 35.73 larvae.m⁻² (mean = 6.47 larvae.m⁻², sd = 8.36). In September 2005 the range was from 0.20 larvae.m⁻² to 14.75 larvae.m⁻² (mean= 2.37 larvae.m⁻², sd= 3.33).

Larvae of the following sciaenids were present in the shelf area: *Bairdiella ronchus*, *Ctenosciaena gracilicirrhus* (Metzelaar, 1919), *Cynoscion spp.*, *Isopisthus parvipinnis*, *Macrodon atricauda*, *Menticirrhus spp.*, *Nebris microps*, *Stellifer rastrifer* and *Stellifer spp.* *Cynoscion spp.* were the most abundant during both cruises, with a mean of 0.49 larvae.m⁻² (sd = 1.81) and FO of 22.2% in September 2005, and 1.65 larvae.m⁻² (sd = 3.44) and FO of 40.5% in March 2006. *Menticirrhus spp.* (mean= 0.12 larvae.m⁻²; sd = 0.27) and *Stellifer spp.* (mean=0.40 larvae.m⁻²; sd = 1.48) were the second in abundance, but respectively in September 2005 and in

March 2006, both with the FO of approximately 22% (Table 4).

The Main Taxa Identified

Bairdiella ronchus. In the estuarine system the highest abundance of *B. ronchus* occurred in November 2004 (0.10 larvae.m⁻²), when larvae were widespread throughout the area (Bay, Port and Channel). In January and December 2005 they occurred only in the Bertioga channel and in February 2005 only in the Santos bay. The *B. ronchus* larvae found in the estuarine system accounted for only 0.93% of the sciaenids present there. On the adjacent continental shelf, the abundance of *B. ronchus* was highest in March 2006 (0.08 larvae.m⁻²), their occurrence being restricted, however, to the southern part. In September 2005 their occurrence was restricted to only one station in the north. In the shelf area this species accounted for 1.10% of all the sciaenids. In the estuarine system the length frequency ranged from 2.29 mm to 4.28 mm, and on the shelf from 2.63 mm to 4.66 mm.

Table 3. Size range, abundance (larvae.m⁻²), frequency of occurrence (%) and absolute number (N) of Sciaenidae larvae in the Santos estuarine system, Brazil, from November 2004 to December 2005.

| Sciaenidae | Size | | Abundance | | FO | N |
|----------------------------------|------|------|-----------|-------|------|-----|
| | Min | Max | Mean | SD | | |
| <i>Bairdiella ronchus</i> | 2.9 | 4.3 | 0.011 | 0.029 | 33.3 | 14 |
| <i>Isopisthus parvipinnis</i> | 3.9 | 12.2 | 0.063 | 0.084 | 66.7 | 66 |
| <i>Macrodon atricauda</i> | 4.2 | 13.4 | 0.027 | 0.051 | 50.0 | 23 |
| <i>Micropogonias furnieri</i> | 3 | 13.7 | 0.111 | 0.146 | 83.3 | 123 |
| <i>Nebris microps</i> | 2.6 | 3.2 | 0.003 | 0.011 | 8.3 | 2 |
| <i>Ophioscion punctatissimus</i> | 4.5 | 4.5 | 0.001 | 0.005 | 8.3 | 1 |
| <i>Paralichthys brasiliensis</i> | 4.5 | 7.3 | 0.004 | 0.007 | 25.0 | 3 |
| <i>Stellifer rastrifer</i> | 2.6 | 8.3 | 0.361 | 0.351 | 91.7 | 360 |
| <i>Cynoscion spp.</i> | 2.3 | 8.3 | 0.030 | 0.039 | 66.7 | 35 |
| <i>Menticirrhus spp.</i> | 2.4 | 6.6 | 0.057 | 0.089 | 66.7 | 67 |
| <i>Stellifer spp.</i> | 2.5 | 11.9 | 0.212 | 0.401 | 50.0 | 256 |
| Non-identified | | | | | | 551 |
| Total | | | | | | 15 |

Table 4. Size range, abundance (larvae.m⁻²), frequency of occurrence (FO %) and absolute number (N) of Sciaenidae larvae on the continental shelf adjacent to the Santos-São Vicente estuarine system, Brazil, in September 2005 and March 2006.

| Sciaenidae | September 2005 | | | | | | March 2006 | | | | | |
|------------------------------------|----------------|-----|-----------|------|------|-----|------------|-----|-----------|------|------|-----|
| | Size | | Abundance | | FO | N | Size | | Abundance | | FO | N |
| | Min | Max | Mean | SD | | | Min | Max | Mean | SD | | |
| <i>Bairdiella ronchus</i> | 3.0 | 4.7 | 0.03 | 0.16 | 2.8 | 3 | 2.6 | 4.1 | 0.08 | 0.22 | 13.5 | 10 |
| <i>Ctenosciaena gracilicirrhus</i> | - | - | 0.01 | 0.04 | 2.8 | 1 | - | - | - | - | - | - |
| <i>Isopisthus parvipinnis</i> | 3.8 | 5.0 | 0.02 | 0.09 | 5.6 | 2 | - | - | - | - | - | - |
| <i>Macrodon atricauda</i> | 3.6 | 4.0 | 0.01 | 0.05 | 5.6 | 2 | - | - | - | - | - | - |
| <i>Nebris microps</i> | 5.5 | 7.7 | 0.03 | 0.09 | 8.3 | 3 | - | - | - | - | - | - |
| <i>Stellifer rastrifer</i> | - | - | - | - | - | - | 2.6 | 6.2 | 0.08 | 0.48 | 2.7 | 13 |
| <i>Cynoscion spp.</i> | 2.2 | 4.2 | 0.49 | 1.81 | 22.2 | 56 | 1.8 | 6.1 | 1.65 | 3.44 | 40.5 | 224 |
| <i>Menticirrhus spp.</i> | 3.4 | 1.4 | 0.12 | 0.27 | 22.2 | 13 | 2.3 | 4.5 | 0.21 | 0.39 | 24.3 | 28 |
| <i>Stellifer spp.</i> | 3.5 | 9.8 | 0.06 | 0.20 | 8.3 | 10 | 2.2 | 6.9 | 0.40 | 1.48 | 21.6 | 51 |
| Non-identified | | | | | | 187 | | | | | | 573 |
| Total | | | | | | 277 | | | | | | 899 |

Isopisthus parvipinnis. The larvae of this species were almost exclusively estuarine, having their highest abundance in November 2004 (0.14 larvae.m⁻²), January 2005 (0.24 larvae.m⁻²) and March 2005 (0.19 larvae.m⁻²). They occurred in the three estuarine areas (Bay, Port and Channel). Their body length ranged from 3.85 mm to 12.20 mm in the estuary. In the shelf area only three larvae were collected in September/2005, ranging from 5.07 mm to 7.24 mm.

Macrodon atricauda. Larvae of this species occurred in the estuarine system and on the adjacent continental shelf, but among the estuarine sites *M. atricauda* larvae were absent from the Bertioga channel. In the estuarine system the abundance was the highest in July 2005 (0.16 larvae.m⁻²). In the shelf area this species was rare, with only two specimens collected in September 2005, representing a mean abundance of 0.01 larvae.m⁻². In the estuary the body length of *M. atricauda* ranged from 4.18 mm to 13.41 mm. The two larvae collected on the shelf were of 3.55 mm and 4.03 mm. *Micropogonias furnieri*. This species was absent from the adjacent continental shelf. On the other hand, its occurrence was observed throughout the estuarine system, mainly in October 2005 (mean abundance = 0.43 larvae.m⁻²). In February 2005 this species was also abundant (mean = 0.39 larvae.m⁻²), though it occurred only in the Bertioga channel. *M. furnieri* accounted for 8.19% of all the sciaenid larvae in the estuarine system. Body length ranged from 2.98 mm to 13.68 mm.

Stellifer rastrifer. This species occurred mostly in the estuarine system, the highest abundance (1.05 larvae.m⁻²) being observed in January 2005. High abundance was also observed in November 2004, December 2004, and February 2005. *S. rastrifer* was one of most common sciaenids in the estuarine area, accounting for 23.99% of all the Sciaenidae larvae. No significant differences among the three estuarine areas were observed (P<0.0001). On the adjacent shelf area the larval *S. rastrifer* occurred only in March 2006, with low abundance (mean = 0.07 larvae.m⁻²). In the estuary the length class ranged from 2.54 mm to 8.64 mm.

Cynoscion spp. This genus was present in the estuarine and the shelf areas. In the estuarine system *Cynoscion spp.* abundance was relatively low, the highest abundance occurring in November 2004 (0.12 larvae.m⁻²). In the estuary the *Cynoscion spp.* represented 2.33% of sciaenids. On the adjacent continental shelf this genus was more common, with wider distribution and higher abundance observed in March 2006 (mean = 1.65 larvae.m⁻²). In September 2005 the mean abundance was 0.49 larvae.m⁻². These

larvae accounted for 23.70% of all the sciaenids in the shelf area.

Menticirrhus spp. This genus occurred in both estuarine and shelf areas. It contributed with a total of 4.46% of all the sciaenids in the estuary and 3.47% of those on the shelf. In the estuarine system the highest abundance was observed in November 2004 (mean = 0.31 larvae.m⁻²) and in December 2004 (mean = 0.13 larvae.m⁻²). On the adjacent continental shelf *Menticirrhus spp.* was more abundant (mean = 0.21 larvae.m⁻²) with wide distribution in March 2006.

Stellifer spp. In the estuarine system larvae of this genus were more abundant in November 2004 (1.92 larvae.m⁻²) and December 2004 (1.49 larvae.m⁻²). They were present in all the estuarine areas and contributed with 17.05% of the taxonomic composition of Sciaenidae. On the adjacent continental shelf, the genus presented its highest abundance and widest distribution in March 2006. It represented 5.16% of the sciaenids in the shelf area.

Abiotic influence on the Sciaenidae larvae

The abiotic variables explained 42.21 % of the variability of the biological data. The first two axes were responsible for 66.30% of biological data variation. Axis 1 was negatively correlated with local depth, South Shelf, North Shelf and salinity. Axis 2 was positively correlated with the months of July and October 2005 and negatively correlated with January 2005. The species-environment correlation presented high values on the first (0.89) and second (0.79) axes. The combined sum of canonical eigenvalues (1.675) equaled 42.2% of that for all the eigenvalues (3.968), showing the effect of an adequate building of an environmental relationship into the CCA model (Table 5). The canonic axes were significantly different by the Monte Carlo permutation test (p= 0.0001). Figure 6 represents the distribution of the species in relation to the significant abiotic variables.

Analyzing the environmental matrix, only salinity was significant by the Monte Carlos Permutation test, with an explanation of 7.18 % of the biological data variation (sum of all canonic eigenvalues = 0.28). *S. rastrifer* and *I. parvipinnis* were negatively correlated with salinity.

The temporal data were those with the highest influence on the biological data variation (percentage of explanation = 34.30%; sum of all canonic eigenvalues = 1.36). *I. parvipinnis* was positively correlated with January 2005. *M. furnieri* and *M. atricauda* were positively correlated with July and October 2005. *Cynoscion spp.* and *S. rastrifer* were more abundant during months of high temperature, during the rainy period.




Fig. 6. Ordination diagram (biplot) from CCA including Sciaenidae species and significant environmental variables (represented by vectors) (BARO = *Bairdiella ronchus*; CYPSP = *Cynoscion spp.*; ISPA = *Isopisthus parvipinnis*; MAAT = *Macrodon atricauda*; MESP = *Menticirrhus spp.*; MIFU = *Micropogonias furnieri*; STRA = *Stellifer rastrifer*; STSP = *Stellifer spp.*).

Table 5. Summary of CCA performed on the abundance of eight most numerous Sciaenidae larval species, in the Santos estuarine system, from November 2004 to December 2005, and on the continental shelf adjacent to the Santos-São Vicente estuarine system, Brazil, in September 2005 and March 2006.

| | Axes | | | |
|--|-------|-------|-------|-------|
| | 1 | 2 | 3 | 4 |
| Summary statistics for ordination axes | | | | |
| Eigenvalues | 0.675 | 0.437 | 0.311 | 0.118 |
| Species-environment correlations | 0.892 | 0.787 | 0.674 | 0.483 |
| Cumulative percentage variance: | | | | |
| of species data | 17.0 | 28.0 | 35.8 | 38.8 |
| of species-environment relation | 40.3 | 66.3 | 84.9 | 91.9 |
| Sum of all eigenvalues | | | | 3.968 |
| Sum of all canonical eigenvalues | | | | |

The spatial data explained 17.97 % of the biological data variation (sum of all canonic eigenvalues = 0.71), only the influence of south and north shelf and local depth being statistically significant. *B. ronchus*, *Menticirrhus spp.* and *Cynoscium* were positively correlated with the south and north shelf.

One-way analysis of similarity (ANOSIM) showed significant differences between the continental

shelf and the estuarine areas. Within the estuarine areas, only Bay and Channel showed significant differences. The differences were more evident between the more distant regions, as exemplified by those between Shelf and Channel (R=0.44) (Table 6). These results showed a gradual spatial variation of the Sciaenid larvae between the continental shelf and the inner estuarine area.

Table 6. Similarity analysis of Sciaenidae larval groups between the different sampling areas: Shelf, Bay, Port and Channel.

| | R | p |
|-----------------|--------|-------|
| Shelf - Bay | 0.273 | 0.001 |
| Shelf - Port | 0.402 | 0.001 |
| Shelf - Channel | 0.440 | 0.001 |
| Bay - Port | -0.006 | 0.526 |
| Bay - Channel | 0.130 | 0.005 |
| Port - Channel | 0.113 | 6.900 |

According to SIMPER analysis, the species that most contributed to the formation of the Shelf group were: *Cynoscion spp.* (64.44%), *Menticirrhus spp.* (20.15%) and *Stellifer spp.* (10.33%); to the Bay group: *S. rastrifer* (43.71%), *Stellifer spp.* (17.97%), *M. furnieri* (10.54%), *Menticirrhus spp.* (10.29%) and *I. parvipinnis* (7.96%), and to the Port group: *S. rastrifer* (72.61%), *M. atricauda* (10.78%) and *Menticirrhus spp.* (7.53%). Finally, the Channel group consisted mainly of: *S. rastrifer* (78.09%), *Stellifer spp.* (10.16%) and *M. furnieri* (6.03%).

DISCUSSION

Although Sciaenidae do not constitute a dominant family, they can be recognized as an important component of the ichthyoplankton in the present study area. As indicated by our results, the sciaenid accounted for 12% and 15% of total fish larvae in the estuary and on the adjacent continental shelf, respectively. Some similar results have been observed in various estuarine and coastal areas around the world. For instance, Rakocinski et al. (1996), by analyzing the ichthyoplankton assemblage structure in Mississippi Sound, described the sciaenids as being among the seven major groups. According to Katsuragawa and Dias (1997), in a region very close to that of the present study, the inner shelf of São Sebastião, the Sciaenidae constituted the 8th family among all the larvae identified, but the second among the demersal group.

The literature indicates that the sciaenids, mainly during their early life stages, prefer protected environments, such as those provided by mangrove swamps, estuaries and bays (CHAO, 1978). The

notable presence of Sciaenidae in estuarine and coastal ecosystems has long been known, mainly as a result of studies of adult specimens. Several studies may be cited, especially relating to the southeastern Brazilian coast, e.g. Chaves and Corrêa (1998), Chaves and Bouchereau (2000), Pessanha et al. (2000), Araújo et al. (2002) and Spach et al. (2004). Concerning the area investigated by the present study, Paiva Filho and Schmiegelow (1986) reported the Sciaenidae as a family of higher biomass and number of specimens, according to their estimates based on their bottom trawling method in the Santos bay.

A feature of the sciaenids collected in the present study was the large number of small larvae, which caused us some difficulty in identifying the specific levels. However, nine species (*Bairdiella ronchus*, *Ctenosciaena gracilicirrhus*, *Isopisthus parvipinnis*, *Macrodon atricauda*, *Micropogonias furnieri*, *Nebrius microps*, *Ophioscion punctatissimus*, *Paralonchurus brasiliensis* and *Stellifer rastrifer*) and three genera (*Cynoscion spp.*, *Menticirrhus spp.* and *Stellifer spp.*) were classified. As compared with the São Sebastião region (Katsuragawa; Dias, 1997), it may be observed that the taxonomic composition of the sciaenids in the two areas differed by virtue of the presence of five species (*Cynoscion leiarchus*, *Larimus breviceps*, *Macrodon ancylodon*, *Menticirrhus americanus* and *P. brasiliensis*) and the absence of *M. atricauda* in São Sebastião. Unfortunately, due to their small size, the *Menticirrhus* and *Cynoscion* genera of the present study could not be refined to species level, but probably *M. americanus*, the main species and widely distributed in São Sebastião (KATSURAGAWA; DIAS, 1997), was also present in Santos. The species *S. rastrifer*, *I. parvipinnis*, and *M. atricauda* were also reported with great frequency and abundance among the adult specimens in the Santos region (GIANNINI; PAIVA FILHO, 1990; SCHMIDT; DIAS, 2012).

The considerable presence of early larval stages suggests an effective use of the Santos estuarine system as a spawning or nursery ground by sciaenid species. On the other hand, the shelf area was in general poor in terms of abundance, and among the species identified, the larval *M. furnieri* was absent. Our results concerning the role of the estuarine system are coherent with the presence of adult sciaenids there, e.g. Chaves (1995) reported that the reproduction of *B. ronchus* occurs inside the Guaratuba bay and that, thereafter, the adults disperse in the coastal area. *M. furnieri* is known as a species highly dependent on semi-enclosed systems such as bays and estuaries (GIANNINI; PAIVA FILHO, 1990; VIEIRA; CASTELLO, 1997; COSTA; ARAÚJO, 2003), and

juveniles and pre-recruits are not present in the shelf area (CASTELLO, 1986; VAZZOLER, 1991). According to Vieira and Castello (1997), living the first year of life inside the Patos lagoon may help the early stages to avoid predation.

Many authors refer to the estuaries as nursery ground for fish (e.g. GODEFROID et al., 1999; COSTA; ARAÚJO, 2003). Lowe-McConnell (1987) included the sciaenids in a group of migrant marine fishes which breed in the sea and whose young enter the estuaries to feed and shelter until their gonads start to mature. Able and Fahay (1998), in a study on the sciaenid species found in the Middle Atlantic Bight, reported that spawning areas are diversified, occurring close to the coast for species such as *Cynoscion regalis*, within harbors, estuaries and sounds for *Bairdiella chrysoura*, and on the continental shelf for *Leiostomus xanthurus* and *Micropogonias undulatus*, all of them, however, being found in estuaries during their first year of life. The sciaenids in the study area seem to follow these trends, with spawning occurring throughout the shelf area, as evidenced by our results for many species, whereas the higher abundance of larvae in the estuarine system may be a clue to spawning activity within this area. At least four species (*Stellifer brasiliensis*, *S. rastrifer*, *I. parvipinnis* and *P. brasiliensis*) are known to complete their reproductive cycle inside the Santos bay.

Temporal change, the abiotic variable that presented the highest percentage of explanation of the occurrence and abundance of sciaenid larvae, is also reflected in their patterns of reproduction and intensity of spawning. Chaves (1995) found that *B. ronchus* reproduces during spring and summer, the period of larval peak abundance for this species in the present study area. Based on the reproductive biology studies of *S. rastrifer* from Guanabara bay, Chaves and Vendel (1997) observed more intense reproductive activity during the winter and especially in spring. In the present study the peak of mean abundance of larval *S. rastrifer* was observed to occur in January, though this species was well represented throughout the year. This result corroborates the study of Sinque (1980), which found large numbers of *S. rastrifer* larvae throughout the year in the Cananéia-Iguape estuarine system. This larval occurrence and abundance pattern also matches the continuous spawning behavior of adults observed by Schmidt (2011) evidenced by the presence of spawning females in samples caught in at least eight months of the year. Another species with variable abundance along the year was *M. furnieri*, with two peaks (February and October 2005). This result is coherent with the study on reproductive

biology conducted by Isaac-Nahum and Vazzoler (1987) who reported a long spawning period during the four seasons.

Historically the Santos estuarine system has been affected by anthropic activities. However, the three estuarine areas investigated in this study present distinct features concerning the source and relative level of the impacts caused. The Santos port area is supposed to have been the most heavily impacted by the several activities related to industry, port and urbanization, and one of the thoughts entertained at the beginning of the ECOSAN project was that, of the areas involved, the Bertioga channel would be the least severely impacted. Our results suggest that the differences among areas have not affected the abundance of sciaenid larvae, as indicated by the ANOSIM.

It is not easy to identify the main environmental factors that influence the distribution and abundance of fish (larvae or adults) in an estuarine system, because in estuaries the lives of organisms are controlled by the interaction of many different abiotic and biotic factors (ALLEN, 1982; LOWE-MCCONNELL, 1987; SPACH et al., 2004). Estuaries are characterized by varying salinity, temperature, turbidity, muddy bottoms, strong currents and freshwater flowing out over the denser more saline water (LOWE-MCCONNELL, 1987). The biotic factors include availability of prey, competitors and predators (ALLEN, 1982). In the present study it has not been possible to evaluate the influence of biotic factors, but the CCA analysis indicated that the abiotic variables measured in the present study showed a high percentage of explanation of the variability of larval sciaenids (42.21 %). This larval variation was explained mainly in terms of the different months sampled (temporal influence), local depth, the south and the north part of the shelf area (spatial influence) and salinity. In this case, the variation in the abundance of larvae in the estuarine system seems, in general, to follow the seasonal patterns observed in the marine coastal waters along the southeastern Brazil coast, with higher abundance during spring and summer.

Although the influence of the temperature on the sciaenid larvae species variation was not significant in this study, it is known that temperature is one of the most important physical properties of the marine environment as it exerts an influence on many physical, chemical, geochemical and biological events (LALLI; PARSONS, 1995). We believe that temperature has been best represented by the temporal data analysis.

In the Santos estuarine system the average temperature is always higher than 20°C, the lowest average being 22.60°C, in the winter, and the highest average 26.09°C, in summer. The large difference between the mean temperatures during spring 2004 (25.28°C) and spring 2005 (22.60°C) may suggest an interannual oscillation in the thermic structure of this area. However, the seasonal range of temperature (from the minimum of 20.77°C to the maximum of 28.50°C) in the present study is far smaller than that observed in other areas, e.g. 9.0°C - 33.5°C in the Mississippi Sound (Rakocinski et al., 1996) and 0°C - 28°C in the Chesapeake Bay (Able; Fahay, 1998). The relatively small variation in temperature perhaps contributed to the all-year-round patterns of distribution and abundance of larval sciaenids in the Santos estuarine system with a tendency to greater abundance in the warmer months.

In the shelf area the thermal dynamics were very different, varying from almost homogeneous and colder in September 2005 (19.00°C - 20.95°C) to a stratified March 2006 scenario with warm water (25.17°C - 28.01°C) at the surface but cold water (minimum of 16.98°C) at the 25 m layer, suggesting the presence of SACW. However, according to the ECOSAN reports (Pires-Vanin et al., 2008), this intrusion of SACW did not reach the euphotic zone, and the new primary production was not significant. Pires-Vanin et al. (2008) concluded that on the shelf off Santos the contribution of SACW to the fertilization of the surface water is not as effective as it is off the Ubatuba and São Sebastião coast, perhaps due to the greater width of the continental shelf in the study region. In this case, the role of the estuarine system in providing organic matter for, at least, the neighboring coast must be of great importance in promoting primary production.

The results of the present study show that the abundance in the shelf area was considerably higher in March 2006 both for larval sciaenids and for total fish larvae collected, indicating a seasonal variation with an increase during the warm season, similar to that observed on other parts of the southeastern coast by several authors, e.g. Matsuura et al. (1992), Katsuragawa et al. (1993) and Lopes et al. (2006).

ACKNOWLEDGEMENTS

We thank FAPESP (Fundação de Amparo à Pesquisa do Estado de São Paulo) for providing the financial support for the ECOSAN project. We are grateful to the captains and crews of R.V. "Prof. W. Besnard" and the boats for their cooperation and assistance in the sampling described in this study. We would also thank Drs. Luiz Bruner de Miranda and

Belmiro Mendes de Castro for providing the CTD data used in this project.

REFERENCES

- ABLE, K. W.; FAHAY, M. P. **The first year in the life of estuarine fishes in the Middle Atlantic Bight**. New Brunswick: Rutgers University Press, 1998. 342 p.
- ALLEN, J. R. L. Simple models for the shape and symmetry of tidal sand waves: (3) dynamically stable asymmetrical equilibrium forms without flow separation. *Mar. Geol.*, v. 48, n. 3/4, p. 321-336, 1982.
- ARAÚJO, F. G.; AZEVEDO, M. C. C.; SILVA, M. A.; PESSANHA, A. L. M.; GOMES, I. D.; CRUZ-FILHO, A. G. Environmental influences on the demersal fish assemblages in the Sepetiba Bay, Brazil. *Estuaries*, v. 25, n. 3, p. 441-450, 2002.
- BORCARD, D.; LEGENDRE, P.; DRAPEAU, P. Partialling out the spatial component of ecological variation. *Ecology*, v. 73, n. 3, p. 1045-1055, 1992.
- BRAY, J. R.; CURTIS, J. T. An ordination of the upland forest communities of Southern Wisconsin. *Ecol. Monogr.*, v. 27, n. 4, p. 325-349, 1957.
- BRUNO, M. A.; MUELBERT, J. H. Padrões temporais de distribuição da abundância de ovos e larvas de *Brevoortia pectinata*, *Lycengraulis grossidens* e *Micropogonias furnieri* para o estuário da Lagoa dos Patos, Rio Grande - RS. In: SIMPÓSIO BRASILEIRO DE OCEANOGRAFIA, 1., São Paulo, 2002. **Resumo**. São Paulo: IO/USP, 2002. 1 CDROOM.
- BRUNO, M. A.; MUELBERT, J. H. Distribuição espacial e variações temporais da abundância de ovos e larvas de *Micropogonias furnieri*, no estuário da Lagoa dos Patos: registros históricos e forçantes ambientais. *Atlântica*, v. 31, n. 1, p. 51-68, 2009.
- CASATTI, L.; MENEZES, N. A. Sciaenidae. In: MENEZES, N. A.; BUCKUP, P. A.; FIGUEIREDO, J. L.; MOURA, R. L. (Eds.). **Catálogo das espécies de peixes marinhos do Brasil**. São Paulo: Museu de Zoologia da Universidade de São Paulo, 2003, 160 p.
- CASTELLO, J. P. Distribución, crecimiento y maduración sexual de la corvina juvenil (*Micropogonias furnieri*) en el estuario de la "Lagoa dos Patos", Brasil. *Physis*, v. 44, n. 106, p. 21-36, 1986.
- CASTRO, B. M.; LORENZETTI, J. A.; MIRANDA, L. B.; SILVEIRA, I. C. A. Estrutura Termohalina e circulação na região entre Cabo de São Tomé (RJ) e o Chuí (RS). In: ROSSI-WONGTSCHOWSKI, C. L. D. B.; MADUREIRA, L. S. P. **O ambiente oceanográfico da Plataforma Continental e do Talude na região sudeste-sul do Brasil**. São Paulo: EDUSP, 2006. p. 11-120.
- CETESB. **Sistema estuarino de Santos e São Vicente**. São Paulo: Companhia de Tecnologia de Saneamento Ambiental, 2001. 183p. (Relatório técnico).
- CHAO, L. N. **A basis for classifying western Atlantic Sciaenidae (Teleostei: Perciformes)**. Washington: U.S. Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, 1978. 64 p. (NOAA Technical Report NMFS Circular, 415). Available on: <http://spo.nmfs.noaa.gov/Circulars/CIRC415.pdf>. Accessed in: 07/15/2013.
- CHAO, L. N. Sciaenidae. Croakers (drums). In: CARPENTER, K. E. (Ed.). **The living marine resources of the Western Central Atlantic**. Rome: Food and Agriculture Organization of the United Nations, 2002. p. 1583-1653. (FAO Species Identification Guide for Fishery Purposes) (Special Publications - American Society of Ichthyologists and Herpetologists, n. 5).
- CHAVES, P. T. C. Atividade reprodutiva de *Bairdiella ronchus* (Cuvier) (Pisces, Sciaenidae) na Baía de Guaratuba, Paraná, Brasil. *Rev. Bras. Zool.*, v. 12, n. 4, p. 759-766, 1995.
- CHAVES, P. T. C.; VENDEL, A. L. Reprodução de *Stellifer rastrifer* (Jordan) (Teleostei, Sciaenidae) na Baía de Guaratuba, Paraná, Brasil. *Rev. Bras. Zool.*, v. 14, n. 1, p. 81-89, 1997.
- CHAVES, P. T. C.; CÔRREA, M. F. M. Composição ictiofaunística da área de manguezal da Baía de Guaratuba, Paraná, Brasil. *Rev. Bras. Zool.*, v. 15, n. 1, p. 195-202, 1998.
- CHAVES, P.; BOUCHEREAU, J. L. Use of mangrove habitat for reproductive activity by the fish assemblage in the Guaratuba Bay, Brazil. *Oceanol. Acta*, v. 23, n. 3, p. 273-280, 2000.
- CLARKE, K. R. Non-parametric multivariate analyses of changes in community structure. *Aust. J. Ecol.*, v. 18, n. 1, p. 117-143, 1993.
- COSTA, M. R.; ARAÚJO, F. G. Use of a tropical bay in Southeastern Brazil by juvenile and subadult *Micropogonias furnieri* (Perciformes, Sciaenidae). *ICES J. Mar. Sci.*, v. 60, n. 2, p. 268-277, 2003.
- DENMAN, K. L.; POWELL, T. M. Effects of physical processes on planktonic ecosystems in the coastal ocean. *Oceanogr. Mar. Biol.*, v. 22, p. 125-168, 1984.
- DITTY, J. G.; SHAW, R. F.; FAROQI, T. Sciaenidae. In: RICHARDS, W. J. (Ed.). **Early stages of atlantic fishes: an identification guide for the Western Central North Atlantic**. Boca Raton: CRC Press, Taylor & Francis, 2006. p. 1669-1723. (CRC marine biology series).
- FAHAY, M. P. **Early stages of fishes in the Western North Atlantic Ocean (Davis Strait, Southern Greenland and Flemish Cap to Cape Hatteras)**. Nova Scotia: Northwest Atlantic Fisheries Organization, 2007. 1696 p.
- GIANNINI, R.; PAIVA FILHO, A. M. Os Sciaenidae (Teleostei: Perciformes) da baía de Santos (SP), Brasil. *Bol. Inst. Oceanogr.*, v. 38, n. 1, p. 69-86, 1990.
- GODEFROID, R. S.; HOFSTAETTER, M.; SPACH, H. L. Larval fish in the surf zone of Pontal do Sul Beach, Pontal do Paraná, Paraná, Brazil. *Rev. Bras. Zool.*, v. 16, n. 4, p. 1005-1011, 1999.
- GODEFROID, R. S.; SANTOS, C.; HOFSTAETTER, M.; SPACH, H. L. Occurrence of larvae and juveniles of *Eucinostomus aregenteus*, *Eucinostomus gula*, *Menticirrhus americanus*, *Menticirrhus littoralis*, *Umbrina coroides* and *Micropogonias furnieri* at Pontal do Sul Beach, Paraná. *Braz. Arch. Biol. Tecnol.*, v. 44, n. 4, p. 411-418, 2001.

- GUILLE, A. Benthic bionomy of continental shelf of the French Catalane coast. II: Benthic communities of the macrofauna. **Vie Milieu**, v. 21, n. 8, p. 149-280, 1970.
- HANNA, S. S. Institutions for marine ecosystems: economic incentives and fishery management. **Ecol. Appl.**, v. 8, n. 1, suppl., p. S170-S174, 1998.
- HILDEBRAND, S. F.; CABLE, L. E. Reproduction and development of whiting or kingfishes, drums, spot, croaker, and weakfishes or sea trouts, family Sciaenidae of the Atlantic coast of the United States. **Bull. U. S. Bur. Fish.**, v. 48, n. 16, p. 41-117, 1934.
- ISAAC-NAHUM, V. J.; VAZZOLER, A. E. A. M. Biologia reprodutiva de *Micropogonias furnieri* (Desmarest, 1823) (Teleostei, Sciaenidae). 2. Relação gonadossomática, comprimento e peso dos ovários como indicadores do período de desova. **Bol. Inst. Oceanogr.**, v. 35, n. 2, p. 123-134, 1987.
- ITAGAKI, M. K.; OHKAWARA, M. H.; DIAS, J. F.; KATSURAGAWA, M. Description of larvae and juveniles of *Bairdiella ronchus* (Sciaenidae: Teleostei) in Southeastern Brazil. **Sci. Mar.**, v. 71, n. 2, p. 249-257, 2007.
- KATSURAGAWA, M.; MATSUURA, Y.; SUZUKI, K.; DIAS, J. F.; SPACH, H. L. The Ichthyoplankton of the Ubatuba Region (São Paulo State, Brazil): composition, distribution and seasonal occurrence (1985-1988). **Publ. Esp. Inst. Oceanogr.**, v. 1, n. 10, p. 85-121, 1993.
- KATSURAGAWA, M.; DIAS, J. F. Distribuição do ictioplâncton na região de São Sebastião, SP. **Relat. Téc. Inst. Oceanogr.**, v. 41, p. 15-28, 1997.
- LALLI, C. M.; PARSONS, T. R. **Biological Oceanography: an introduction**. Oxford: Butterworth-Heinemann, 1995. 301 p.
- LEGENDRE, P.; LEGENDRE, L. **Numerical ecology**. 2. ed. Amsterdam: Elsevier Science B.V., 1998. 870 p. (Developing in Environmental Modelling, 20).
- LEIS, J. M. The pelagic stage of reef fishes: the larval biology of coral reef fishes. In: SALE, P. F. (Ed.). **The ecology of fishes on coral reefs**. San Diego: Academic Press, 1991. p. 183-230.
- LOPES, R. M.; KATSURAGAWA, M.; DIAS, J. F.; MONTÚ, M. A.; MUELBERT, J. H.; GORRI, C.; BRANDINI, F. P. Zooplankton and ichthyoplankton distribution on the Southern Brazilian shelf: an overview. **Sci. Mar.**, v. 70, n. 2, p. 189-202, 2006.
- LOWE-MCCONNELL, R. H. **Ecological studies in tropical fish communities**. Cambridge: New York: Cambridge University Press, 1987. 382 p. (Cambridge tropical biology series).
- MATSUURA, Y.; NAKATANI, K. Ocorrência de larvas e jovens de peixes na Ilha Anchieta (SP), com algumas anotações sobre a morfologia da castanha, *Umbrina coroides* Cuvier, 1830. **Bol. Inst. Oceanogr.**, v. 28, n. 1, p. 165-183, 1979.
- MATSUURA, Y.; SPACH, H. L.; KATSURAGAWA, M., Comparison of spawning patterns of the Brazilian sardine (*Sardinella brasiliensis*) and anchoita (*Engraulis anchoita*) in Ubatuba region, Southern Brazil during 1985 through 1988. **Bol. Inst. Oceanogr.**, v. 40, n. 1/2, p. 101-115, 1992.
- MIRANDA, L. B.; KATSURAGAWA, M. Estrutura Térmica na Região Sudeste do Brasil (outubro/novembro de 1998). **Publ. Esp. Inst. Oceanogr.**, v. 8, p. 1-14, 1991.
- NELSON, J. S. **Fishes of the world**. 4. ed. Hoboken: John Wiley, 2006. 601 p.
- NORCROSS, B. L.; SHAW, R. F. Oceanic and estuarine transport of fish eggs and larvae: a review. **Trans. Am. Fish. Soc.**, v. 113, n. 2, p. 153-165, 1984.
- PAIVA FILHO, A. M.; SCHMIEGELOW, J. M. M. Estudo sobre a ictiofauna acompanhante da pesca do camarão sete-barbas (*Xyphopeneaus kroyeri*) nas proximidades da Baía de Santos, SP. **Bol. Inst. Oceanogr.**, v. 34, p. 79-85, 1986.
- PAIVA FILHO, A. M.; GIANNINI, R.; RIBEIRO NETO, F. B.; SCHMIEGELOW, J. M. M. **Ictiofauna do complexo baía-estuário de Santos e São Vicente, SP, Brasil**. São Paulo: Instituto Oceanográfico/Universidade de São Paulo, 1987. p.1-10. (Relatório Interno, v. 17).
- PESSANHA, A. L. M.; ARAÚJO, F. G.; AZEVEDO, M. C. C. GOMES, I. D. Variações temporais e espaciais na composição e estrutura da comunidade de peixes jovens da baía de Sepetiba, Rio de Janeiro. **Rev. Bras. Zool.**, v. 17, n. 1, p.251-261, 2000.
- PIRES-VANIN, A. M. S.; CASTRO, B. M.; MIRANDA, L. B.; DIAS, J. F.; BICEGO, M. C.; KATSURAGAWA, M.; LOPES, R.; GIANESELLA, S. M. F.; CORBISIER, T. N. Síntese dos resultados e conclusões. In: PIRES-VANIN, A. M. S. (Coord.). **A influência do complexo estuarino da Baixada Santista sobre o ecossistema da plataforma adjacente (ECOSAN)**. São Paulo: ECOSAN, 2008. 643 p. (Technical report, n. 4).
- RAKOCINSKI, C. F.; LYCZKOWSKI-SHULTZB, J.; RICHARDSON, S. L. Ichthyoplankton assemblage structure in Mississippi Sound as revealed by Canonical Correspondence Analysis. **Estuarine, Coastal Shelf Sci.**, v. 43, n. 2, p. 237-257, 1996.
- SASSA, C.; KAWAGUCHI, K.; OOZEKI, Y.; KUBOTA, H.; SUGISAKI, H. Distribution patterns of larval Myctophid fishes in the transition region of the Western North Pacific. **Mar. Biol.**, v. 144, n. 3, p. 417-428, 2004.
- SCHAEFFER-NOVELI, Y.; CINTRON, G. **Guia para estudo de áreas de manguezal: estrutura, função e flora**. São Paulo: Caribbean Ecological Research, 1986. 150 p.
- SCHMIDT, T. C. S.; DIAS, J. F. Pattern of distribution and environmental influences on the Sciaenidae community of the Southeastern Brazilian Coast. **Braz. J. Oceanogr.**, v. 60, n. 2, p. 235-245, 2012.
- SINQUE, C. Larvas de Sciaenidae (Teleostei) identificadas na região estuarina lagunar de Cananéia. **Bol. Zool.**, v. 5, p. 39-77, 1980.
- SMITH, P. E.; RICHARDSON, S. L. **Standard techniques for pelagic fish egg and larvae surveys**. Rome: F.A.O., 1977. 100 p. (FAO Fisheries Technical Papers, v. 175).
- SPACH, H. L.; GODEFROID, R. S.; SCHWARZ Jr., C. S. R.; QUEIROZ, G. M. L. Temporal variation in fish assemblage composition on a tidal flat. **Braz. J. Oceanogr.**, v. 52, n. 1, p. 47-58, 2004.

- VALENTINI, H.; PEZZUTO, P. R. **Análise das principais pescarias comerciais da região Sudeste-Sul do Brasil com base na produção controlada do período 1986-2004**. São Paulo: Instituto Oceanográfico-USP, 2006. 56 p. (Série Documentos Revizee: Score-Sul).
- VAZZOLER, A. E. A. M. Síntese de conhecimentos sobre a biologia da corvina *Micropogonias furnieri* (Desmarest, 1823), na costa do Brasil. *Atlântica*, v. 13, n. 1, p. 55-74, 1991.
- VIEIRA, J. P.; CASTELLO, J. P. Environment and biota of the Patos Lagoon estuary – fish fauna. In: SEELIGER, U.; ODEBRECHT, C.; CASTELLO, J. P. (Eds.). **Subtropical convergence environments: the coast and sea in the Southwestern Atlantic**. Berlin: London: Springer Verlag, 1997, p. 56-62.

(Manuscript received 29 July 2013; revised 13 March 2014; accepted 20 March 2014)