

## Reproductive rates and performance traits in beef cattle infected by Bovine Herpesvirus

Claudia DEL FAVA<sup>1</sup>  
 Edviges Maristela PITUCO<sup>1</sup>  
 Leopoldo Andrade de  
 FIGUEIREDO<sup>2</sup>  
 Alexander George RAZOOK<sup>2</sup>  
 Joslaine Noely dos Santos  
 Gonçalves CYRILLO<sup>2</sup>  
 José Victor de OLIVEIRA<sup>3</sup>  
 Roberto Hauck REICHERT<sup>4</sup>  
 José Luiz D'ANGELINO<sup>5,6</sup>

**Correspondence to:**  
 CLAUDIA DEL FAVA  
 Centro de Pesquisa e Desenvolvimento de  
 Sanidade Animal  
 Instituto Biológico  
 Agência Paulista de Tecnologia dos  
 Agronegócios  
 Av. Conselheiro Rodrigues Alves, 1252  
 04014-002 - São Paulo - SP  
 delfava@biologico.sp.gov.br

Received: 28/03/2005  
 Accepted: 13/02/2006

1 - Centro de Pesquisa e Desenvolvimento de Sanidade Animal do Instituto Biológico da Agência Paulista de Tecnologia dos Agronegócios, São Paulo - SP  
 2 - CAPTA Bovinos de Corte do Instituto de Zootecnia da Agência Paulista de Tecnologia dos Agronegócios, Sertãozinho - SP  
 3 - APTA Regional – PRDTA Alta Mogiana, Colina - SP  
 4 - APTA Regional – PRDTA Vale do Ribeira, Registro - SP  
 5 - Faculdade de Medicina Veterinária e Zootecnia da Universidade de São Paulo, São Paulo - SP  
 6 - Faculdade de Medicina Veterinária da Universidade de Santo Amaro, São Paulo - SP

### Abstract

Reproductive rates and performance traits in beef cattle females infected by Herpesvirus Bovine-1 (BoHV-1) were evaluated in a farm at São Paulo state, Brazil. Gir, Guzerá, Nelore and Caracu purebred animals were examined to BoHV-1 (ELISA) at the beginning of the breeding season and the occurrence of reactors was 54.2% (386/712). BoHV-1 did not interfere in the pregnancy rates of both reagent - 80.3% (310/386) and non-reagent - 74.5% (243/326) females. It did not reduce the parturition rate of both reagent - 97.7% (300/307) and non-reagent - 93.8% (225/240) females. Total rate of stillbirths in BoHV-1 reagent females - 1.3% (4/300) did not differ from non-reagent females - 2.2% (5/225). BoHV-1 did not affect performance traits for reagent and non-reagent females, respectively, to daily weight gain during the breeding season ( $459.90 \pm 2.82$  g and  $466.63 \pm 2.87$  g), body condition score at the beginning of the breeding season ( $6.89 \pm 0.08$  and  $6.99 \pm 0.08$ ), body condition score at the end of the breeding season ( $7.73 \pm 0.06$  and  $7.71 \pm 0.06$ ), weight at parturition ( $419.17 \pm 3.34$  kg and  $425.97 \pm 3.22$  kg). It was concluded that non-vaccinated beef cattle females infected by BoHV-1 and bred under adequate extensive management, with body condition score over 5 and gain of weight during the breeding season presented good pregnancy, parturition and birth rates, no matter the breed, genetic group, age and seroconversion.

### Introduction

Bovine infectious rhinotracheitis / infectious pustular vulvovaginitis (IBR/IPV), caused by Bovine Herpesvirus type 1 (BoHV-1)<sup>1</sup> is a virosis that affects the respiratory and reproductive systems. Its economic impact can be not only observed in the losses it causes in infected animals, but also in the restrictions to the international trade of animals and animal products<sup>2</sup>.

Infected bovines are carriers of

BoHV-1 for their entire life and the virus remains in neurons of the trigeminal and sacral ganglia, in a non-infectious or latent form. It may be reactivated when animals are exposed to predisposing stressful events that decrease their immunological resistance, such as transportation, treatment with glucocorticoids and calving<sup>3</sup>. Carriers may reactivate and eliminate viral particles, most of the times without presenting clinical signs<sup>4,5,6</sup>.

In pregnant cows, viremia may lead

**Key-words:**  
 Bovine Herpesvirus 1.  
 Beef Cattle.  
 Zebu breeds.  
 Reproductive.  
 performance.  
 Performance traits.

to abortion, even if the female does not present rhinotracheitis symptoms<sup>3</sup>. BoHV-1 infection may lead to losses due to respiratory and reproductive symptoms as infertility, stillbirths or birth of weak calves. This virus interferes in reproductive rates of infected herds, affecting both embryonic and fetal development; abortion is more frequently observed between 3 and 9 months of pregnancy<sup>8,9,10</sup>. Infection of the reproductive tract may lead to endometritis and infertility<sup>11,12,13,14</sup>.

BoHV-1 is disseminated in Brazilian beef cattle herds<sup>15,16,17</sup>. However, the effects of the infection on reproductive and productive rates and on productive performance traits of beef cattle breeds endemically infected should be evaluated in field conditions. The profitability of beef cattle production is directly related to the number of animals born per year, which is dependent on the reproductive performance and performance traits, such as age at first calving, calving interval, weight at calving and at weaning. Therefore, performance traits should be analyzed together with sanitary evaluations in order to exclude the negative effects of sanitary problems on fertility of animals bred in extensive conditions.

Body condition score is an indirect evaluation of the effect of nutrition on reproduction. A body condition score equal to five or six at calving is necessary for females to present ovarian activity at the beginning of the breeding season, besides a quicker return to heat<sup>19,20,21,22</sup>.

The absence of data demonstrating the reproductive and productive effects of BoHV-1 in non-vaccinated beef cattle bred under extensive management and adapted to tropical conditions lead to the design of the present study. The objective of this trial was to evaluate performance traits and reproductive performance in females bred under the management conditions stated above.

## Materials and methods

This trial was performed in a rural property located in the northern region of the

state of São Paulo, Brazil, average altitude 548 meters above sea level, tropical humid climate, AW subtype, according to Köppen's classification. Meteorological indices observed during the period of study were average precipitation equal to 148.3 mm, mean minimum temperature equal to 15.3°C and mean maximum temperature equal to 32.5°C.

The herd was composed of purebred Gir, Guzerá, and Nelore *Bos indicus* and Caracu *Bos taurus*. Nelore animals are classified into three different genetic groups: control (selected for average postweaning weights), selection (both selected for high post weaning weights) and traditional, while Gir, Guzerá and Caracu animals come only from the selected lines.

Breeding season lasted 3 months, starting in November 1998 and ended in February 1999. Cows from the Selection and Traditional herds were mated: a group of 15 and 25 cows were, respectively, designed to bulls with two and three years old. In the control herd, cows were also mated with bulls two and three years old and each bull, no matter the age, was mated with 15 cows. A similar age distribution was followed within mating groups.

Pregnancy diagnosis was performed 60 days after the end of the breeding season and females were separated in groups of pregnant and non-pregnant cows<sup>24</sup>. The clinical evaluation of the reproductive tract of bulls and spermiogram were performed and males not approved in these examinations were not included in this trial<sup>25</sup>.

Pregnant cows were regrouped and sent to pastures that could stand a higher load, while non-pregnant cows were culled. Heifers that were not pregnant were kept in the herd in order to have a second chance during the next breeding season. All pregnant cows were supplemented before and after calving with silage or chopped pasture, corn and cotton meal.

Weight, height and body condition score of the cows were evaluated in the beginning and end of the breeding season, at calving and at weaning. Evaluation of body condition score ranged from 1 to 9, for *Bos indicus* breeds<sup>26</sup> and

*Bos taurus* breeds<sup>20</sup>.

Blood collection to identify the occurrence of BoHV-1 reagent and non-reagents<sup>27</sup> was performed in all animals in the mating groups, at the beginning of the breeding season, in November 1999. A second collection was performed in November 2000, in order to identify the animals that seroconverted until the end of the calving season<sup>27</sup>. BoHV-1 serodiagnosis was performed by indirect ELISA, as recommended by the Office International des Epizooties<sup>28</sup>. Commercial ELISA kits were used (Idexx; *HerdCheK*) according to the manufacturer's instructions.

Reproductive performance of BoHV-1 reagent and non-reagent females was evaluated by pregnancy rates, calving and stillbirth coefficient. Two-proportion tests comparing reagent and non-reagent animals were used, such as chi-square, exact Fisher's test; a was established at 5% for two-tailed test<sup>29,30</sup>. Pregnancy and calving rates, and stillbirth coefficient were analyzed, considering the different animal categories of the cows, such as breed (Gir, Guzerá, Nelore and Caracu), genetic group of the Nelore breed (Control, Selection and Traditional), age (2 to 3 years old, 3 to 4 years old, 4 to 5 years old and<sup>29</sup> 5 years old) and body conditions scores (from 2 to 5, from 5 to 6, from 6 to 7, from 7 to 8 and from 8 to 9).

BoHV-1 effect on performance traits of the cows, such as mean weight gain during breeding season, body condition scores at the beginning and the end of the breeding season were evaluated by the F test, through a variance analysis of random trials<sup>30</sup>. The *Statistical Analyses System* (SAS) was used, establishing a at 5%.

BoHV-1 effect on weight gain traits during breeding season and body condition score of the cows at the beginning and end of the breeding season took into account the following sources of variation: breed (Gir, Guzerá, Nelore and Caracu); age of the cow (2, 3, 4, 5, 6 and<sup>29</sup> 7 years old); age and breed interaction; calving in the previous

breeding season (1998-1999); calving in the 1999-2000 breeding season; BoHV-1 reagent or non-reagent; interaction between 1998-1999 calving and 1999-2000 calving; interaction between calving (1999-2000) and being BoHV-1 reagent or non-reagent.

BoHV-1 effect on weight of the cows at calving took into account the following sources of variation: breed (Gir, Guzerá, Nelore and Caracu); age of the cow (2, 3, 4, 5, 6 and<sup>29</sup> 7 years old); calving in the 1999-2000 breeding season; being BoHV-1 reagent or non-reagent; interaction between calving (1999-2000) and being BoHV-1 reagent or non-reagent.

## Results

Total pregnancy rate among reagent – 80.3% (310/386) and non-reagent females – 74.5% (243/326) did not differ statistically by the chi-square test ( $p > 0.05$ ) (Table 1). Pregnancy rates of reagent and non-reagent females did not vary according to age and breed ( $p > 0.05$ ). However, when each Nelore genetic group was considered, those in the Selection herd presented greater pregnancy rates in reagent females – 92.7% (51/55) than in non-reagent ones – 79.1% (53/67) ( $p < 0.05$ ), that is, the virus did not reduce fertility in the Nelore herd.

Considering the total number of females, the calving rate of reagent females – 97.7% (300/307) was greater than that observed for non-reagent ones – 93.8% (225/240). Difference was significant in the chi-square test ( $p < 0.05$ ). That is, BoHV-1 did not decrease calving (Table 1). No effect was observed in the calving rate of reagent and non-reagent females in each breed, age and Nelore genetic groups, either ( $p > 0.05$ ).

Condition of the calf at birth (live or stillborn), according to the conditions of the mother (reagent / non-reagent) showed that the stillbirth coefficient of the BoHV-1 females – 1.3% (4/300) did not differ from that of the females BoHV-1 non-reagent – 2.2% (5/225) (Table 1), even when breed, genetic group and age were considered

( $p > 0.05$ ); that is, BoHV-1 did not interfere with the stillbirth coefficient.

Body condition score of the females at the beginning and end of the breeding season was analyzed together with the reproductive and sanitary evaluation of the herd. Females that presented a body condition score below 5 at the beginning of the breeding season were also the ones that presented lower pregnancy rates – 65.1% (28/43) – than those with body condition scores above 5 – 82.2% (282/343) (Table 2) ( $p < 0.05$ ). At the end of the breeding season, pregnancy rates of reagent females with body condition score below 6 – 36.4%

(8/22) was lower than those with body condition score above 6 – 82.9% (301/363) ( $p < 0.05$ ), according to the chi-square test (Table 3).

Although seroconversion occurred in 10.3% (58/561) of the herd from the beginning to the end of the breeding season, there was no effect of BoHV-1 on the stillbirth coefficient, and pregnancy and calving rates, when compared that seroconverted with those that remained non-reagent and reagent during the same period (Table 4).

No statistical significant effects of BoHV-1 were observed ( $p > 0.05$ ) on the performance traits evaluated (Table 5).

Table 1 – Comparison between pregnancy and calving rates and stillbirth coefficients between cows reagent and non-reagent to BoHV-1, and level of statistical significance. São Paulo - SP, 2001

Variable	BoHV-1 reagent	BoHV-1 non-reagent	statistical significance
Total pregnancy rate	80.3 % a (310/386)	74.5 % a (243/326)	$\chi^2 = 3.07$ $P > 0.05$
Total calving rate	97.7 % a (300/307)	93.8 % b (225/240)	$\chi^2 = 4.52$ $P < 0.05$
Total stillbirth coefficient	1.3 % a (04/300)	2.2 % a (05/225)	Exact Fisher's test = 0.51 $P > 0.05$

Different letters show statistical significance between columns.

Table 2 – Chi-square test between the proportion of pregnant and non-pregnant BoHV-1 reagent cows, according to the body condition score at the beginning of the breeding season. São Paulo - SP, 2001

Body condition scores	BoHV-1 reagents		
	Pregnant	Non-pregnant	Statistical significance
2  - 5	5.1 % (28/43)	34.9% (15/43)	$\chi^2 = 6.03 *$
5  - 9	82.2% (282/343)	17.8% (61/343)	$p < 0.05$

Table 3 – Chi-square test between the proportion of pregnant and non-pregnant BoHV-1 reagent cows, according to the body condition score at the end of the breeding season. São Paulo - SP, 2001

Body condition scores	BoHV-1 reagents		
	Pregnant	Non-pregnant	Statistical significance
2  - 6	36.4% (08/22)	63.6% (14/22)	$\chi^2 = 25.5 *$
6  - 9	82.9% (301/363)	17.1% (62/363)	$p < 0.0001$

Table 4 – Two-proportion test between the groups Seroconversion, Non-reagents and Reagents by ELISA BoHV-1, for pregnancy and calving rates, and stillbirth coefficients. São Paulo - SP, 2001

Rates	BoHV-1 Seroconversion			BoHV-1 Non-reagents			BoHV-1 Reagents		
	No.	%	Total	No.	%	Total	No.	%	Total
Pregnancy	49	84.5 <sup>a</sup>	58	172	82.7 <sup>a</sup>	208	261	88.5 <sup>a</sup>	295
Calving	46	79.3 <sup>a</sup>	58	167	80.3 <sup>a b</sup>	208	258	87.5 <sup>a c</sup>	295
Stillbirth	0	0.0 <sup>a</sup>	46	04	2.4 <sup>a</sup>	167	05	2.0 <sup>a</sup>	257

z critical = 1.96 <sup>a</sup> = 5% Different letters show statistical significance between columns.

Table 5 – Summary of the variance analysis for some performance traits of the cows in the breeding season. São Paulo - SP, 2001

Performance traits	BoHV-1 reagents mean $\pm$ SD	BoHV-1 non-reagents mean $\pm$ SD	F	Pr > F
Mean daily weight gain (kg)	459.90 $\pm$ 2.82 <sup>a</sup>	466.63 $\pm$ 2.87 <sup>a</sup>	0.05	0.82
Body condition score at the beginning of the breeding season	6.89 $\pm$ 0.08 <sup>a</sup>	6.99 $\pm$ 0.08 <sup>a</sup>	0.52	0.47
Body condition score at the end of the breeding season	7.73 $\pm$ 0.06 <sup>a</sup>	7.71 $\pm$ 0.06 <sup>a</sup>	0.02	0.89
Weight at calving (Kg)	419.17 $\pm$ 3.34 <sup>a</sup>	425.97 $\pm$ 3.22 <sup>a</sup>	2.48	0.12

Different letters show statistical significance between columns.

## Discussion

BoHV-1 impact was studied in clinical, reproductive and nutritional evaluations in a herd naturally infected and non-vaccinated of beef cattle, made up of genetically improved Gir, Guzerá, Nelore and Caracu animals, adapted to tropical climate. Reproductive and productive effects were based on the comparison of BoHV-1 reagent and non-reagent animals bred under the same environmental and animal husbandry conditions, what validated the results obtained. In the Brazilian specialized literature there is a lack of information on reproductive and productive rates in herds of beef cattle breeds adapted to this kind of climate and to extensive animal husbandry, which is the main process of beef cattle breeding in Brazil.

The occurrence of 54.2% (386/712) BoHV-1 reagent females in the herd studied

was similar to the rate described in several Brazilian states<sup>15,16,17</sup>. A high rate of natural infection may characterize a condition of naturally acquired immunity<sup>7</sup>. This observation justifies the study of epidemiological, clinical and animal husbandry characteristics in the interpretation of BoHV-1 impact in endemically infected herds. Seroreagent animals in the present trial did not present genital or respiratory symptoms, condition that was also described by other authors<sup>7,31</sup>.

Although BoHV-1 infection has been described as causing abortion<sup>8,9,10</sup>, endometritis and infertility<sup>11,12,13,14</sup>, the herd studied did not present significant reproductive disorders, when BoHV-1 reagent females were compared to non-reagent ones in the control groups. In spite of the fact that literature reports that pregnant females infected by BoHV-1 may abort more frequently in the second or third

trimesters of pregnancy<sup>8,9,15</sup>, BoHV-1 infection did not cause a decrease in calving rates in the herd studied. The virus is also reported to lead to the birth of weak calves or stillbirths<sup>7,32</sup>. This effect, however, was not observed in the herd evaluated, when infected and non-infected females were compared. Besides the effect of BoHV-1 on stillbirth coefficient, other factors, such as age and breed of the mother were also evaluated, for they may be associated with the death of neonates<sup>33</sup>. There was no interference in relation to breed, Nelore genetic groups and age, though.

Females that seroconverted did not present any effects on stillbirth coefficient, and pregnancy and calving rates, what is in accordance with the results of other trials<sup>34,13</sup>. However, it has been reported that primo infections may lead to losses due to reproductive and respiratory problems<sup>35</sup>.

Nutritional status of beef cattle is considered to be a limiting factor in the reproduction of females bred under extensive husbandry. Because of this, evaluations of body condition score were performed together with the sanitary and reproductive evaluation of the herd, in order to exclude possible nutritional effects that may interfere with fertility. Pregnancy rates in reagent females showed that body condition score below 5, both in the beginning and end of the breeding season, was the factor that reduced animal fertility, and not virus infection. Body condition score above 5 is technically recommended when good reproductive rates are aimed, for it favors ovarian activity at the beginning and end of the breeding season, besides a quicker return to heat<sup>20</sup>.

BoHV-1 reagent or non-reagent pregnant animals gained weight and

improved body condition during the breeding season, even with most of the cows in lactation, as seen by the mean weight gain obtained. Although lactation has a suppressive effect on ovulation, this is less evident in animals that present moderate body condition scores at calving, or animals that did not present a drastic weight loss after calving<sup>19,22</sup>. Lactation is very demanding on the organism and together with the inhibition of GnRH release during milking, it makes difficult for cows presenting body condition scores below four to return to heat<sup>27</sup>. Therefore, the good nutritional status of the herd is favorable for reproduction. However, there are no published data to draw a possible comparison with BoHV-1 effects observed here.

## Conclusions

Beef cattle females infected and non-vaccinated against BoHV-1, submitted to extensive management, presenting body condition scores above five at the beginning and end of the breeding season and presenting weight gain during the breeding season showed good fertility and calving rates, as well as low stillbirth coefficients, no matter the breed, genetic group, age and seroconversion.

## Acknowledgements

The authors would like to thank Mr. Lúcio Aparecido Furtado for animal husbandry; to Mrs. Rosângela Maria Furtado and Mr. Sebastião Aparecido Teixeira for organizing the data files; to Mr. Antônio José de Lima, for providing all the breeding book information. The authors would like to thank FAPESP (process no. 99/05068-3) for the financial support.

## **Índices reprodutivos e características de desempenho em bovinos de corte infectados pelo Herpesvírus Bovino 1**

### **Resumo**

Avaliou-se índices reprodutivos e características de desempenho em fêmeas bovinas de corte, infectadas pelo Herpesvírus Bovino-1

**Palavras-chave:**  
Herpesvirus Bovino 1.  
Bovinos de Corte.  
Zebuíños.

(BoHV-1), em um rebanho no Estado de São Paulo, Brasil. Animais das raças Gir, Guzerá, Nelore e Caracu foram monitorados no início da estação de monta e a sororeatividade ao BoHV-1 pelo teste ELISA foi 54,2% (386/712). O BoHV-1 não interferiu no índice de prenhez de matrizes reagentes - 80,3% (310/386) e não reagentes - 74,5% (243/326) e nem reduziu a taxa de parição de matrizes reagentes - 97,7% (300/307) e não reagentes - 93,8% (225/240). O coeficiente de natimortalidade de matrizes reagentes ao BoHV-1 - 1,3% (4/300) não diferiu da encontrada para as não reagentes - 2,2% (5/225). O BoHV-1 não afetou a média de algumas características de desempenho de fêmeas reagentes e não reagentes, respectivamente, como ganho de peso médio diário durante a estação de monta ( $459,90 \pm 2,82$  g e  $466,63 \pm 2,87$  g), condição corporal na entrada da estação de monta ( $6,89 \pm 0,08$  e  $6,99 \pm 0,08$ ), condição corporal na saída da estação de monta ( $7,73 \pm 0,06$  e  $7,71 \pm 0,06$ ) e peso à parição ( $419,17 \pm 3,34$  kg e  $425,97 \pm 3,22$  kg). Concluiu-se que matrizes de corte infectadas pelo BoHV-1 e não vacinadas, criadas sob condições adequadas de manejo zootécnico extensivo e nutricional, como escore corporal acima de 5 e ganho de peso durante a estação de monta, apresentaram bons índices de prenhez, parição e natalidade, independente da raça, grupo genético, faixa etária e soroconversão.

Performance reprodutiva.  
Características de desempenho.

## References

- 1 VAN REGENMORTEL, M. H. V. et al. (Ed.). *Virus Taxonomy: the classification and nomenclature of viruses*. 7th Report. San Diego: Academic Press, 2000. 1167p. Chapter 00.031: Family Herpesviridae.
- 2 OIE. WORLD ORGANIZATION FOR ANIMAL HEALTH. Infectious Bovine Rhinotracheitis. In: **Terrestrial Animal Health Code**. 13th Ed., 2004. Paris: OIE. Disponível em <[http://www.oie.int/eng/publicat/em\\_code.htm](http://www.oie.int/eng/publicat/em_code.htm)> Acesso em: 31 dez. 2004.
- 3 TIKOO, S. K.; CAMPOS, M.; BABIUK, L. A. Bovine Herpesvirus 1 (BHV-1): biology, pathogenesis and control. In: MARAMOROSCH, K.; MURPHY, F.A.; SHATKIN, A.J. **Advances in virus research**. San Diego: Academic Press, 1995. v. 45, p. 191-223.
- 4 ASHBAUGH, S. E. et al. Specific detection of shedding and latency of bovine herpesvirus 1 and 5 using a nested polymerase chain reaction. **Journal of Diagnostic Investigation**, v. 9, n. 4, p. 387-394. 1997.
- 5 ENGELS, M.; ACKERMANN, M. Pathogenesis of ruminant Herpesvirus infections. **Veterinary Microbiology**, v. 53, n. 1-2, p. 3-15, 1996.
- 6 KAASHOEK, M. J.; RIJSEWIJK, F. A. M.; OIRSCHOT, J.T. Persistence of antibodies against bovine herpesvirus 1 and virus reactivation two to three years after infection. **Veterinary Microbiology**, v. 53, n. 1/2, p. 103-110, 1996.
- 7 LEMAIRE, M.; PASTORET, P. P.; THIRY, E. Le contrôle de l'infection pas le virus de la rhinotracéite infectieuse bovine. **Annales de Médecine Vétérinaire**, v. 138, n. 3, p. 167180, 1994.
- 8 BARR, B. C.; ANDERSON, M. L. Infectious diseases causing bovine abortion and fetal loss. **Veterinary Clinics of North America. Food Animal Practice**, v. 9, n. 2, p. 343-368, 1993.
- 9 KIRKBRIDE, C.A. Managing an outbreak of livestock abortion - 2: diagnosis and control of bovine abortion. **Veterinary Medicine**, v. 80, n. 5, p. 70-79, 1985.
- 10 PITUCO, E. M. et al. Detecção do Herpesvírus Bovino1 (HVB-1) e do vírus da Diarréia Viral Bovina (BVDV) pela Imunofluorescência Direta (IFD) em fetos bovinos abortados. **Arquivos do Instituto Biológico**, v. 66, p. 44, 1999. Suplemento
- 11 KAHRS, R. F. Infectious Bovine Rhinotracheitis: a review and update. **Journal of the American Veterinary Medical Association**, v. 171, n. 10, p. 1055-1064, 1977.
- 12 KENDRICK, J. W.; McENTEE, K. The effect of artificial insemination with semen contaminated with IBR-IPV virus. **The Cornell Veterinarian**, v. 57, n. 1, p. 3-11, 1967.
- 13 PARSONSON, I. M.; SNOWDON, W. A. The effect of natural and artificial breeding using bulls infected with, or semen contaminated with, infectious bovine rhinotracheitis virus. **Australian Veterinary Journal**, v. 51, n. 8, p. 365-369, 1975.
- 14 WHITE, M. B.; SNOWDON, W. A. The breeding record of cows inseminated with a batch of semen contaminated with infectious bovine rhinotracheitis virus. **Australian Veterinary Journal**, v. 49, n. 11, p. 501-506, 1973.
- 15 PITUCO, E. M. et al. Detecção de anticorpos contra o Herpesvírus Bovino tipo 1 (HVB-1) em rebanhos de corte e leite com problemas reprodutivos no Brasil.

- Arquivos do Instituto Biológico**, v. 66, p. 126, 1999. Suplemento.
- 16 RICHTZENHAIN, L. J. et al. Rinotraqueíte infecciosa bovina: levantamento sorológico nos Estados de Minas Gerais, Mato Grosso do Sul, São Paulo, Rio de Janeiro, Paraná e Rio Grande do Sul. **Arquivos do Instituto Biológico**, v. 66, n. 1, p. 83-88, 1999.
- 17 VIDOR, T. et al. Herpes Bovino Tipo 1 (BHV 1): I. Sorologia de rebanhos com problemas reprodutivos. **Ciência Rural**, v. 25, n. 3, p. 421-424, 1995.
- 18 LOBO, R. B. **Programa de melhoramento genético da raça Nelore**. Ribeirão Preto: KBL, 1992. 56 p.
- 19 BOLAÑOS, J. M. et al. Influence of body condition and restricted suckling on post-partum reproductive performance of zebu cows in the humid tropics. **Reproduction of Domestic Animals**, v. 31, n. 2, p. 363-367, 1996.
- 20 RICE, L. E. The effects of nutrition on reproductive performance of beef cattle. **Veterinary Clinics of North America. Food Animal Practice**, v. 7, n. 1, p. 1-26, 1991.
- 21 RICHARDS, M. W.; SPITZER, J. C.; WARNER, M. B. Effect of varying levels of postpartum nutrition and body condition at calving on subsequent reproductive performance in beef cattle. **Journal of Animal Science**, v. 62, n. 2, p. 300-306, 1986.
- 22 SELK, G. E. et al. Relationships among weight change, body condition and reproductive performance of range beef cows. **Journal of animal science**, v. 66, n. 12, p. 3153-3159, 1988.
- 23 RAZOOK, A. G. et al. Seleção para peso pós-desmame em bovinos Nelore e Guzerá. Diferenciais e intensidades de seleção. **Boletim de Indústria Animal**, v. 45, n. 2, p. 241-271, 1988.
- 24 GRUNERT, E.; BIRGEL, E. H. **Obstetrícia veterinária**. 2. ed. Porto Alegre: Sulina, 1984. 323 p.
- 25 COLÉGIO BRASILEIRO DE REPRODUÇÃO ANIMAL. **Manual para exame andrológico e avaliação de sêmen animal**. 2. ed. Belo Horizonte: Colégio Brasileiro de Reprodução Animal, 1998. 49 p.
- 26 NICHOLSON, M. J.; BUTTERWORTH, M. H. A guide to condition scoring of Zebu Cattle. Addis Ababa: International Livestock Centre for Africa, 1986. 29 p.
- 27 CÔRTES, J. D. Indicadores da ocorrência de doenças em populações. In: CÔRTES, J. D. **Epidemiologia: conceitos e princípios**. São Paulo, Varela, 1993. p. 143-157.
- 28 OIE. WORLD ORGANIZATION FOR ANIMAL HEALTH. Infectious Bovine Rhinotracheitis. In: **manual of diagnostic tests and vaccines for terrestrial animals**. 5th Ed., 2004. Paris: OIE. Disponível em <[http://www.oie.int/eng/normes/mmanual/A\\_00055.htm](http://www.oie.int/eng/normes/mmanual/A_00055.htm)> Acesso em: 31 dez. 2004
- 29 BERQUÓ, E. S.; SOUZA, J. M. P.; GOTLIEB, S. L. D. **Bioestatística**. 2. ed. São Paulo: EPU, 1981. 350 p.
- 30 VIEIRA, S. **Introdução à bioestatística**. 3. ed. Rio de Janeiro: Campus, 1998. 196 p.
- 31 WYLER, R.; ENGELS, M.; SCHWYZR, M. Infectious bovine rhinotracheitis/vulvovaginites (BHV-1). In: WITTMANN, G. Ed., **Herpesvirus diseases of cattle, horses and pigs**. Massachusetts: Keuwer Academic Publishers, 1989. p. 1-72.
- 32 ROCHA, M. A. Diagnóstico da Rinotraqueíte Infecciosa Bovina (IBR). **Revista Brasileira de Reprodução Animal**, v. 23, n. 4, p. 535-539, 1999.
- 33 LASTER, D. B.; GREGORY, K. E. Factors influencing peri- and early postnatal calf mortality. **Journal of Animal Science**, v. 37, n. 5, p. 1092-1097, 1973.
- 34 PARSONSON, I.M. Infectious pustular vulvovaginitis in dairy cattle in Victoria. **Australian veterinary journal**, v. 40, n. 7, p. 257-260, 1964.
- 36 MILLER, J. M. The effect of IBR virus infection on reproductive function of cattle. **Veterinary Medicine**, v. 86, n. 1, p. 95-98, 1991.
- 37 MILLER, J. M.; VAN DER MAATEN, M. J. Effect of primary and recurrent infectious bovine rhinotracheitis virus infection on the bovine ovary. **American Journal of Veterinary Research**, v. 46, n. 7, p. 1434-1437, 1985.