

GENERATION OF REDUCED NICOTINAMIDE
ADENINE DINUCLEOTIDE PHOSPHATE
MEDIATED BY GLUCOSE-6-PHOSPHATE
DEHYDROGENASE AND 6-
PHOSPHOGLUCONATE DEHYDROGENASE IN
NEUTROPHILS OF THOROUGHBRED HORSES*

AMAURY ANTIQUERA LEITE
Assistent Teacher
Faculdade de Ciências Farmacêuticas de Araraquara -
UNESP

ORLANDO CESAR DE OLIVEIRA BARRETTO
Associate Professor
Faculdade de Medicina da USP

LUCILA FREIRE MEDEIROS
Associate Professor
Instituto de Ciências Biomédicas da USP

LUIZ OCTÁVIO MEDEIROS
Associate Professor
Instituto de Ciências Biomédicas da USP

LEITE, A.A.; BARRETTO, O.C. de O.; MEDEIROS, L.F.;
MEDEIROS, L.O.; Generation of reduced nicotinamide
adenine dinucleotide phosphate mediated by glucose-
6-phosphate dehydrogenase and 6-phosphogluconate
dehydrogenase in neutrophils of thoroughbred horses.
Braz. J. vet. Res. anim. Sci., São Paulo, v.28, n.1, p.7-9,
1991.

SUMMARY: Twenty adult thoroughbred horses were
investigated for neutrophil glucose-6-phosphate
dehydrogenase and 6-phosphogluconate dehydrogenase
specific activities which were found to be 945 ± 288
 mIU mg^{-1} of protein and 375 ± 88 mIU mg^{-1} of protein
respectively, per minute at 37°C .

UNITERMS: NADP; Glucose-6-phosphate dehydrogenase; 6-
Phosphogluconate dehydrogenase;
Neutrophils; Horses, PSI

INTRODUCTION

Reduced nicotinamide adenine dinucleotide
phosphate (NADPH) plays a key role in microbicidal
process carried out by neutrophils, as it is involved
in superoxide generation by NADPH oxidase (BABIOR et
al. ², 1981). In resting neutrophils, NADPH oxidase is
barely detected, but when they are stimulated for
phagocytosis the enzyme activity is considerably
increased (BABIOR et al. ¹, 1976; MC PHAIL et al. ⁹,
1976; HOHN; LEHRER ⁶ 1975; DE CHATELET et al. ⁴,
1974). This work aimed at assaying in thoroughbred
horse neutrophils, the two sequential enzymes of the
pentose shunt responsible for the NADP reduction to
NADPH, the glucose-6-phosphate dehydrogenase (E.C.
1.1.1.49) and the 6-phosphogluconate dehydrogenase.
Both enzymes keep the NADPH pool at optimal levels in
order to supply NADPH to the NADPH oxidase activity
during the neutrophil respiratory burst.

MATERIAL AND METHOD

Neutrophils from twenty adult thoroughbred horses
from the Jockey Club of São Paulo were studied; 20 ml
of blood were collected after the morning exercise in
heparin (10 IU ml^{-1} of blood). Soon after drawing the
blood, the neutrophils were separated according to
standard procedures (FERRANTE; THONG ⁵, 1980), by
using Ficoll 400.000 (Sigma Co.) and Hypaque 90%
(Wintrop Products Inc.). The neutrophils (95% of
purity-ascertained by examining 200 leucocytes in a
Romanowski stained smear) were suspended in 1.0 ml of
saline, lysed by freeze-and-thawing (conic tubes with
blood immersed in an acetone and dry ice mixture, and
defrost at 37°C), and centrifuged at 16.000 G. The
supernatant was employed for enzyme assay, and the
protein was measured according to LOWRY et al. ⁸
(1951).

Glucose-6-phosphate-dehydrogenase activity was
determined in a reaction system containing 100 mM
TRIS-HCl pH 8.0, 100 mM magnesium chloride, 0.2 mM
NADP and 0.6 mM glucose-6-phosphate (G-6-P) (BEUTLER
³, 1984). 6-phosphogluconate dehydrogenase activity
was followed in a reagent system containing 100 mM
TRIS-HCl pH 8.0, 100 mM magnesium chloride, 0.2 mM
NADP and 0.6 mM G-6-P (BEUTLER ³, 1984). Enzyme
activities expressed as international units, were
calculated as micromoles of NADP reduced to NADPH at
 340 nm per minute, per miligram of protein, at 37°C .
A Gilford spectrophotometer model 2400-2 with recorder
was employed.

* This work was performed in the Instituto de Ciências
Biomédicas da USP and in Instituto dos Laboratórios
de Investigação Médica, Faculdade de Medicina da USP.

RESULTS AND DISCUSSION

Glucose-6-phosphate dehydrogenase (G6PD) and 6-phosphogluconate dehydrogenase (6-PGD) activities assayed in 20 horses are shown in Tab. 1. The neutrophil G6PD specific activity exhibited values of 945 ± 288 mIU mg^{-1} of protein per minute at 37°C , and 375 ± 88 mIU mg^{-1} of protein per minute at 37°C for 6-PGD, indicating a clearly higher G6PD specific activity. Such an increase of activity has also been found in human neutrophils (492 mIU mg^{-1} of protein for G6PD and 259 mIU mg^{-1} for 6-PGD), according to LANE et al.⁷ (1984) and personal communication **.

Although from a different lineage, the erythrocytes may help understanding the neutrophil characteristics. In fact MEDEIROS et al.¹⁰ (1982) found in thoroughbred horse erythrocytes a great imbalance favouring G6PD activity, with a ratio G6PD/6-PGD = 16, whereas in neutrophils this ratio is 2.5 (Tab. 1).

These findings suggest that in erythrocytes there is a limiting role for 6-PGD but in neutrophils a limiting role for 6-PGD does not seem to occur. This is suggestive of a harmonic relationship between both enzymes, which must work together in order to maintain the NADPH generation, the crucial step for keeping the efficiency of phagocytic process.

There is, moreover, a striking increase in enzyme specific in neutrophils as compared to that in erythrocytes, disclosing an overwhelming increase in gene expression for both G6PD and 6-PGD, mostly for the later, which is 200 times more active in neutrophils than in erythrocytes.

A close correlation between G6PD and 6-PGD specific activities is disclosed by a Fisher's coefficient of 0.75 suggesting that both specific activities vary together.

Both G6PD and 6-PGD are involved in transforming NADP to the reduced state NADPH. It is known that NADPH inhibits G6PD (YOSHIDA¹¹, 1973). Inasmuch as in neutrophil respiratory burst there is a high NADPH consumption by NADPH oxidase, the decrease of NADPH concentration would allow G6PD, and possibly 6-PGD as well, to work at maximum rate, without the inhibitory effect of NADPH. The maximum rate would be reached depending only upon the availability of its substrate glucose-6-phosphate.

LEITE, A.A.; BARRETTO, O.C. de O.; MEDEIROS, L.F.; MEDEIROS, L.O.; Geração da nicotinamida adenina dinucleotídeo fosfato reduzida mediada pela glicose-6-fosfato desidrogenase e 6-fosfogliconato

desidrogenase em neutrófilos de cavalos Puro-Sangue Inglês. *Braz. J. vet. Res. anim. Sci.*, São Paulo, v.28, n.1, p.7-9, 1991.

RESUMO: Em vinte cavalos Puro-Sangue Inglês determinou-se a atividade da glicose-6-fosfato-desidrogenase e da 6-fosfogliconato-desidrogenase de neutrófilos, encontrando-se atividades específicas de 945 ± 288 mIU mg^{-1} de proteína e 375 ± 88 mIU mg^{-1} de proteína, respectivamente, por minuto a 37°C .

UNITERMOS: NADP; Glicose-6-fosfato desidrogenase; 6-Fosfogliconato desidrogenase; Neutrófilos; Cavalos, PSI

REFERENCES

- 01-BABIOR, B.M.; CURNUTTE, J.T.; MC MURRICH, B.J. The particulate superoxide-forming system from human neutrophils. Properties of the system and further evidence supporting its participation in the respiratory burst. *J. clin. Invest.*, v.58, p.989-96, 1976.
- 02-BABIOR, G.L.; ROSIN, R.E.; MC MURRICH, B.J.; PETERS, W.A.; BABIOR, B.M. Arrangement of the respiratory burst oxidase in the plasma membrane of the neutrophil. *J. clin. Invest.*, v.67, p.1724-8, 1981.
- 03-BEUTLER, E. Red cell metabolism: a manual of biochemical methods. 3. ed. Orlando, Grune & Stratton, 1984.
- 04-DE CHATELET, L.R.; MC CALL, C.E.; MC PHAIL, L.C.; JOHNSTON, R.B. Superoxide dismutase activity in leucocytes. *J. clin. Invest.*, v.53, p.1197-201, 1974.
- 05-FERRANTE, A.; THONG, Y.H. Simultaneous preparation of mononuclear and polymorphonuclear leucocytes from horse blood on Ficoll-Hypaque medium. *J. Immunol. meth.*, v.34. p.279-85, 1980.
- 06-HOHN, D.C.; LEHRER, R.I. NADPH oxidase deficiency in X-linked chronic granulomatous disease. *J. clin. Invest.*, v.55, p.707-13, 1975.
- 07-LANE, T.A.; BEUTLER, E.; WEST, C.; LAMKIM, G. Glycolytic enzymes of stored granulocytes. *Transfusion*, Philadelphia, v.24, p.153-6, 1984.

** BEUTLER, E. Personal communication, 1989.

- 08-LOWRY, O.H.; ROSEBROUGH, N.J.; FARR, A.L.; RANDALL, R.J. Protein measurement with the Folin phenol reagent. *J. biol. Chem.*, v.193, p.265-75, 1951.
- 09-MC PHAIL, L.C.; DE CHATELET, L.R.; SHIRLEY, P.S. Further characterization of NADPH oxidase activity of human polymorphonuclear leucocytes. *J. clin. Invest.*, v.58, p.774-80, 1976.
- 10-MEDEIROS, L.F.; MEDEIROS, L.O.; BERCIANO, M.A. Energy metabolism in the erythrocytes of thoroughbred horses connected with perinatal physiological hemolysis. *Comp. Biochem. Physiol.*, v.71 B, p.541-4, 1982.
- 11-YOSHIDA, A. Hemolytic anemia and G6PD deficiency. *Science*, v.179, p.532-7, 1973.

Recebido para publicação em 03/08/90

Aprovado para publicação em 12/04/91

TABLE 1 - Thoroughbred horse neutrophils glucose-6-phosphate dehydrogenase and 6-phosphogluconate dehydrogenase. São Paulo, 1989.

Enzyme activity in mIU mg ⁻¹ of protein ERYTHROCYTES (E)* NEUTROPHILS (N) N/E per minute at 37 °C			
Glucose-6-phosphate dehydrogenase	29.2 ± 4.5	945 ± 288	32
6-phosphogluconate dehydrogenase	1.84 ± 0.2	375 ± 88	208
G6PD/6-PGD	16	2.5	

* MEDEIROS et al. ¹⁰ (1982)