



Original articles

Bleeding in patients hospitalized with acute pulmonary embolism in Brazil

Leonardo Jordan Hansen Vizzotto¹, Corina dos Reis Sepeda¹, Carlos Henrique Miranda^{1*}*Division of Emergency Medicine, Department of Internal Medicine, Faculdade de Medicina de Ribeirão Preto, Universidade de São Paulo (FMRP-USP), Ribeirão Preto, SP, Brazil*

ARTICLE INFO

Keywords:

Pulmonary embolism
Bleeding risk
Adverse events

ABSTRACT

Objective: Acute Pulmonary Embolism (APE) is a disease with increasing incidence worldwide. Antithrombotics are the cornerstone of the treatment. Bleeding is an adverse event related to this therapy. The objective was to evaluate the prevalence of bleeding in a sample of Brazilian patients hospitalized with APE and the impact of this complication on mortality. Additionally, the performance of some bleeding predictive scores was evaluated in this sample.

Methods: A retrospective cohort study was carried out on patients hospitalized with APE from January 2009 through August 2017. The medical records of these patients were reviewed, and the bleeding recorded during hospital stay was classified according to the “Thrombolysis in Myocardial Infarction (TIMI) bleeding risk.” Five different predictive scores for bleeding after APE (RIETE, PE-SARD, VTE-BLEED, Kujjer, and ATRIA) were applied. Overall mortality at 30 days and one year were assessed.

Results: One hundred fifty-nine patients were included. The prevalence of any bleeding was 36/159 (23 %), major bleeding was 10/159 (06 %), minor bleeding was 11/159 (07 %), and bleeding requiring attention was 15/159 (10 %). Only major bleeding was associated with higher mortality at one-year follow-up with a Relative Risk (RR) of 2.00 (95 % CI 1.16–3.57; $p = 0.044$). All bleeding predictive scores evaluated showed low accuracy in identifying patients at higher risk of bleeding.

Conclusion: Patients hospitalized with APE in Brazil had a high prevalence of bleeding. The major bleeding increased the one-year mortality. The bleeding predictive scores assessed showed limited accuracy in identifying patients at high risk of bleeding.

Introduction

Acute Pulmonary Embolism (APE) is a highly prevalent disease considered among the top three causes of cardiovascular mortality.^{1,2} In the United States, there are approximately 370,000 cases per year, with mortality rates ranging from 16 % to 27 %.^{3,4} There were 42,000 hospitalizations due to APE, 1500 deaths per year, and >76 million reais invested in treatment between 2015 and 2019 in Brazil.⁵ The APE incidence is increasing globally. The aging population, increase in associated comorbidities, such as obesity and cancer, and greater access to imaging tests for diagnosis justified this fact.^{1,3}

Antithrombotics (heparins or direct-acting anticoagulants) must be promptly started when the diagnosis of APE is established. Reperfusion therapy with fibrinolytics is indicated in the presence of hemodynamic instability. The objective of this approach is to interrupt or reduce the clot's progression, thereby reducing morbidity, mortality, and event recurrence.⁶⁻⁹

Unfortunately, bleeding is a common and severe adverse event associated with this therapy. Based on international literature, this complication occurs in approximately 7 % of these patients. It leads to an estimated mortality of 20 %, with the highest occurrence of this complication occurring in the first seven days of treatment.⁶⁻⁹

However, to our knowledge, no scientific investigation evaluated the prevalence of bleeding, as well as the impact of this complication on mortality, in a sample of Brazilian patients with APE.

The early identification of patients at risk of bleeding is helpful in clinical practice because it allows for the selection of antithrombotic drugs that are more appropriate to the patient's profile and ensures more surveillance. Several clinical scores are proposed for predicting the risk of bleeding after APE; however, none of them were validated for the Brazilian population.⁶⁻¹⁰

The objective of this study was to evaluate the prevalence of bleeding in a sample of Brazilian patients with APE, as well as the impact of this complication on the mortality of these individuals. Furthermore, to

* Corresponding author.

E-mail address: chmiranda@fmrp.usp.br (C.H. Miranda).

<https://doi.org/10.1016/j.clinsp.2024.100573>

Received 2 August 2024; Accepted 18 December 2024

Available online 6 January 2025

1807-5932/© 2025 HCFMUSP. Published by Elsevier España, S.L.U. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Table 1
Bleeding risk predictive scores after acute pulmonary embolism.

| Score/Year of publication | Items of the score | Risk categories | Original goal |
|-------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------|----------------------------------|
| RIETE, ¹¹ 2008 | Recent major bleeding (2) Creatinine > 1.2 mg/dL (1.5) Anemia (1.5) Previous malignancy (1) Clinical-overt APE (1) Age > 75-years (1) | Low: 0-points Intermediate: 1–4-points High: > 4-points | DVT and/or APE |
| Kuijjer, ¹² 1999 | Age ≥ 60-years (1.6) Female (1.3) Malignancy (2.2) | Low: 0-points Intermediate: 1–2-points High: ≥ 3-points | DVT and/or APE |
| PE-SARD, ⁶ 2021 | Anemia (2.5) Syncope (1.5) Renal dysfunction (1) | Low: 0-points Intermediate: 1–2.5-points High: > 2.5-points | Only APE |
| VTE-BLEED, ¹³ 2016 | Active malignancy (2) Male with uncontrolled hypertension (1) Anemia (1.5) Previous bleeding (1.5) Renal dysfunction (1.5) Age > 60-years (1.5) | Low: < 2-points High: ≥ 2-points | DVT and/or APE |
| ATRIA, ¹⁴ 2011 | Anemia (3) Renal or liver disease (3) Age ≥ 75-years (2) Previous bleeding (1) Hypertension (1) | Low: 0–3-points Intermediate: 4-points High: > 4-points | Non-valvular atrial fibrillation |

DVT, Deep Vein Thrombosis; APE, Acute Pulmonary Embolism.

evaluate the prognostic performance of some bleeding predictive scores in this sample.^{11–15}

Methods

A cohort, retrospective, single-center study carried out in patients with a definitive APE diagnosis admitted to the Emergency Unit of the Hospital das Clínicas of the Ribeirão Preto School of Medicine of the University of São Paulo from January 2009 through August 2017. The study was approved by the research ethics committee of this institution under number CAAE 51,979,515.9.0000.5440 and followed the guidelines of the Declaration of Helsinki.

Participants and data

Data were collected from the medical records of patients admitted with a primary diagnosis of APE in the electronic hospital discharge sheet record using codes I26.0 (pulmonary embolism with mention of acute cor pulmonale) and I26.9 (pulmonary embolism without mention of acute cor pulmonale) according to the International Classification of Diseases version 10 (ICD-10).

The confirmation of APE diagnosis was performed through a compatible clinical presentation associated with at least one confirmatory test, which could be Computed Tomography Pulmonary Angiography (CTPA) ventilation and perfusion scintigraphy, or necropsy.

After inclusion, demographic, clinical, laboratory data, and treatment options were collected from the patient's medical records. The "Pulmonary Embolism Severity Index (PESI)" score was calculated for all patients. Five different scores predicting the occurrence of bleeding were applied for these patients: RIETE, PE-SARD, VTE-BLEED, Kuijjer, and ATRIA. The criteria included in each score are shown in Table 1.

Table 2
Bleeding Stratification according to "Thrombolysis in Myocardial Infarction (TIMI) bleeding risk".¹⁵

| TIMI bleeding stratification | Definition |
|------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Major | Intracranial hemorrhage; Fatal bleeding; Significant clinical bleeding (hemoglobin decrease ≥ 5 g/dL or hematocrit decrease ≥ 15 %). |
| Minor | Observable blood loss: decrease in the hemoglobin 3–5 g/dL or decrease in the hematocrit ≥ 10 %; Non-observable blood loss: hemoglobin decrease ≥ 4 g/dL or hematocrit decrease ≥ 12 %. |
| Attention required | Observable blood loss with hemoglobin decrease < 3 g/dL or hematocrit decrease < 9 %. |

When bleeding was present during hospitalization, it was classified according to the "Thrombolysis in Myocardial Infarction (TIMI) bleeding risk" criteria into major, minor, and bleeding requiring attention. These criteria are shown in Table 2.

The outcomes evaluated were general mortality at 30 days and one year. When this outcome did not occur during hospitalization, follow-up was conducted via telephone contact by a properly trained employee of the institution's clinical research unit.

Statistical analysis

Categorical variables were expressed as frequency and percentage. Continuous variables with normal distribution were expressed as mean and standard deviation, and the others as median and Interquartile Range (IQR). Category variables were compared through the Chi-Square test. Two quantitative variables with normal distribution were compared through the unpaired Student's *t*-test, and two variables with another type of distribution were compared through the Mann-Whitney test. Three or more quantitative variables with normal distribution were compared through the ANOVA test, and three or more variables with another type of distribution were compared through the Kruskal-Wallis test.

The Relative Risk (RR) and its respective 95 % Confidence Interval (95 % CI) were calculated to evaluate the association between bleeding and mortality.

The accuracy of different types of scores for predicting bleeding was evaluated through the Area Under (AUC) the Receiver Operating Characteristic Curve (ROC). A convenience sample was employed and no estimative was made to determine the sample size. A two-tailed *p*-value ≤ 0.05 was considered significant. Stata software version 13.1 (StataCorp LP, College Station, TX, USA) was used for statistical analysis.

Results

One hundred fifty-nine patients who met the diagnostic criteria for APE were included. CTPA was the primary diagnostic tool (78 %), followed by ventilation and perfusion scintigraphy (8 %) and necropsy (5 %). Necropsy was reserved for unstable patients who died before performing an imaging test. The mean age of the patients was 58 ± 17 years; white race (79 %) and female gender (56 %) were predominant. The average PESI score was 100 ± 43 and its distribution among the risk stratification groups was: I – very low (20.7 %), II – low (20.7 %), III – moderate (19.5 %); IV – high (12.5 %); V – very high (26.4 %). Circulatory shock occurred in 16/159 (10 %) of patients, and cardiorespiratory arrest in 11/159 (7 %).

The average length of stay was 7 ± 17 days; during this period, anticoagulation with low molecular weight heparin was used in 67 % of patients, and anticoagulation with unfractionated heparin in 22 %. Thrombolysis was performed using alteplase in 37 patients (23 %). After

Table 3

Comparison of demographic, clinical, laboratorial and therapeutic features in hospitalized patients with acute pulmonary embolism divided according to the presence or absence of bleeding events and TIMI bleeding categories (major, minor and with attention required).

| Features | Bleeding events | | | Bleeding category | | | p |
|------------------------------------------------------------|---------------------|---------------------|-------|-------------------|-------------------|-----------------------------|-------|
| | Absent (n = 123) | Present (n = 36) | p | Major (n = 10) | Minor (n = 11) | Attention required (n = 15) | |
| Demographics | | | | | | | |
| Male gender; n (%) | 56 (45) | 13 (36) | 0.316 | 03 (30) | 03 (27) | 07 (46) | 0.534 |
| White people; n (%) | 100 (81) | 27 (75) | 0.407 | 08 (80) | 06 (54) | 13 (86) | 0.174 |
| Age | | | | | | | |
| > 75-years, n (%) | 22 (17) | 05 (13) | 0.574 | 01 (10) | 01 (09) | 03 (20) | 0.801 |
| > 65-years, n (%) | 51 (41) | 17 (47) | 0.539 | 05 (50) | 04 (36) | 08 (53) | 0.761 |
| Age; years, mean ± SD | 57.2 ± 17.3 | 58.6 ± 16.0 | 0.668 | 60.8 ± 14.7 | 56.8 ± 14.7 | 58.6 ± 18.4 | 0.926 |
| Clinical findings | | | | | | | |
| Duration of symptoms; days, mean ± SD | 6.8 ± 10.9 | 6.8 ± 9.3 | 0.995 | 5.7 ± 6.1 | 5.9 ± 9.0 | 8.2 ± 11.4 | 0.925 |
| Dyspnea, n (%) | 98 (79) | 30 (83) | 0.751 | 08 (80) | 08 (72) | 14 (93) | 0.574 |
| Hemoptysis, n (%) | 10 (08) | 05 (13) | 0.306 | 02 (20) | 01 (09) | 02 (13) | - |
| Syncope, n (%) | 18 (14) | 05 (13) | 0.897 | 01 (10) | 03 (27) | 01 (06) | 0.503 |
| Cough, n (%) | 38 (30) | 12 (33) | 0.804 | 04 (40) | 03 (27) | 05 (33) | 0.929 |
| Fever, n (%) | 15 (12) | 04 (11) | 0.848 | 0 (0) | 01 (09) | 03 (20) | 0.500 |
| Pleuritic chest pain, n (%) | 35 (28) | 14 (38) | 0.245 | 04 (40) | 02 (18) | 08 (53) | 0.170 |
| Any chest pain, n (%) | 44 (35) | 14 (38) | 0.757 | 02 (20) | 05 (45) | 07 (46) | 0.527 |
| Cardiac arrest, n (%) | 07 (05) | 04 (11) | 0.260 | 02 (20) | 01 (09) | 01 (06) | - |
| Altered mental status, n (%) | 27 (21) | 10 (27) | 0.467 | 04 (40) | 03 (27) | 03 (20) | 0.599 |
| Circulatory shock, n (%) | 12 (09) | 04 (11) | 0.812 | 02 (20) | 01 (09) | 01 (06) | 0.728 |
| RR > 20 breaths/min, n (%) | 81 (65) | 25 (69) | 0.644 | 05 (50) | 06 (54) | 14 (93) | 0.190 |
| HR > 100 bpm, n (%) | 52 (42) | 14 (38) | 0.810 | 05 (50) | 01 (09) | 08 (53) | 0.153 |
| Signs of DVT, n (%) | 35 (28) | 12 (33) | 0.592 | 02 (20) | 05 (45) | 05 (33) | 0.591 |
| PESI, categories | | | 0.813 | | | | 0.354 |
| I, n (%) | 26 (21) | 07 (19) | | 0 (0) | 02 (18) | 05 (33) | |
| II, n (%) | 24 (19) | 09 (25) | | 04 (40) | 04 (36) | 01 (06) | |
| III, n (%) | 26 (21) | 05 (13) | | 01 (10) | 01 (09) | 03 (20) | |
| IV, n (%) | 16 (13) | 04 (11) | | 01 (10) | 02 (18) | 01 (06) | |
| V, n (%) | 31 (25) | 11 (30) | | 04 (40) | 02 (18) | 05 (33) | |
| PESI; mean ± SD | 99.5 ± 43.9 | 103.4 ± 44.6 | 0.639 | 114.0 ± 40.6 | 96.8 ± 48.5 | 101.1 ± 46.0 | 0.784 |
| RR; breaths/min, mean ± SD | 24.3 ± 7.8 | 23.6 ± 6.5 | 0.597 | 23.4 ± 8.80 | 20.6 ± 5.9 | 25.4 ± 5.0 | 0.466 |
| HR; bpm, mean ± SD | 95.2 ± 20.1 | 95.4 ± 17.8 | 0.944 | 96.2 ± 12.6 | 85.6 ± 17.9 | 101.6 ± 18.6 | 0.260 |
| SBP; mmHg, mean ± SD | 122.0 ± 27.0 | 118.2 ± 30.4 | 0.489 | 118.3 ± 44.4 | 121.4 ± 29.0 | 115.9 ± 22.3 | 0.868 |
| DBP; mmHg, mean ± SD | 75.0 ± 15.9 | 74.9 ± 21.4 | 0.969 | 68.3 ± 28.8 | 78 ± 24.3 | 76.5 ± 13.3 | 0.616 |
| SpO ₂ ; %, mean ± SD | 90.3 ± 7.8 | 92.3 ± 5.3 | 0.239 | 90.0 ± 5.8 | 96.4 ± 1.14 | 92.0 ± 5.5 | 0.314 |
| SI (HR/SBP), mean ± SD | 0.82 ± 0.27 | 0.89 ± 0.40 | 0.211 | 0.97 ± 0.51 | 0.76 ± 0.30 | 0.94 ± 0.39 | 0.230 |
| Risk factors | | | | | | | |
| Previous DVT, n (%) | 24 (19) | 10 (27) | 0.287 | 01 (10) | 04 (36) | 05 (33) | 0.287 |
| Active malignancy, n (%) | 10 (08) | 03 (08) | 0.969 | 01 (10) | 01 (09) | 01 (06) | - |
| Recent surgery < 1-month, n (%) | 20 (16) | 04 (11) | 0.438 | 0 (0) | 02 (18) | 02 (13) | 0.315 |
| Immobilization > 3-days, n (%) | 31 (25) | 11 (30) | 0.522 | 04 (40) | 04 (36) | 03 (20) | 0.589 |
| Fracture, n (%) | 16 (13) | 04 (11) | 0.763 | 01 (10) | 01 (09) | 02 (13) | 0.973 |
| Previous stroke, n (%) | 16 (13) | 04 (11) | 0.763 | 01 (10) | 01 (09) | 02 (13) | 0.973 |
| Oral contraceptive use, n (%) | 14 (11) | 04 (11) | 0.952 | 01 (10) | 01 (09) | 02 (13) | 0.988 |
| Obesity, n (%) | 44 (35) | 13 (36) | 0.920 | 03 (30) | 05 (45) | 05 (33) | 0.966 |
| Heart failure, n (%) | 16 (13) | 04 (11) | 0.763 | 0 (0) | 02 (18) | 02 (13) | 0.619 |
| COPD, n (%) | 14 (11) | 04 (11) | 0.964 | 01 (10) | 02 (18) | 01 (06) | 0.836 |
| Thrombophilia, n (%) | 04 (03) | 08 (22) | 0.001 | 01 (10) | 04 (36) | 03 (20) | - |
| Laboratory findings | | | | | | | |
| Anemia, n (%) | 53 (43) | 18 (50) | 0.487 | 06 (60) | 06 (54) | 06 (40) | 0.663 |
| Hemoglobin, g/dL, mean ± SD | 12.7 ± 2.3 | 12.5 ± 2.0 | 0.661 | 12.3 ± 1.9 | 12.2 ± 2.2 | 12.8 ± 2.1 | 0.887 |
| Platelets < 150 × 10 ³ , n (%) | 21 (17) | 10 (27) | 0.161 | 02 (20) | 03 (27) | 05 (33) | 0.449 |
| Platelets < 50 × 10 ³ , n (%) | 02 (1.6) | 01 (2.0) | - | 0 (0) | 0 (0) | 01 (06) | - |
| Platelets × 10 ³ (/mm ³), mean ± SD | 223.684 ± 96.815 | 210.361 ± 98.743 | 0.471 | 250.300 ± 126.316 | 193.909 ± 53.476 | 195.800 ± 102.451 | 0.416 |
| Creatinine, mg/dL, mean ± SD | 1.2 ± 0.5 | 1.2 ± 0.4 | 0.772 | 1.3 ± 0.4 | 1.1 ± 0.3 | 1.3 ± 0.5 | 0.837 |
| Creatinine clearance*, mean ± SD | 68.2 ± 28.1 | 61.6 ± 27.0 | 0.221 | 52.7 ± 22.3 | 67.2 ± 24.1 | 63.5 ± 31.6 | 0.388 |
| Lactate, mg/dL, mean ± SD | 2.4 ± 1.8 | 3.3 ± 4.1 | 0.176 | 2.3 ± 1.0 | 5.2 ± 6.6 | 2.3 ± 1.5 | 0.031 |
| Arterial pH, mean ± SD | 7.40 ± 0.1 | 7.37 ± 0.1 | 0.282 | 7.34 ± 0.1 | 7.34 ± 0.1 | 7.42 ± 0.07 | 0.231 |
| PT (INR), mean ± SD | 1.37 ± 0.88 | 1.58 ± 0.97 | 0.249 | 1.38 ± 0.55 | 1.96 ± 1.37 | 1.40 ± 0.75 | 0.239 |
| Troponin, ng/mL, mean ± SD | 0.16 ± 0.35 | 0.17 ± 0.38 | 0.900 | 0.16 ± 0.24 | 0.22 ± 0.54 | 0.12 ± 0.18 | 0.949 |
| NT-pro-BNP, pg/mL, mean ± SD | 3.546 ± 4.642 | 3.719 ± 3.630 | 0.878 | 4.162 ± 4.729 | 4.605 ± 3.979 | 2.327 ± 1.820 | 0.779 |
| Treatment | | | | | | | |
| Thrombolysis, n (%) | 26 (21) | 11 (30) | 0.240 | 03 (30) | 04 (36) | 04 (26) | 0.633 |
| UH, n (%) | 25 (20) | 11 (30) | 0.197 | 04 (40) | 03 (27) | 04 (26) | 0.499 |
| LMWH, n (%) | 81 (65) | 27 (75) | 0.329 | 05 (50) | 08 (72) | 14 (93) | 0.102 |
| Warfarin, n (%) | 79 (64) | 23 (63) | 0.892 | 04 (40) | 09 (81) | 10 (66) | 0.130 |
| DOAC, n (%) | 13 (10) | 02 (1.6) | 0.354 | 01 (10) | 0 (0) | 01 (06) | - |
| Echocardiographic findings | | | | | | | |
| RV dilatation | 55 (44) | 21 (58) | 0.102 | 07 (70) | 07 (63) | 07 (46) | 0.402 |
| RV failure | 30 (24) | 06 (16) | 0.243 | 02 (20) | 02 (18) | 02 (13) | 0.714 |
| Acute pulmonary hypertension | 57 (46) | 21 (58) | 0.177 | 07 (70) | 06 (54) | 08 (53) | 0.509 |

(continued on next page)

Table 3 (continued)

| Features | Bleeding events | | | Bleeding category | | | |
|------------------------------------------------------|---------------------|---------------------|-------|-------------------|-------------------|-----------------------------|-------|
| | Absent (n = 123) | Present (n = 36) | p | Major (n = 10) | Minor (n = 11) | Attention required (n = 15) | p |
| Estimated pulmonary artery pressure, mmHg, mean ± SD | 51.3 ± 24.6 | 54.2 ± 14.3 | 0.598 | 59.5 ± 5.2 | 48.0 ± 22.5 | 55.2 ± 6.8 | 0.717 |

SD, Standard Deviation; RR, Respiratory Rate; HR, Heart Rate; DVT, Deep Vein Thrombosis; PESI, Pulmonary Embolism Severity Index; SBP, Systolic Blood Pressure; DBP, Diastolic Blood Pressure; SpO₂, Peripheral Oxygen Saturation; SI, Shock Index; COPD, Chronic Obstructive Pulmonary Disease; PT, Prothrombin Time; UH, Unfractionated Heparin; LMWH, Low Molecular Weight Heparin; DOAC, Direct Oral Anticoagulants; RV, Right Ventricle.

Table 4

Association with different types of bleeding according to TIMI (Thrombolysis in Myocardial Infarction bleeding risk) classification and 30-day and 1-year overall mortality.

| Bleeding category (TIMI) | Outcome | | |
|--------------------------|--------------------|-------------|-------|
| | 30-day mortality | | |
| | RR | 95 %CI | p |
| Any bleeding | 1.28 | 0.65 – 2.50 | 0.475 |
| Required attention | 1.49 | 0.66 – 3.34 | 0.351 |
| Minor | 0.42 | 0.06 – 2.79 | 0.322 |
| Major | 2.05 | 0.90 – 4.70 | 0.121 |
| | One-year mortality | | |
| Any bleeding | 1.46 | 0.91 – 2.36 | 0.133 |
| Required attention | 1.59 | 0.90 – 2.80 | 0.142 |
| Minor | 0.56 | 0.15 – 2.00 | 0.326 |
| Major | 2.00 | 1.16 – 3.57 | 0.044 |

RR, Relative Risk; 95 % CI, 95 % Confidence Interval.

hospital discharge, anticoagulation maintenance was performed with warfarin in 64 % and direct oral anticoagulants (rivaroxaban) in 9.5 %. The inferior vena cava filter was used in 2.5 % of cases. The in-hospital mortality rate was 15 %. Overall 30-day mortality was 20.7 % and 1-year mortality was 31 %. The other demographic, clinical, and laboratory characteristics of the included patients are shown in Table 3.

The prevalence of any bleeding was 36/159 (23 %), of major bleeding was 10/159 (06 %), of minor bleeding was 11/159 (06 %), and of bleeding requiring attention was 15/159 (10 %) during hospitalization.

Table 5

Bleeding risk categories according to different bleeding prediction scores in patients with acute pulmonary embolism.

| Bleeding risk score | Bleeding events | | | Bleeding category | | | |
|------------------------------|---------------------|---------------------|----------------|-------------------|-------------------|--------------------------------|----------------|
| | Absent (n = 123) | Present (n = 36) | p ^a | Major (n = 10) | Minor (n = 11) | Attention required (n = 15) | p ^b |
| RIETE | | | | | | | |
| Low and intermediate, n (%) | 113 (92) | 33 (92) | 0.969 | 09 (90) | 10 (90) | 14 (93) | 0.847 |
| High, n (%) | 10 (08) | 03 (08) | | 01 (10) | 01 (09) | 01 (07) | |
| Mean ± SD | 2.41 ± 1.34 | 2.70 ± 1.31 | 0.254 | 3.05 ± 1.18 | 2.68 ± 1.50 | 2.50 ± 1.28 | 0.510 |
| Kuijjer | | | | | | | |
| Low, n (%) | 26 (21) | 29 (80) | 0.826 | 01 (10) | 02 (18) | 04 (27) | 0.838 |
| Intermediate and high, n (%) | 97 (79) | 07 (20) | | 09 (90) | 09 (82) | 11 (73) | |
| Mean ± SD | 1.62 ± 1.16 | 1.86 ± 1.25 | 0.290 | 1.96 ± 0.97 | 2.01 ± 1.29 | 1.69 ± 1.44 | 0.643 |
| PE-SARD | | | | | | | |
| Low, n (%) | 41 (33) | 10 (28) | 0.297 | 3 (30) | 3 (27) | 4 (26) | 0.355 |
| Intermediate, n (%) | 62 (51) | 16 (44) | | 3 (30) | 4 (36) | 9 (60) | |
| High, n (%) | 20 (16) | 10 (28) | | 4 (40) | 4 (36) | 2 (13) | |
| Mean ± SD | 1.52 ± 1.37 | 1.75 ± 1.52 | 0.409 | 1.85 ± 1.84 | 2.13 ± 1.53 | 1.40 ± 1.29 | 0.482 |
| VTE-BLEED | | | | | | | |
| Low, n (%) | 67 (54) | 13 (36) | 0.05 | 2 (20) | 4 (36) | 7 (47) | 0.141 |
| High, n (%) | 56 (46) | 23 (64) | | 8 (80) | 7 (64) | 8 (53) | |
| Mean ± SD | 2.21 ± 1.73 | 2.66 ± 1.52 | 0.161 | 3.2 ± 1.65 | 2.54 ± 1.66 | 2.36 ± 1.51 | 0.300 |
| ATRIA | | | | | | | |
| Low, n (%) | 98 (80) | 27 (75) | 0.553 | 8 (80) | 8 (73) | 11 (73) | 0.903 |
| Intermediate and high, n (%) | 25 (20) | 09 (25) | | 2 (20) | 3 (27) | 4 (27) | |
| Mean ± SD | 2.08 ± 2.12 | 2.38 ± 1.79 | 0.432 | 2.60 ± 1.71 | 2.18 ± 1.83 | 2.4 ± 1.92 | 0.843 |

^a Comparison between absent vs. present bleeding.

^b Comparison among absent vs. major vs. minor vs. require attention bleeding events.




Fig. 1. Receiver Operating Characteristic Curve (ROC) depicting the accuracy of different scores in predicting any bleeding (A), major bleeding (B), minor bleeding (C) and bleeding that required attention (D) in patients hospitalized with acute pulmonary embolism in Brazil. AUC, Area Under the Curve; 95 % CI, Confidence Interval of 95 %.

Discussion

This investigation showed a high prevalence of bleeding during hospitalization for APE in Brazil. Furthermore, major bleeding was associated with higher one-year mortality. Bleeding predictive scores showed unsatisfactory performance in identifying patients at higher risk of bleeding.

Kresoja et al. showed an in-hospital incidence of major bleeding of 3.5 % in patients with APE. Furthermore, patients with major bleeding had a higher risk of death within one-year follow-up with a Relative Risk (RR) of 3.6 (95 % CI 2.0–6.6; $p < 0.001$).¹⁶ Budaj-Fidecka et al. reported an incidence of major bleeding of 2.4 % and any bleeding of 6 % in individuals with APE during a three-month follow-up. In this same time, there was higher mortality in individuals who had major bleeding with a RR of 2.75 (95 % CI 1.29–5.87; $p = 0.009$).¹⁷ Compared to these two studies, the present investigation showed a higher in-hospital bleeding rate (6 % of major bleeding and 23 % of any bleeding). However, similar to these two reported studies, major bleeding was associated with a higher one-year mortality with an RR of 2.00 (95 % CI 1.16–3.57; $p = 0.044$).

An important detail that must be highlighted is the lack of standardization in the definition and classification of bleeding between studies. Several classifications are used for this purpose. This study used the TIMI classification for bleeding¹⁵; however, currently, the International Society on Thrombosis and Haemostasis (ISTH) criteria is the most used.¹⁸ In this last classification, major bleeding is considered to be any fatal bleeding and/or symptomatic bleeding in a critical area or organ, such as intracranial, intraspinal, intraocular, retroperitoneal, intra-articular or pericardial, or intramuscular with compartment

syndrome, and/or bleeding causing a fall in hemoglobin levels > 2.0 g/dL or leading to a transfusion of 2 units or more of whole blood or red cells. If the authors had used this last classification, the prevalence of major bleeding would have been even higher in the present study.

The higher prevalence of bleeding in this study could be explained in part by the greater severity of patients hospitalized with APE in Brazil. In a cohort of 1880 individuals with APE, Pollack et al. showed an overall 30-day mortality of 5.4 %.¹⁹ Laporte et al. showed a 30-day mortality of 3.3 % in a sample of 6518 patients diagnosed with APE.²⁰ In the present investigation, the authors observed an in-hospital mortality of 15 %, a 30-day mortality of 20.7 %, and a one-year mortality of 31 %. The incidence of circulatory shock upon hospital admission was higher in the present study's sample (10 %) when compared to studies by Pollack et al. (3.0 %; $p = 0.002$) and Laporte et al. (3.80 %; $p = 0.006$).^{19,20} Furthermore, our study included individuals with higher PESI scores (100 ± 43 vs. 88 ± 34 ; $p = 0.0001$) compared with Pollack's sample.¹⁹

In a cohort of 727 Brazilian patients with APE, Volschan et al. showed an in-hospital mortality of 19.5 % and circulatory shock prevalence of 19.9 %.²¹ In another Brazilian study, Soriano et al. also observed high circulatory shock and 30-day mortality rates, respectively 11 % and 23 %. In this last study, 59 % of the sample presented PESI > 84 -points.²² Probably, only the most severe APEs are diagnosed and hospitalized in Brazil, which ends up contributing to the higher bleeding in these patients.

Regarding predictive bleeding scores, Klok et al. evaluated the KUIJIER, RIETE, HEMORR2HAGES, HAS-BLED, and ATRIA scores to predict bleeding within 30-days in individuals with APE. As in the present investigation, the prognostic performance for predicting bleeding was limited with an AUC-ROC of 0.57 to 0.64.⁷ In agreement,

Zhu Zhang et al. showed an unsatisfactory accuracy of the Kuijer and RIETE scores for predicting major bleeding within three months after APE diagnosis with an AUC-ROC of 0.57 and 0.56; respectively.⁸

In a study to externally validate the PE-SARD score, 50,686 individuals with APE were included, Chopard et al. showed an AUC-ROC of 0.65 for this score in discriminating patients at risk of major bleeding within 30-days.⁶ Kresoja et al. reported an AUC-ROC of 0.69 for the VTE-BLEED score in predicting major bleeding in patients hospitalized with APE.¹⁶

Some limitations deserve to be highlighted. First, the study was unicentric and carried out in a highly complex tertiary referral hospital. This may have affected the findings by introducing a selection bias, leading to the inclusion of patients with greater severity and comorbidities. Second, the bleeding classification was carried out exclusively using the TIMI. Because of this, the comparison with other studies is difficult and it may have even underestimated the major bleeding in the present investigation. Third, this study was carried out when warfarin was the primary anticoagulant used in the long term; only 9.5 % of patients used Direct Oral Anticoagulants (DOACs), which may have contributed to the higher bleeding risk.

To our knowledge, this is the only study that evaluated the prevalence of bleeding and the impact of this complication on mortality during APE treatment in the Brazilian population. Furthermore, no validation study of predictive bleeding scores in patients with APE in Brazil was found.

In conclusion, the authors observed a high prevalence of bleeding in this sample of Brazilian patients hospitalized with APE. The presence of major bleeding was associated with a higher mortality within one year. The evaluated bleeding predictive scores showed unsatisfactory performance in identifying patients at high risk of bleeding. Therefore, the authors do not recommend using these scores when making decisions regarding antithrombotic therapy for these patients. National, multicenter, and prospective registries are necessary to more accurately determine the prevalence of bleeding in hospitalized patients with APE in Brazil and to develop appropriate bleeding predictive scores for this population.

Conflicts of interest

The authors declare no conflicts of interest.

References

1. Wang LT, Yang H, Zhang HD. Very early major bleeding in acute pulmonary embolism: could the French Pulmonary Embolism-Syncope, Anemia, and Renal Dysfunction score be applied to the Swiss cohort? *J Thromb Haemost.* 2023;21(10):2711–2714.
2. Mathew D, Seelam S, Bumrah K, Sherif A, Shrestha U. Systemic thrombolysis with newer thrombolytics vs anticoagulation in acute intermediate risk pulmonary embolism: a systematic review and meta-analysis. *BMC Cardiovasc Disord.* 2023;23(1):1–7.
3. Di Nisio M, van Es N, Büller HR. Deep vein thrombosis and pulmonary embolism. *Lancet.* 2016;388(10063):3060–3073.

4. Freund Y, Cohen-Aubart F, Bloom B. Acute Pulmonary Embolism. *JAMA.* 2022;328(13):1336 [Internet].
5. Pinheiro da Silva J, Barros Souza R, Costa de Oliveira L, De Barros Rocha L, J Luís Monteiro Spinelli, Simões Herminia. Hamad Farias do Couto M. Perfil Epidemiológico do Tromboembolismo Pulmonar no Brasil de 2015 a 2019. *BEPA Bol Epidemiológico Paul.* 2021;18(208):1–10.
6. Chopard R, Piazza G, Falvo N, Ecarnot F, Besutti M, Capellier G, et al. An Original Risk Score to Predict Early Major Bleeding in Acute Pulmonary Embolism: the Syncope, Anemia, Renal Dysfunction (PE-SARD) Bleeding Score. *Chest.* 2021;160(5):1832–1843.
7. Klok FA, Niemann C, Dellas C, Hasenfuß G, Konstantinides S, Lankeit M. Performance of five different bleeding-prediction scores in patients with acute pulmonary embolism. *J Thromb Thrombolysis.* 2016;41(2):312–320.
8. Zhang Z, Lei J, Zhai Z, Yang Y, Wan J, Xie W, et al. Comparison of prediction value of four bleeding risk scores for pulmonary embolism with anticoagulation: a real-world study in Chinese patients. *Clin Respir J.* 2019;13(3):139–147.
9. Mathonier C, Meneveau N, Besutti M, Ecarnot F, Falvo N, Guillon B, et al. Available bleeding scoring systems poorly predict major bleeding in the acute phase of pulmonary embolism. *J Clin Med.* 2021;10(16):1–14.
10. Badescu M, Ciociu M, Badulescu O, Vlădeanu MC, Bojan I, Vlad C, et al. Prediction of bleeding events using the VTE-BLEED risk score in patients with venous thromboembolism receiving anticoagulant therapy (Review). *Exp Ther Med.* 2021;22(5):1–6.
11. Ruíz-Giménez N, Suárez C, González R, Nieto JA, Todolí JA, Samperiz ÁL, et al. Predictive variables for major bleeding events in patients presenting with documented acute venous thromboembolism. Findings from the RIETE Registry. *Thromb Haemost.* 2008;100(1):26–31.
12. Kuijer PMM, Hutten BA, Prins MH, Büller HR. Prediction of the risk of bleeding during anticoagulant treatment for venous thromboembolism. *Arch Intern Med.* 1999;159(5):457.
13. Klok FA, Hösler V, Clemens A, Yollo WD, Tilke C, Schulman S, et al. Prediction of bleeding events in patients with venous thromboembolism on stable anticoagulation treatment. *Eur Respir J.* 2016;48(5):1369–1376.
14. Fang MC, Go AS, Chang Y, Borowsky LH, Pomernacki NK, Udaltsova N, et al. A new risk scheme to predict warfarin-associated hemorrhage: the ATRIA (Anticoagulation and Risk Factors in Atrial Fibrillation) study. *J Am Coll Cardiol.* 2011;58(4):395–401.
15. Rao AK, Pratt C, Berke A, Jaffe A, Ockene I, Schreiber TL, et al. Thrombolysis in myocardial infarction (TIM) trial-Phase I: hemorrhagic manifestations and changes in plasma fibrinogen and the fibrinolytic system in patients treated with recombinant tissue plasminogen activator and streptokinase. *J Am Coll Cardiol.* 1988;11(1):1–11.
16. Kresoja KP, Ebner N, Rogge NIJ, Sentler C, Keller K, Hobohm L, et al. Prediction and prognostic importance of in-hospital major bleeding in a real-world cohort of patients with pulmonary embolism. *Int J Cardiol.* 2019;290:144–149.
17. Budaj-Fidecka A, Kurzyna M, Fijałkowska A, Zylkowska J, Wieteska M, Florczyk M, et al. In-hospital major bleeding predicts mortality in patients with pulmonary embolism: an analysis of ZATPOL Registry data. *Int J Cardiol.* 2013;168(4):3543–3549.
18. Schulman S, Kearon C. Definition of major bleeding in clinical investigations of antihemostatic medicinal products in non-surgical patients. *J Thromb Haemost.* 2005;3(4):692–694.
19. Pollack CV, Schreiber D, Goldhaber SZ, Slattery D, Fanikos J, O'Neil BJ, et al. Clinical characteristics, management, and outcomes of patients diagnosed with acute pulmonary embolism in the emergency department: initial report of EMPEROR (multicenter emergency medicine pulmonary embolism in the real-world registry). *J Am Coll Cardiol.* 2011;57(6):700–706.
20. Laporte S, Mismetti P, Décousus H, Uresandi F, Otero R, Lobo JL, et al. Clinical predictors for fatal pulmonary embolism in 15 520 patients with venous thromboembolism: findings from the Registro Informatizado de la Enfermedad TromboEmbolica venosa (RIETE) registry. *Circulation.* 2008;117(13):1711–1716.
21. Volschan A, Albuquerque DC de, Tura BR, Knibel M de F, Souza PCP da SE, Toscano ML. Pulmonary embolism: multicenter registry in tertiary hospitals. *Rev Bras Ter Intensiva.* 2009;21(3):237–246.
22. Soriano L de A, Castro TT, Vilalva K, Borges M de C, Pazin-Filho A, Miranda CH. Validation of the Pulmonary Embolism Severity Index for risk stratification after acute pulmonary embolism in a cohort of patients in Brazil. *J Bras Pneumol.* 2019;45(1), e20170251.