

Three-dimensional soft tissue comparison in non-growing patients treated for skeletal Class II Division I malocclusion: mandibular advancement surgery vs. Herbst Appliance

Gökhan ÇOBAN¹ | Taner ÖZTÜRK¹ | Gökhan TÜRKER² | Ebru Topcuoğlu HASHIMLI¹ | Yusuf Nuri KABA³

¹Erciyes University, Department of Orthodontics, Faculty of Dentistry, Kayseri, Turkey.

²Autonomous researcher, Mersin, Turkey.

³Uludag University, Faculty of Dentistry, Department of Oral and Maxillofacial Surgery, Bursa, Turkey.

Abstract

Objective: This study used three-dimensional (3D) stereophotogrammetry to compare changes in facial soft tissues of patients with skeletal class II malocclusion with mandibular retrognathia treated with orthognathic surgery (OS) or Herbst appliance (HA). **Methodology:** This retrospective study included 15 adults treated with bilateral sagittal split osteotomy without genioplasty (OS) and 16 adults treated with HA. The patients were treated successfully, with class I occlusal relationships with normal overjet and overbite. Three-dimensional stereophotogrammetric records acquired in habitual occlusion from before and after fixed orthodontic treatment were analyzed. Lateral cephalometric radiographs were used to evaluate skeletal parameters and sagittal oropharyngeal airway length. **Results:** In HA-group, mandibular corpus length, anterior facial height, posterior facial height, B⁰, vertical angle, and mentolabial angle increased; lower facial width, mandibular angle, mandibular convexity angle, ANB⁰, and A⁰ decreased. In OS-group, mandibular length, mandibular corpus length, and facial convexity angle increased significantly compared with those in HA-group; posterior facial height, B⁰, vertical angle, and mentolabial angle also increased, but mandibular angle and ANB⁰ decreased significantly. Only a significant difference in coordinate changes was observed for the pogonion in the sagittal direction. According to the cephalometric analysis, SNB⁰ and Pg-NA perpendicular measurements increased significantly in the OS group compared with the HA group. In both groups, sagittal oropharyngeal airway length increased significantly post-treatment, with no significant intergroup difference. **Conclusion:** In patients treated with OS, skeletal advancement resulted in greater increases in mandibular and corpus lengths, along with forward positioning of the pogonion. Based on 3D soft tissue and cephalometric comparisons, the Herbst appliance seems to be a viable non-surgical alternative for young adults with moderate Class II malocclusion. Both treatment modalities also contributed to significant improvements in sagittal oropharyngeal airway length.

Keywords: Orthognathic surgery. Anthropometric soft tissue landmarks. Herbst appliance. Dentofacial orthopedics. Stereophotogrammetry.

Correspondence:
Taner Öztürk - Erciyes University - Faculty of Dentistry - Department of
Orthodontic - 38039 - Melikgazi - Kayseri - Turkey.
e-mail: tanertr35@gmail.com



Received: May 09, 2025
Revised: July 31, 2025
Accepted: September 15, 2025

Editor: Ana Carolina Magalhães
Associate Editor: Daniela Rios Honório

Introduction

One-fourth of the population have class II malocclusions, which predominantly result from genetic and environmental factors (i.e., parafunctional behaviors, such as persistent finger and tongue sucking habits, lip trap, or early loss of deciduous teeth) that can be observed at variable rates over time.^{1,2}

Class II, division I malocclusions are the most common form of malocclusion³ and are often characterized by proclined maxillary incisors with or without a relatively narrow maxilla, an increased overjet, incomplete lip closure, and mandibular skeletal retrusion, which is the most common diagnostic result.^{1,4,5}

The backward positioning of the tongue, due to mandibular retrusion, may cause a narrowing of the pharyngeal airway and breathing difficulties.⁶ Moreover, retrognathic mandible can cause sleep-disordered breathing, which ranges from snoring to stroke.⁷ Moreover, maxillary incisor trauma increases in class II malocclusions, and oral health-related quality of life decreases.⁸⁻¹⁰

The treatment of skeletal class II malocclusion, which is observed with all those characteristics, is important not only to address chewing and biting but also to improve functions such as breathing and speaking as well as aesthetic, psychosocial, and self-image development.¹¹

Three basic approaches have been proposed in the treatment of skeletal malocclusion: growth modification, camouflage, and orthognathic surgery.¹² Although growth modification is the first option for individuals who are still growing and developing, camouflage treatment can be also applied. Late adolescents and adults no longer have significant growth, so the treatment options are camouflage and orthognathic surgery.¹³

However, effective correction can also be achieved with the Herbst appliance (HA), the most frequently used fixed bite jumping orthopedic device,¹⁴ in class II malocclusions, even in adults, and it may be an alternative to extraction treatments and orthognathic surgery.¹⁵ The HA keeps the mandible in a constantly forced protruded position when the mouth is open or closed. Because of this mechanical stress, different degrees of changes can be seen in the mandible (and its position¹²), muscle function,

and temporomandibular joint (condyle-glenoid fossa) structures, which, combined with dentoalveolar effects, depend on age and maturity stage.^{3,16} Additionally, the HA stimulates condylar growth and remodels the glenoid fossa in children and adults.¹⁷ For all these reasons, the HA can be considered a fixed functional appliance.¹⁸ Evidence indicates that growth-adapting treatment with the HA is possible even after 20 years and that it can replace or be an alternative to surgery, especially in treatment for mild skeletal class II¹⁵, and its stability was as high as that of orthognathic surgery in a three-year long-term follow-up.¹⁹

Various types of HAs have been used, depending on the method of attachment to the teeth. These are banded HA, cast splints, acrylic splints, and stainless-steel crowns.²⁰ The cast splint and banded versions are cemented to the teeth whereas the acrylic splint version requires an interocclusal acrylic layer, and it can design to be complete or partially removable.²¹ The stainless-steel cast splint HA has several advantages over other types: it fits perfectly, it is rigid and hygienic, the chair time is short, and problems such as band or crown breaks are less common.¹⁸

Combined orthodontic and orthognathic surgery (OS) is a well-known method that has been used for about 70 years; It has a high success rate in resolving the underlying skeletal discrepancy in skeletal class II malocclusions, correcting function, and improving soft tissues by reducing facial convexity, thus improving aesthetics.²² The most common mandibular surgical procedure is mandibular bilateral sagittal split osteotomy (BSSO).²³

The stereophotogrammetry method, which was introduced to dentistry in the 1940s and is frequently used in 3D imaging of the face²⁴, is fast, easy, and non-invasive and enables the evaluation of facial soft tissues without radiation.²⁵ This provides comprehensive 3D visualization and precise assessment of facial structures, making it particularly suitable for evaluating and comparing morphological changes in orthodontic and surgical treatments.²⁶

Although studies have compared the treatment effectiveness and outcomes of dentofacial orthopedics and OS,^{3,15,27} no studies have compared the treatment effectiveness of these two methods using 3D images. Therefore, this study compared the changes in the soft tissues of patients (whose growth was completed)

who were treated with the cast splint stainless-steel HA and mandibular advancement with BSSO (without genioplasty) using the stereophotogrammetry method.

Methodology

This was a retrospective, cross-sectional, and observational study involving a 3D outcome analysis of two groups of patients, which was approved by the Erciyes University Ethics Committee (Approval Code: 2023/695). The study was conducted ethically in accordance with the guidelines of the World Health Organization's Declaration of Helsinki.

Sample size calculation

The G*Power (v. 3.1.9.7, Franz Faul, Universität Kiel, Germany) software was used to determine the sample size. The results reported by Ruf and Pancherz (N–Sn–Pog facial convexity angle) were used for power analysis, and the following parameters were applied: $1-\alpha=0.05$, 90% power, and $d=0.812$. The calculation indicated that at least 15 individuals should be included in each group.¹⁵

Patients

The study population comprised 15 adults (11 women and four men, mean age: 19.9 ± 2.43 years, range: 17.2–25 years) treated with OS and 16 adults (ten women and six men, mean age: 16.01 ± 0.78 years, range: 15.1–17.1 years) treated with the HA. The patients were recruited between June 2020 and January 2023 and were treated at the Departments of Orthodontics and Oral and Maxillofacial Surgery, Faculty of Dentistry, Erciyes University.

The following patients were included:

- Patients who had skeletal class II, division 1 malocclusion and were treated without extraction (except for the third molars) ($ANB^\circ > 4^\circ$).
- Patients who had normal vertical growth pattern ($30^\circ < SN/GoGn < 35^\circ$).
- Patients who had appropriate 3D stereophotogrammetric recordings at the beginning and end of treatment, attended their appointments regularly, had complete follow-up, and did not report any incompatibilities, such as device breakage or usage errors.
- Patients who were treated with HA and OS

and had class I occlusions with normal overjet and overbite at the end of treatment.¹⁵

Patients who had syndromes, congenital anomalies, craniofacial deformities, excessive vertical growth patterns, missing teeth, an overjet of less than 6 mm, severe facial asymmetry, or severe maxillary transverse deficiency, along with those who had previously received orthodontic treatment, and those whose growth and development was not complete (according to hand–wrist radiography evaluation) were excluded from the study.

Skeletal maturity of the patients was classified as described by Ruf and Pancherz²⁸ according to hand and wrist roentgens, and all patients were found to be in the R-J stage. Accordingly, all patients in both groups were evaluated as young adults and adults.

Herbst appliance group

The HA type used in this study was a stainless-steel cast splint with a multibracket system. In this type, cast crowns cover the maxillary and mandibular molars and premolar teeth, and a lingual arch connects the mandibular part. A 10 mm hyrax screw (Dentaurum, Pforzheim, Germany) was also used to bring the maxillary parts together and provide expansion. This was activated via a semi-rapid protocol (two turns on the first three days, one turn on the following four days, one turn on every following two days) until adequate expansion was achieved. A Dentaurum type 1 telescopic system (Herbst Set I; Dentaurum, Ispringen, Germany) was used.¹⁶ Guide tubes were soldered to the steel crowns on the upper second molar teeth, and slide pins were soldered to the steel crowns on the lower first premolar teeth. The construction process of this device was performed by experienced technicians in a laboratory on plaster models taken from the patient. The HA was cemented to the teeth with 3M ESPE glass ionomer cement (Ketac-Cem; 3M ESPE, Seefeld, Germany). Moreover, because a multibracket HA was used, premolar brackets were placed on the stainless-steel crowns on the upper first premolar teeth, and Roth prescription 0.018" brackets (Mini Master Series, American Orthodontics, Sheboygan, WI, USA) were placed on the upper anterior teeth (Figure 1). Orthodontic leveling and alignment was performed up to 0.016" × 0.022" stainless-steel (SS) wire. Sagittal activation was performed in the first appointment in which the HA was applied,




Figure 1- Intraoral images (A-Profile view, B-Frontal view) of the appliance applied in the Herbst appliance group.

and activations after the first 4-5 mm of forward movement of the mandible were conducted by adding shims until normal overjet was achieved.²⁹ After the HA was removed, all remaining teeth were bonded, and 3D stereophotogrammetric recordings were taken before the HA was applied (T0) and immediately after debonding (T1). As a result of fixed functional and subsequent orthodontic treatment, class I occlusion was achieved in molar and canine relations, normal overjet values, and good interdigitation, then all devices were removed.

Orthognathic surgery group

This group included patients who underwent mandibular advancement alone with BSSO without genioplasty. BSSO was performed by the same experienced surgical team according to Hunsuck's modification.¹⁵ The same surgical and anesthesia techniques were performed for all patients.

Pre- and post-surgical orthodontic procedure

For all patients, 0.018" slot Roth prescription brackets (Mini Master Series, American Orthodontics, Sheboygan, WI, USA) with straight wire technique were used. Treatment was initiated with 0.012" or 0.014" Nickel-titanium (NiTi) wires for alignment and leveling, and 0.016", 0.016"x0.022" NiTi, and

0.016"x0.022" Stainless Steel (SS) arch-wires were used for incisor decompensation and arch coordination, respectively. Patients were seen for four to six weeks. In all cases, a final surgical splint produced conventionally or with a virtual surgery planning software (Nemo FAB, Nemotec, Leganés, Madrid, Spain) was used to ensure the planned occlusion. The pre-surgical orthodontic treatment phase was ten to 12 months. Surgical arch wires with crimpable hooks were placed in the mouth one day before the surgery and replaced with 0.016"x0.022" NiTi arch-wire one month after the surgery.

Post-surgical orthodontics is necessary to perform decompensation or minor dental movements that cannot be performed because of the iatrogenic origin or disto-occlusion or mastication function and force.³⁰ The post-surgical orthodontic treatment is focused on finishing and detailing, so after achieving Class I occlusion, normal overjet values, and good interdigitation, the treatment was finished, and the orthodontic devices were removed.

Three-dimensional stereophotogrammetric recordings were taken immediately before surgery (T0) and immediately after debonding (T1), which was performed at least six months after surgery to ensure complete resolution of edema and soft tissue adaptation to BSSO.

Surgical procedure

Surgical procedures were performed according to the procedure described below.³¹ The same team performed the operations under general anesthesia. Bilateral inferior alveolar nerve block and local infiltrating anesthesia were applied. A mucosal incision was made with the electrocautery, and the mucoperiosteal flap was lifted to expose the medial and the lateral surface of the ramus and the lower border. Medial horizontal osteotomy was performed with ultrasonic devices just above the lingula and extending behind the lingula. Oblique osteotomy was performed starting from the anterior of the medial osteotomy to the mesial part of the second molar tooth. Vertical osteotomy was extended starting from the anterior border of the oblique osteotomy to the lingual surface of the lower border. Osteotomy lines related to chisel, and the split was performed with a border separator and bone spreader. The mandible was moved to the new position with the orthodontic surgical splint. The proximal and distal segments were fixed with mini plates (KLS Martin, Tuttlingen, Germany) and mini screws (KLS Martin, Tuttlingen, Germany). The mucosa was sutured with 4-0 Vicryl. The patient was extubated and transferred to the postoperative care unit.

Three-dimensional stereophotogrammetric records

Three-dimensional stereophotogrammetric records of all patients of this study were obtained with a 3dMDface™ (3dMD Ltd, Atlanta, GA, USA) imaging system in a private room in the orthodontics

department. Prior to photography, patients were instructed to remove their glasses and any jewelry, and men were required to shave their beards and/or mustaches. Images were taken with the teeth in slight occlusion, eyes open, and lips relaxed and together (Figure 2 and Figure 3). Care was taken to avoid any distortions or artifacts in the image recordings.

Outcomes

The primary outcome was the difference in facial soft tissue changes—including the mandibular area (pogonion, gonion, and lower lip) as well as other regions, such as the upper lip, nose, and maxilla—between the HA and OS groups, as measured by 3D facial images taken at T0 and T1. Secondary outcomes were skeletal and dental changes, along with pharyngeal airway dimension—specifically the minimum posterior airway space (PASmin)—assessed via lateral cephalometric radiographs.

Images taken at T0 and T1 were evaluated with 3dMDvultus™ (3dMD Ltd, Atlanta, GA, USA) software. These images were manually aligned in the same software program and automatically registered by marking the broad surface of the forehead, the surface from the root of the nose to the hump of the nose, and the lateral parts of the bilateral Ex.³² The root mean square (RMS) value calculated for the acceptability of registration was below the recommended value of 0.5³³ (RMS: 0.09–0.49, mean value: 0.30±0.11).

Figures 4 and 5 show, respectively, the definitions and abbreviations of the anthropometric landmarks³⁴ and the measurements used in the study. For bilateral




Figure 2- The 3dMD images showing the three-dimensional view before (A) and after treatment (B) of a patient who underwent mandibular advancement surgery.



Figure 3- The 3dMD images showing the three-dimensional view before (A) and after treatment (B) of a patient who underwent Herbst appliance therapy.

Soft tissue landmarks	Abb	Definition
glabella	g	The most anterior midpoint on the fronto-orbital soft tissue contour.
nasion	na	The point in the midline of both the nasal root and the frontonasal suture.
endocanthion	en	The soft tissue point at the inner commissure of each eye fissure.
pronasale	prn	The most prominent point on the nasal tip.
columella constructed	cc	The midpoint of the columella crest at the level of the nostril top points.
tragion	tr	The point at the upper margin of each tragus.
gonion	go	The most lateral point on the soft tissue contour of each mandibular angle.
alare	al	The most lateral point on each alar contour.
alare curvature	ac	The point at the facial insertion of each alar base.
subnasale	sn	The midpoint on the nasolabial soft tissue contour between the columella crest and the upper lip.
crista philtrum	cph	The points at the intersection of the vermilion border and the elevated margin of the right and left philtrum.
subspinale	ss	The most posterior midpoint of the philtrum.
cheilion	ch	The point at each labial commissure.
labrale superior	ls	The midpoint of the vermilion line of the upper lip .
labrale inferior	li	The midpoint of the vermilion line of the lower lip .
stomion	sto	The midpoint between the lower border of the upper lip and upper border of the lower lip.
sublabiale	sl	The most posterior midpoint on the labiomental soft tissue contour that defines the border between the lower lip and the chin.
pogonion	pog	The most anterior midpoint of the chin.
menton	me	The lowest point on the midline of the chin.

Abb: Abbreviations

Figure 4- Definitions of soft tissue landmarks and reference planes.

points, the letter 'r' is added on the right, and 'l' is added on the left. After the landmarking process of the T0 images was completed, all points were projected onto the superimposed T1 image. The points that were not affected by the surgery (G, N, En_r, and En_l) were left as they were; All other points were repositioned on the T1 images, and the 3D changes (in the horizontal (x), vertical (y), and

sagittal (z) directions) of the points at T0 and T1 were compared.

The measurements marked with an asterisk were measured separately for the right and left sides for all patients, and no statistically significant differences were detected between the measurements on either side. Therefore, right-side measurements are provided.

3D stereophotogrammetric measurements	
*Mandibular Length (mm)	The line extending from the tragion_r to the pogonion
*Mandibular Corpus Length (mm)	The line extending from the gonion_r to the pogonion
Maxillary Height=Upper Lip Height (mm)	The line extending from the subnasale to the stomion
Anterior Facial Height (mm)	The line extending from the nasion to the menton
Lower Facial Width=Bigonial Width (mm)	The line extending from the gonion_r to the gonion_l
Lower Face Height (mm)	The line extending from subnasale to menton
Nasal Ala Width (mm)	The line extending from the alar_r to the alar_l
Alar Base Width (mm)	The line extending from the alar curvature_r to the alar curvature_l
Mandibular Height (mm)	The line extending from the stomion to the menton
*Posterior Facial Height (mm)	The line extending from the tragion_r to the gonion_r
Upper 1/3 \ Lower1/3	The ratio of the distance between the glabella and subnasale to the distance between the subnasale and menton
A Angle (°)	The angle formed between the tragion_r, nasion, and subnasale
B Angle (°)	The angle formed between the tragion_r, nasion, and sublabiale
ANB Angle (°)	The angle formed between the subnasale, nasion, and sublabiale
Mandibular Angle (°)	The angle formed between the tragion_r, pogonion, and tragus_l
Mandibular Convexity Angle (°)	The angle formed between the gonion_r, pogonion, and gonion_l
Facial Convexity Angle (°)	The angle formed between the nasion, subnasale, and pogonion
*Vertical Angle (°)	The angle formed between the nasion, gonion_r, and menton
Nasolabial Angle (°)	The angle formed between the columella constructed, subnasale, and labiale superius
Mentolabial Angle (°)	The angle formed between the labiale inferius, sublabiale, and pogonion
*Gonial Angle (°)	The angle formed between the tragion_r, gonion_r, and menton
Cephalometric measurements	
SNA (°)	Angle between Sella-Nasion line and A-point; reflects maxillary position
SNB (°)	Angle between Sella-Nasion line and B-point; reflects mandibular position
ANB (°)	Difference between SNA and SNB; indicates skeletal sagittal discrepancy
SN/GoGn (°)	Mandibular plane angle relative to anterior cranial base
U1/PP (°)	Inclination of the upper incisor to the palatal plane
IMPA (°)	Inclination of the lower incisor to the mandibular plane
SN/PP (°)	Angle between the anterior cranial base and palatal plane
A-NA perp (mm)	Linear distance from A-point to the N-A perpendicular line
Pg-NA perp (mm)	Linear distance from Pogonion to the N-A perpendicular line
Occ/SN (°)	Occlusal plane angle relative to the cranial base
OJ (mm)	Overjet; horizontal overlap between upper and lower incisors
OB (mm)	Overbite; vertical overlap between upper and lower incisors
<i>Airway Measurement</i>	
Sagittal Oropharyngeal Length (mm) **	The anteroposterior soft tissue distance between the posterior median pharyngeal wall (MPW) and the tongue base at the level of the Second Cervical Vertebra (Cv2'), measured parallel to the Frankfort horizontal plane, is defined as Cv2'–MPW.

Sagittal Oropharyngeal Length (mm) ** The anteroposterior soft tissue distance between the posterior median pharyngeal wall (MPW) and the tongue base at the level of the Second Cervical Vertebra (Cv2'), measured parallel to the Frankfort horizontal plane, is defined as Cv2'–MPW.

* Since there is no statistical difference between the measurements evaluated separately for the right and left sides, right side measurements are given for the relevant measurements.

** Reference 4

Figure 5 - Measurements and their explanations.

The changes in the 17 anthropometric landmark coordinate changes and the then linear, ten angular, and one rational measurements (Figure 5, Table 1 and 2) were evaluated. Additionally, eight angular

and four linear cephalometric parameters were analyzed, along with one linear airway measurement, to provide a comprehensive assessment of skeletal and functional changes (Table 5).

Statistical analysis

The data obtained in the study were recorded using Microsoft Office Excel (Microsoft Office 365, Microsoft, Redmond, Washington, USA) software and statistically analyzed with Statistical Package of the Social Sciences (v. 24.0, IBM Inc., Armonk, NY, USA) software. The Shapiro-Wilk test was used for normality analysis, and Levene's test was used for homogeneity analysis. All data were normally and homogeneously distributed; therefore, comparisons before and after treatment were made using paired samples t-tests, and comparisons between groups were made using independent samples t-tests. Data are shown as the mean and standard deviation, and statistical significance was defined as $p < 0.05$.

Method error

The same researcher (ETH) re-analyzed the data from ten randomly selected patients one month after the first measurement. The intraclass correlation coefficient was used to examine the reliability between the first and second measurements. Cronbach's alpha was 0.978 (range: 0.962-0.989), indicating sufficient reliability.

Results

All subjects in both treatment groups were treated to a Class I dental arch relationship. The HA treatment time was seven to eight months, and the total treatment time was 19.8 months. The average body mass index (BMI) of the patients in the HA group was 20.48 at the pre- and 20.8 at post-treatment. The total treatment time was 24.2 months in the OS group, with a post-surgical phase lasting approximately six months. The mean BMI in this group was 21.9 at the pre- and 21.2 at post-treatment.

Table 1 shows the comparison of pre-treatment values between groups. None of the parameters differed between the groups at the beginning of treatment.

Table 2 shows the between-group comparison of the changes that occurred after treatment. In the HA group, the lower facial width ($p = 0.008$), mandibular angle ($p = 0.027$), mandibular convexity angle ($p = 0.006$), ANB angle ($p < 0.001$), and A angle ($p = 0.039$) decreased significantly after treatment, whereas the mandibular corpus length ($p < 0.001$),

Table 1- Pre-treatment values of the Herbst Appliance (HA) and Orthognathic Surgery (OS) Patients.

Measurements	HA	OS	P values*
Mandibular Length (mm)	131.46±7.24	129.58±7.56	0.483
Mandibular Corpus Length (mm)	98.13±7.0	95.21±6.50	0.240
Maxillary Height=Upper Lip Height (mm)	21.17±1.92	21.25±2.92	0.916
Anterior Facial Height (mm)	115.99±5.27	116.40±10.55	0.892
Lower Facial Width=Bigonial Width (mm)	114.18±12.26	110.31±8.73	0.319
Lower Face Height (mm)	68.76±5.56	68.40±9.10	0.895
Nasal Ala Width (mm)	34.59±4.43	32.68±3.51	0.195
Alar Base Width (mm)	28.48±3.96	27.61±2.52	0.477
Mandibular Height (mm)	48.32±4.49	48.06±8.78	0.921
Posterior Facial Height (mm)	55.65±19.94	53.04±8.81	0.646
Upper 1/3 \ Lower1/3	0.98±0.11	0.99±0.11	0.758
A Angle (°)	80.79±2.15	80.18±3.59	0.571
B Angle (°)	72.88±2.14	71.47±3.45	0.179
ANB Angle (°)	13.38±2.99	13.08±1.69	0.726
Mandibular Angle (°)	66.95±2.57	68.30±3.50	0.236
Mandibular Convexity Angle (°)	71.23±4.65	71.27±5.92	0.981
Facial Convexity Angle (°)	155.33±7.65	155.32±4.74	0.992
Vertical Angle (°)	62.99±2.68	62.93±5.80	0.973
Nasolabial Angle (°)	118.52±7.95	112.02±10.97	0.068
Mentolabial Angle (°)	114.52±18.69	120.93±17.22	0.329
Gonial Angle (°)	135.27±5.01	134.04±6.21	0.548

Data was given mean ± standard deviation. T0: Pre-treatment. T1: Post-treatment. * Results of Independent Samples- t- test. Statistically significant value: $p < 0.05$.

anterior facial height ($p=0.019$), posterior facial height ($p=0.013$), B angle ($p=0.001$), vertical angle ($p=0.028$), and mentolabial angle ($p<0.001$) increased significantly after treatment.

In the OS group, mandibular length ($p<0.001$), mandibular corpus length ($p<0.001$), facial convexity angle ($p<0.001$), posterior facial height ($p<0.001$), B angle ($p<0.001$), vertical angle ($p=0.007$), and mentolabial angle ($p=0.001$) increased significantly after treatment, whereas the mandibular angle ($p<0.001$) and ANB angle ($p<0.001$) decreased significantly after treatment.

Table 3 shows the intergroup comparison of changes after treatment. In the OS group, the mandibular length (OS: 4.07 ± 2.54 ; HA: 0.92 ± 2.12 ; $p=0.001$), mandibular corpus length (OS: 4.94 ± 2.51 ; HA: 2.63 ± 1.46 ; $p=0.005$), and facial convexity angle (OS: 5.10 ± 3.29 ; HA: 1.51 ± 2.96 ; $p=0.003$) increased significantly, and the increase was greater than that in the HA group. Moreover, a significant difference was observed in alar base width between the HA group and the OS group (HA: 0.37 ± 0.75 ; OS: -0.12 ± 0.47 ; $p=0.039$).

Table 4 shows the changes in the coordinates of anthropometric landmarks after treatment and the comparison of these changes between groups. The only point with a significant change in coordinates between groups was the pogonion; in the OS group, this point moved significantly more forward (3.82 ± 2.64 ; $p=0.001$) than the same point in the HA group (PG_z: 0.78 ± 2.13).

Table 5 shows cephalometric comparisons of pre- and post-treatment changes. No statistically significant differences were observed between the HA and OS groups in any cephalometric parameter ($p>0.05$) at pre-treatment (T0). The HA group showed significant increases in SNB angle ($p=0.002$), IMPA ($p=0.001$), SN/GoGn angle ($p=0.032$), Pg-NA perpendicular distance ($p=0.006$), and airway dimension ($p<0.001$) at post-treatment. Additionally, significant reductions were noted in ANB angle ($p<0.001$), overjet ($p=0.003$), and overbite ($p<0.001$). SNB angle ($p<0.001$), Pg-NA perpendicular distance ($p<0.001$), SN/GoGn angle ($p=0.007$), and airway dimension ($p<0.001$) increased significantly after treatment in the OS

Table 2- Pre- and post-treatment comparisons of the Herbst Appliance (HA) and Orthognathic Surgery (OS) Patients.

Measurements	HA			OS		
	T0	T1	P values*	T0	T1	P values*
Mandibular Length (mm)	131.46±7.24	132.38±7.44	0.105	129.58±7.56	133.65±7.32	<0.001
Mandibular Corpus Length (mm)	98.13±7.0	100.74±6.77	<0.001	95.21±6.50	100.16±6.88	<0.001
Maxillary Height=Upper Lip Height (mm)	21.17±1.92	21.33±2.14	0.638	21.25±2.92	20.86±2.52	0.246
Anterior Facial Height (mm)	115.99±5.27	117.39±5.86	0.019	116.40±10.55	118.08±8.62	0.136
Lower Facial Width=Bigonial Width (mm)	114.18±12.26	110.33±10.97	0.008	110.31±8.73	109.46±9.06	0.440
Lower Face Height (mm)	68.76±5.56	69.52±6.06	0.348	68.40±9.10	69.35±6.87	0.476
Nasal Ala Width (mm)	34.59±4.43	34.50±4.33	0.497	32.68±3.51	32.55±3.34	0.406
Alar Base Width (mm)	28.48±3.96	28.85±3.96	0.067	27.61±2.52	27.49±2.39	0.338
Mandibular Height (mm)	48.32±4.49	48.83±5.46	0.493	48.06±8.78	49.58±7.32	0.268
Posterior Facial Height (mm)	55.65±19.94	58.88±19.81	0.013	53.04±8.81	55.93±9.18	<0.001
Upper 1/3 \ Lower1/3	0.98±0.11	0.97±0.10	0.422	0.99±0.11	0.98±0.10	0.403
A Angle (°)	80.79±2.15	79.88±2.44	0.039	80.18±3.59	79.86±3.72	0.381
B Angle (°)	72.88±2.14	74.17±2.24	0.001	71.47±3.45	73.72±3.95	<0.001
ANB Angle (°)	13.38±2.99	11.53±2.89	<0.001	13.08±1.69	10.14±2.73	<0.001
Mandibular Angle (°)	66.95±2.57	65.96±1.53	0.027	68.30±3.50	66.25±3.38	<0.001
Mandibular Convexity Angle (°)	71.23±4.65	69.71±3.41	0.006	71.27±5.92	70.14±6.04	0.132
Facial Convexity Angle (°)	155.33±7.65	156.84±6.62	0.060	155.32±4.74	160.41±5.93	<0.001
Vertical Angle (°)	62.99±2.68	64.10±3.34	0.028	62.93±5.80	64.34±5.11	0.007
Nasolabial Angle (°)	118.52±7.95	119.90±7.95	0.327	112.02±10.97	111.94±13.53	0.964
Mentolabial Angle (°)	114.52±18.69	132.23±23.67	<0.001	120.93±17.22	134.33±14.71	0.001
Gonial Angle (°)	135.27±5.01	134.77±3.93	0.723	134.04±6.21	134.82±6.32	0.318

Data was given mean ± standard deviation. T0: Pre-treatment. T1: Post-treatment. * Results of Paired Samples -t- test. Statistically significant value: $p<0.05$.

Table 3- Comparison of treatment changes in the Herbst Appliance (HA) and Orthognathic Surgery (OS) Patients.

Measurements	HA	OS	P values*
Overjet Change (mm)	7.41±2.48	7.90±2.97	0.638
Mandibular Length (mm)	0.92±2.12	4.07±2.54	0.001
Mandibular Corpus Length (mm)	2.63±1.46	4.94±2.51	0.005
Maxillary Height=Upper Lip Height (mm)	0.17±1.40	-0.40±1.26	0.251
Anterior Facial Height (mm)	1.40±2.13	1.68±4.12	0.811
Lower Facial Width=Bigonial Width (mm)	-3.85±4.99	-0.85±4.16	0.081
Lower Face Height (mm)	0.76±3.12	0.95±5.00	0.899
Nasal Ala Width (mm)	-0.09±0.51	-0.13±0.56	0.853
Alar Base Width (mm)	0.37±0.75	-0.12±0.47	0.039
Mandibular Height (mm)	0.51±2.89	1.52±5.12	0.497
Posterior Facial Height (mm)	3.24±4.60	2.89±2.37	0.796
Upper 1/3 \ Lower1/3	-0.01±0.05	-0.02±0.07	0.760
A Angle (°)	-0.91±1.60	-0.31±1.35	0.277
B Angle (°)	1.29±1.31	2.25±1.71	0.090
ANB Angle (°)	-1.85±1.00	-2.93±2.13	0.079
Mandibular Angle (°)	-0.99±1.63	-2.05±1.69	0.086
Mandibular Convexity Angle (°)	-1.51±1.92	-1.13±2.73	0.652
Facial Convexity Angle (°)	1.51±2.96	5.10±3.29	0.003
Vertical Angle (°)	1.11±1.82	1.41±2.82	0.727
Nasolabial Angle (°)	1.38±5.43	-0.07±5.97	0.485
Mentolabial Angle (°)	17.72±13.51	13.40±13.14	0.375
Gonial Angle (°)	-0.50±2.76	0.78±2.92	0.220

Data was given mean ± standard deviation. * Results of Independent Samples -t- test. Statistically significant value: $p < 0.05$.

group. ANB angle ($p < 0.001$), overjet ($p = 0.002$), and overbite ($p = 0.010$) also demonstrated statistically significant reductions. When the magnitude of change was compared between the groups, the OS group showed a significantly greater increase in SNB angle ($p = 0.030$) and Pg-NA perpendicular distance ($p = 0.003$). No significant intergroup differences were found for the change in overjet, overbite, ANB angle, or other variables ($p > 0.05$).

Discussion

In this study, the effects of OS and one of its most common alternatives—the HA (a fixed functional device)³⁵—on soft tissues were compared via 3D stereophotogrammetric recordings in patients with skeletal class II malocclusion with mandibular deficiency. All participants had completed growth and development, as determined by hand and wrist X-rays.

The choice of treatment for adult patients with moderate skeletal class II malocclusion varies

depending on the attitudes and behaviors of the patients and the approach of the physician. Physician-related preferences were eliminated because all patients attended the same center. Although previous research has reported that OS provides superior skeletal correction,^{15,19} some patients may refuse surgery. No differences in the initial values were observed between groups in this study, indicating that the patients treated with HA did not have meaningfully different characteristics from patients who preferred OS, who did not demand OS. Therefore, the results of two different treatment methods could be clearly monitored because the patients were similar.

Stereophotogrammetry is very useful in recording the complex 3D morphology and measuring dimensions of the human face in realistic dimensions³⁶ and may cause software errors, requires frequent calibration (every 15 days), is expensive, and cannot evaluate hard tissue without X-ray. Various strategies have been employed for the superimposition of consecutive 3D facial images in literature. For instance, Day and Lee³⁷ (2006) based their registration solely on the broad surface of the forehead. Soncul and Bamber³⁸ (2004) utilized a five-point method involving bilateral

Table 4- Comparison of 3D coordination changes of anthropometric landmarks in the Herbst Appliance (HA) and Orthognathic Surgery (OS) Patients.

Measurements	HA	OS	P values*
G_z	-0.16±0.69	-0.02±0.34	0.502
N_z	-0.25±0.66	-0.13±0.48	0.563
EN_R_z	-0.34±0.68	0.28±0.61	0.283
EN_L_z	-0.19±0.99	0.15±0.44	0.230
AC_R_x	-0.03±0.3	0.12±0.17	0.106
AC_R_z	0.31±1.2	-0.4±1.15	0.103
AC_L_x	0.25±0.6	-0.04±0.28	0.093
AC_L_z	0.57±1.38	-0.07±0.84	0.432
AL_R_x	0.19±0.49	0.06±0.26	0.344
AL_R_z	-0.21±1.09	-0.2±1.13	0.962
AL_L_x	0.05±0.4	-0.08±0.39	0.372
AL_L_z	-0.53±2.28	-0.12±1.19	0.540
Go_R_x	2.21±3.18	0.26±2.17	0.056
Go_R_y	-2.95±3.87	-2.23±2.1	0.523
Go_R_z	0.98±1.92	2.58±2.88	0.078
Go_L_x	-1.64±2.31	-0.6±2.36	0.225
Go_L_y	-2.96±3.87	-2.38±2.12	0.608
Go_L_z	1.09±1.98	2.86±3.48	0.089
CH_R_y	-0.45±1.14	-0.1±1.22	0.420
CH_R_z	0.11±2.51	1.09±2.17	0.258
CH_L_y	-0.31±1.26	-0.07±1.03	0.563
CH_L_z	0.19±2.69	0.73±2.03	0.537
CPH_R_y	0.07±0.28	0.03±0.52	0.807
CPH_R_z	-0.87±1.81	-0.43±1.6	0.473
CPH_L_y	0.12±0.5	-0.08±1.04	0.479
CPH_L_z	-0.84±2.03	-0.42±1.63	0.535
PRN_z	-0.28±0.89	-0.19±0.92	0.790
SN_z	-0.1±1.49	-0.21±0.95	0.810
CC_z	0.34±1.37	0.01±0.62	0.399
SS_y	-0.32±0.68	-0.05±0.38	0.075
SS_z	-0.59±1.87	-0.52±1.93	0.928
LS_y	0.01±0.88	0.33±1.41	0.448
LS_z	-0.69±2.08	-0.38±1.95	0.667
STO_y	-0.38±1.21	0.17±0.93	0.168
STO_z	0.5±2.4	1.46±3.17	0.346
Li_y	-0.46±1.45	-0.19±2.15	0.689
Li_z	1.29±2.28	2.35±3	0.274
SL_y	-2.19±2.35	-1.89±2.08	0.703
SL_z	2.75±2	3.92±3.39	0.245
PG_y	-1.86±3.41	-2.05±2.03	0.855
PG_z	0.78±2.13	3.82±2.64	0.001

Data was given mean ± standard deviation. * Results of Independent Samples -t- test. Statistically significant value: $p < 0.05$. x-axis (horizontal), left (no sign), right (minus sign); y-axis (vertical), superior (no sign), inferior (minus sign); z-axis (sagittal), anterior (no sign), posterior (minus sign).

Parameters are defined in Figure 4

All measurements are given in mm.

En, Ex, and Na landmarks. McCance, et al.³⁹ (1992), on the other hand, combined the standard five-point approach with additional randomly distributed points across the forehead.

Nevertheless, the precision of this process can be compromised by minor inconsistencies in landmark

placement. Specifically, since bilateral En and Ex points tend to lie nearly along the same horizontal line, relying solely on these landmarks may lead to inaccuracies in alignment.⁴⁰ Prior research has also indicated that using only the wide forehead region—especially in individuals with broad and flat forehead

Table 5- Comparison of cephalometric changes with treatment in the Herbst Appliance (HA) and Orthognathic Surgery (OS) Patients.

	HA			OS			Comparisons of T0	Differences with Treatment		
	T0	T1	P values*	T0	T1	P values*		P values**	HA	OS
SNA°	81.47±3.73	81.41±3.72	0.780	80.31±1.67	80.56±1.66	0.373	0.414	-0.06±0.58	0.24±0.78	0.366
SNB°	75.29±3.97	76.99±3.91	0.002	72.48±2.53	75.77±2.68	<0.001	0.096	1.7±1.14	3.29±1.62	0.030
ANB°	6.19±2.33	4.42±2.55	<0.001	7.82±1.16	4.77±2.24	<0.001	0.084	-1.77±0.69	-3.06±1.8	0.062
SN/GoGn°	30.56±8.37	31.57±7.65	0.032	34.1±7.83	34.77±5.69	0.562	0.368	1.01±1.17	0.67±3.31	0.772
U1/PP°	109.68±9.26	113.22±6.84	0.234	109.07±8.96	112.04±6.81	0.386	0.889	3.54±8.25	2.98±9.74	0.896
IMPA°	94.04±8.07	99.88±6.77	0.001	94.22±9.96	95.92±8.59	0.357	0.967	5.83±3.49	1.7±5.22	0.066
SN/PP°	9.62±3.27	9.6±3	0.960	8.17±4.27	6.48±3.02	0.064	0.429	-0.02±1.29	-1.69±2.36	0.087
A-NA perp (mm)	-0.68±3.82	-0.76±3.78	0.669	-2.21±1.8	-1.86±2.06	0.290	0.299	-0.08±0.53	0.36±0.94	0.250
Pg-NA perp (mm)	-9.49±6.68	-7.3±6.31	0.006	-15.14±5.69	-9.62±5.66	<0.001	0.072	2.19±1.79	5.52±2.24	0.003
Occ/SN°	15.58±4.6	17.33±4.41	0.133	14.87±3.86	16.66±3.6	0.182	0.727	1.76±3.15	1.79±3.67	0.983
OJ (mm)	6.53±1.45	3.08±2.48	0.003	8.81±2.89	3.68±1.14	0.002	0.057	-3.46±2.51	-5.13±3.31	0.243
OB (mm)	5.07±2.05	1.86±1.56	<0.001	4.99±3.27	2.16±1.27	0.010	0.953	-3.21±1.39	-2.83±2.53	0.700
Airway (mm)	4.31±1.71	5.64±1.84	<0.001	4.27±1.34	5.47±1.34	<0.001	0.952	1.33±0.58	1.2±0.71	0.683

PP: palatal plane, Occ: Occlusal line, OJ: overjet, OB: overbite. Data was given as mean ± standard deviation. T0: Pre-treatment. T1: Post-treatment. * Results of Paired Samples t-test. ** Results of the Independent samples t-test. Statistically significant value: p<0.05.

contours, as often seen in some Asian populations—can result in lateral deviations of the superimposed images.⁴⁰

To minimize such errors, this study employed a more stable composite registration area, incorporating not only the wide forehead surface but also the nasal bridge and the lateral extensions of the bilateral Ex points, as previously described.^{32,41}

In this study, HA was applied with gradual activation, which provides better bone remodeling in the glenoid fossa and condyle, thus showing more skeletal improvement.⁴² To yield a response in functional treatment, progress was started with 4 mm of activation, which is the minimum amount recommended to exceed the minimum tension threshold.⁴³ Although stepwise mandibular advancement has been postulated to yield superior skeletal adaptations relative to single-step activation, current evidence remains inconclusive.²⁹ While certain investigations have demonstrated enhanced mandibular advancement and attenuated dentoalveolar compensations with gradual activation, other studies report no significant differences in skeletal and dental outcomes between the two approaches, aside from increased proclination and anterior displacement of the mandibular incisors in the stepwise group.²⁹ Furthermore, the adjunctive application of extraoral appliances—particularly high-pull headgear—has been proposed to augment

vertical control and mitigate undesired dentoalveolar effects.⁴⁴ Collectively, these findings suggest that therapeutic response is likely influenced by a complex interplay of factors, including individual growth patterns, appliance configuration, and activation mechanics.

HA is independent of patient cooperation, and transversal expansion can be performed together with sagittal activation and provide skeletal corrections by increasing the total mandibular length.⁴⁵ The mandibular corpus length increased significantly in this study, however, complications such as crown fracture or separation, screw loosening, rod distortion, pivot fracture, and soft tissue injuries may be found in HA treatment.¹⁴ Moreover, the device should be used with extreme caution in patients who are not growing, whose skeletal changes are minimal, and treatment effects are limited to the dentoalveolar area. There is also an increased risk for the development of 'dual bite' with symptoms of dysfunction from the TMJ as a possible consequence.¹⁸ Moreover, the changes seen with the use of the HA in adults may simply be the result of "mandibular repositioning"⁴⁶ and the proclination of the lower incisors.⁴⁷

The first use of BSSO procedures for the correction of mandibular deformities was described in 1953.⁴⁸ The most common problem in patients with OS treatment is cranial nerve damage and sensitivity change (during the BSSO, the risk of damage to the

inferior alveolar nerve of the mandibular branch of the trigeminal nerve is quite high).⁴⁹ These patients also experience a loss of function, such as drooling, unnoticeable food particles remaining on the lips and/or chin, errors in pronunciation, pain, cheek biting, eating, and kissing.⁵⁰

Temporomandibular joint problems, ranging from intra-articular noise to pain and from limitations in mouth opening to condylar resorption, are the second most common problems of OS.⁴⁸ Bleeding, auditory tube function and hearing problems, infections, and bad split were reported as other possible side effects.⁵¹ Also, the cost of OS is approximately five to seven times higher than that of orthodontic treatment alone.⁵¹

A meta-analysis examining the skeletal effect of HA⁵² reported that the SNA angle changed by -0.56 degrees, the SNB angle changed by 1.06 degrees, and the ANB changed by -1.08 degrees. These results are quite close to the relevant values in the HA group in this study. There were no differences when the parameters mentioned were compared with those in the OS group, however, the values in this study were for soft tissues.

Although the meta-analysis revealed that the hard tissue Co-Gn length changed significantly by 1.74 mm, the same authors reported that this small amount may not be clinically significant.⁵² Thus, although the values were like those of the mandibular length and mandibular corpus length in the HA group in this study, these lengths increased significantly in the OS group.

In this study, the anterior facial height increased by 1.5 mm, the posterior facial height increased by 3 mm, and the vertical angle increased by one degree after treatment in the HA group. Although these values differed significantly within the group, they did not differ significantly compared with the changes in the OS group. These results are consistent with the results of the meta-analysis, which reported only a 0.17 -degree increase in the mandibular plane angle following HA treatment and that HA had little effect on the mandibular plane angle in the clinic.⁵² Notably, the meta-analysis was conducted on cephalometric studies, whereas this study evaluated soft tissues. The increase in the posterior facial heights in both groups in this study was almost the same as the increase in the S-Go height (3.4 mm) in a cephalometric study.⁵³ Many other studies reported that, as well as

this study, neither the anterior and posterior facial heights nor the mandibular plane angle (SN-GoMe) were significantly affected.⁵³ Conversely, one study reported a decrease in posterior facial height with OS (2 mm), which the authors attributed to remodeling and reformation during post-operative healing.¹⁵

A study in which only mandibular advancement was evaluated with 3D methods⁵⁴ found that the mentolabial fold tightened after surgery, which is consistent with this study, and the mentolabial angle increased similarly in both groups. This may be attributed to the fact that the lower lip is freed from being trapped between the teeth because of the decrease in overjet with treatment.

In this study, no change in the gonial angle was observed, similar to another study, in which no significant change was noted in the Co-Go-Me angle.⁵⁵ We can conclude that both protocols (HA and OS) can be safely applied in all patients with class II skeletal malocclusion in whom the gonial angle is acceptable (between 128 and 115 degrees⁵⁶).

A cone-beam computed tomography (CBCT) study showed that the location of pogonion in the HA group was significantly different from that in the control group, it was located 1.7 mm anteriorly in the HA group. However, this study was conducted in younger patients and reported short-term results.⁵⁷ Other studies found that the pogonion point was located anteriorly after HA treatment in patients younger than those in this study^{53,58} but no differences were observed when the HA group was compared with the control group. There was a similar result in the HA group in this study, but the pogonion showed a clear average increase in OS of 3 mm more forward in the anteroposterior direction (HA 0.78 mm and OS 3.82 mm). This is very similar to the results of a cephalometric evaluation study that compared the effects of HA and OS (HA: 1.28 mm; OS 4.05 mm).¹⁵ The same study also observed that the increase in facial convexity angle measured from the soft tissue nasion-subnasale and soft tissue pogonion in the OS group (5.45 degrees) was almost the same as this study (5.10 degrees). The most profound differences between the OS and HA groups were greater mandibular base advancement (mandibular length, corpus length, facial convexity angle, and Pg_z), which led to greater reductions in soft tissue profile convexities. The forward movement of the pogonion indicates that skeletal movements were more active

in improving the facial profile and correcting increased overjet in the OS group. In the HA group, we attribute the slight expansion in the alar base width, which we consider clinically insignificant, to the activation of the hyrax screw used for expansion.

The lack of significant differences in baseline cephalometric parameters between the HA and OS groups confirms that the two cohorts were comparable at the outset, thereby strengthening the validity of the treatment effect comparisons. The cephalometric outcomes in this study agree with a previous study, showing greater skeletal advancement in the OS group, particularly in SNB and Pg-NA perpendicular measurements.¹⁵ As in their findings, data in this study revealed more pronounced mandibular advancement and ANB reduction in OS patients, while HA patients showed more dentoalveolar compensation. Regarding airway dimensions, the CV2'-MPW (sagittal oropharyngeal airway length) increased significantly within both groups, but no significant intergroup difference was detected. This supports previous evidence that both modalities may improve pharyngeal airway space, however, the magnitude of this change may not be solely dependent on the skeletal advancement technique.²⁹

Despite the complications and possible risks of OS treatment, many patients are satisfied with the results and recommend OS to those who need it.⁵⁹ Early treatment can also improve oral health and oral health-related quality of life.⁵¹ We agree that HA is a very reasonable and appropriate option when patients refuse surgery, or their main goal is not to improve their facial aesthetics.¹⁹

Limitations

This study has several limitations, including the relatively small sample size, the absence of a control group, the lack of long-term follow-up, and its retrospective design, which may reduce the strength of causal inferences. Furthermore, although a cephalometric skeletal analysis was included to enhance the interpretation of soft tissue changes, the evaluation was limited to 2D lateral cephalometric radiographs, which do not fully capture 3D skeletal dynamics. Three-dimensional hard tissue evaluations using cone-beam computed tomography (CBCT) were not performed, in accordance with the "as low as reasonably achievable" (ALARA) principle.

Radiographic imaging was strictly limited to clinically necessary situations, and additional radiation exposure for research purposes was avoided, consistent with internationally recognized ethical standards. This study mainly differed from previous studies in its evaluation of soft tissues. Although this approach is in line with today's aesthetic concept approach, it also has limitations, such as soft tissue thickness, muscle structures, and biological and individual differences.

Conclusions

Despite the limitations of the study, several key conclusions can be drawn. Patients in the HA and OS groups achieved class I occlusal relationships and reductions in the convexity of the soft tissues over the treatment period (T1-T0). However, more skeletal contribution was observed in the OS group than in the HA group because of the increase in mandibular length and mandibular corpus length. The only significant 3D soft tissue coordinate change between the groups was the anterior displacement of the pogonion in the sagittal plane, which was further corroborated by cephalometric findings showing significant increases in SNB angle and Pg-NA perpendicular measurements in the OS group. This anterior movement of the pogonion contributed to a reduction in soft tissue convexity, as reflected by the increased facial convexity angle. A clinically insignificant increase in alar base width was observed in the HA group. Additionally, both treatment modalities resulted in significant improvements in sagittal oropharyngeal airway length, without significant intergroup difference.

Ethics approval and consent to participate

Prior to this single-center clinical study, the necessary permissions were obtained from the Erciyes University Clinical Research Ethics Committee (Approval Date: 25.10.2023; Approval Number: 2023/695). The study was conducted ethically in accordance with the guidelines of the World Health Organization's Declaration of Helsinki. Potential complications were explained to the patients and their relatives, and informed consent forms were obtained.

Conflict of interest

The authors declare no conflicts of interest concerning this study.

Data availability statement

The datasets used and/or analyzed during this study are available from the corresponding author on reasonable request.

Authors' contributions

Çoban, Gökhan: Conceptualization (Equal); Data curation (Equal); Formal analysis (Equal); Investigation (Equal); Methodology (Equal); Project administration (Equal); Supervision (Equal); Validation (Equal); Visualization (Equal); Writing – original draft (Equal); Writing – review & editing. **Öztürk, Taner:** Conceptualization (Equal); Data curation (Equal); Formal analysis (Equal); Methodology (Equal); Validation (Equal); Visualization (Equal); Writing – original draft (Equal); Writing – review & editing. **Türker, Gökhan:** Conceptualization (Equal); Formal analysis (Equal); Visualization (Equal); Writing – original draft (Equal); Writing – review & editing. **Hashimli, Ebru Topcuoğlu:** Conceptualization (Equal); Data curation (Equal); Methodology (Equal); Validation (Equal); Writing – review & editing. **Kaba, Yusuf Nuri:** Data curation (Equal); Formal analysis (Equal); Investigation (Equal); Methodology (Equal); Validation (Equal); Writing – original draft (Equal); Writing – review & editing.

References

- 1- Bishara SE. Class II malocclusions: diagnostic and clinical considerations with and without treatment. *Semin Orthod.* 2006;12:11-24. doi: 10.1053/j.sodo.2005.10.005
- 2- El Chekie MR, Nemer G, Khalil A, Macari AT, Ghafari JG. Novel genes linked to Class II Division 1 malocclusion with mandibular micrognathism. *Am J Orthod Dentofacial Orthop* 2023;163:667-76. e3. doi: 10.1016/j.ajodo.2022.04.020
- 3- Kinzinger G, Frye L, Diedrich P. Class II treatment in adults: comparing camouflage orthodontics, dentofacial orthopedics and orthognathic surgery – a cephalometric study to evaluate various therapeutic effects. *J Orofac Orthop.* 2009;70:63-71. doi: 10.1007/s00056-009-0821-2
- 4- Çoban G, Gül Amuk N, Yağcı A, Akgün G, Abbood Abbood IH. Evaluation of external apical root resorption caused by fixed functional treatment of class II malocclusion: cast splint Herbst appliance vs. forsus fatigue resistant device. *J Orofac Orthop* 2023;84:50-9. doi: 10.1007/s00056-021-00334-x

- 5- Al-Dboush Re, Soltan R, Rao J, El-Bialy T. Skeletal and dental effects of Herbst appliance anchored with temporary anchorage devices: a systematic review with meta-analysis. *Orthod Craniofac Res.* 2022;25:31-48. doi: 10.1111/ocr.12510
- 6- Jena AK, Singh SP, Utreja AK. Sagittal mandibular development effects on the dimensions of the awake pharyngeal airway passage. *Angle Orthod.* 2010;80:1061-7. doi: 10.2319/030210-125.1
- 7- Liu Y, Zhao T, Ngan P, Qin D, Hua F, He H. The dental and craniofacial characteristics among children with obstructive sleep apnoea: a systematic review and meta-analysis. *Eur J Orthod.* 2023;45:346-55. doi: 10.1093/ejo/cjac074
- 8- Frilund E, Sonesson M, Magnusson A. Patient compliance with Twin Block appliance during treatment of Class II malocclusion: a randomized controlled trial on two check-up prescriptions. *Eur J Orthod* 2023;45:142-9. doi: 10.1093/ejo/cjac046
- 9- Uzuner FD, Aslan BI, Dinçer M. Dentoskeletal morphology in adults with Class I, Class II Division 1, or Class II Division 2 malocclusion with increased overbite. *Am J Orthod Dentofacial Orthop.* 2019;156:248-56. e2. doi: 10.1016/j.ajodo.2019.03.006
- 10- Masood M, Suominen AL, Pietila T, Lahti S. Malocclusion traits and oral health-related quality of life in Finnish adults. *Community Dent Oral Epidemiol* 2017;45:178-88. doi: 10.1111/cdoe.12276
- 11- Giuntini V, McNamara Jr JA, Franchi L. Treatment of Class II malocclusion in the growing patient: Early or late? *Semin Orthod* 2023;29:183-8. doi: 10.1053/j.sodo.2023.04.008
- 12- Pancherza H. Dentofacial orthopedics or orthognathic surgery: is it a matter of age? *Am J Orthod Dentofacial Orthop.* 2000;117:571-4. doi: 10.1016/s0889-5406(00)70203-4
- 13- Proffit WR, Phillips C, Douvartzidis N. A comparison of outcomes of orthodontic and surgical-orthodontic treatment of Class II malocclusion in adults. *Am J Orthod Dentofacial Orthop.* 1992;101:556-65. doi: 10.1016/0889-5406(92)70131-s
- 14- Silva JF, Gerszewski C, Moresca RC, Correr GM, Flores-Mir C, Moro A. Retrospective study of clinical complications during orthodontic treatment with either a removable mandibular acrylic splint Herbst or with a cantilever Herbst. *Angle Orthod.* 2015;85:64-71. doi: 10.2319/122113-936.1
- 15- Ruf S, Pancherz H. Orthognathic surgery and dentofacial orthopedics in adult Class II Division 1 treatment: mandibular sagittal split osteotomy versus Herbst appliance. *Am J Orthod Dentofacial Orthop.* 2004;126:140-52. doi: 10.1016/j.ajodo.2004.02.011
- 16- Amuk M, Gul Amuk N, Yılmaz S. Treatment and posttreatment effects of Herbst appliance therapy on trabecular structure of the mandible using fractal dimension analysis. *Eur J Orthod.* 2022;44:125-33. doi: 10.1093/ejo/cjab048
- 17- Ruf S, Pancherz H. Temporomandibular joint remodeling in adolescents and young adults during Herbst treatment: a prospective longitudinal magnetic resonance imaging and cephalometric radiographic investigation. *Am J Orthod Dentofacial Orthop.* 1999;115:607-18. doi: 10.1016/s0889-5406(99)70285-4
- 18- Pancherz H. The effects, limitations, and long-term dentofacial adaptations to treatment with the herbst appliance. *Semin Orthod.* 1997;3:232-43. doi: 10.1016/s1073-8746(97)80056-4
- 19- Chaiyongsirisern A, Rabie AB, Wong RW. Stepwise advancement Herbst appliance versus mandibular sagittal split osteotomy: treatment effects and long-term stability of adult Class II patients. *Angle Orthod.* 2009;79:1084-94. doi: 10.2319/110308-556r.1
- 20- Lai M, McNamara JA Jr. An evaluation of two-phase treatment with the Herbst appliance and preadjusted edgewise therapy. *Semin Orthod.* 1998;4:46-58. doi: 10.1016/s1073-8746(98)80041-8
- 21- Barnett GA, Higgins DW, Major PW, Flores-Mir C. Immediate skeletal and dentoalveolar effects of the crown-or banded type herbst appliance on class ii division 1 malocclusion: a systematic review. *Angle Orthod.* 2008;78:361-9. doi: 10.2319/031107-123.1

- 22- Te Veldhuis E, Te Veldhuis A, Bramer W, Wolvius E, Koudstaal M. The effect of orthognathic surgery on the temporomandibular joint and oral function: a systematic review. *Int J Oral Maxillofac Surg* 2017;46:554-63. doi: 10.1016/j.ijom.2017.01.004
- 23- Mulier D, Gaitán Romero L, Führer A, Martin C, Shujaat S, Shaheen E, et al. Long-term dental stability after orthognathic surgery: a systematic review. *Eur J Orthod*. 2021;43:104-12. doi: 10.1093/ejo/cjaa022
- 24- Singh P, Bornstein MM, Hsung RT, Ajmera DH, Leung YY, Gu M. Frontiers in three-dimensional surface imaging systems for 3D face acquisition in craniofacial research and practice: an updated literature review. *Diagnostics* 2024;14:423. doi: 10.3390/diagnostics14040423
- 25- Marino Merlo M, Quiroga Souki B, Nieri M, Bonanno A, Giuntini V, McNamara JA Jr, et al. Comparison of the effects on facial soft tissues produced by rapid and slow maxillary expansion using stereophotogrammetry: a randomized clinical trial. *Prog Orthod*. 2024;25:1. doi: 10.1186/s40510-023-00498-9
- 26- Coban G, Yavuz I, Karadas B, Demirbas AE. Three-dimensional assessment of nasal changes after maxillary advancement with impaction using stereophotogrammetry. *Korean J Orthod*. 2020;50:249-57. doi: 10.4041/kjod.2020.50.4.249
- 27- Berger JL, Pangrazio-Kulbersh V, George C, Kaczynski R. Long-term comparison of treatment outcome and stability of Class II patients treated with functional appliances versus bilateral sagittal split ramus osteotomy. *Am J Orthod Dentofacial Orthop*. 2005;127:451-64. doi: 10.1016/j.ajodo.2004.03.028
- 28- Ruf S, Pancherz H. Dentoskeletal effects and facial profile changes in young adults treated with the Herbst appliance. *Angle Orthod*. 1999;69:239-46. doi: 10.1043/0003-3219(1999)069%3C0239:deafpc%3E2.3.co;2
- 29- Gul Amuk N, Kurt G, Baysal A, Turker G. Changes in pharyngeal airway dimensions following incremental and maximum bite advancement during Herbst-rapid palatal expander appliance therapy in late adolescent and young adult patients: a randomized non-controlled prospective clinical study. *Eur J Orthod*. 2019;41:322-30. doi: 10.1093/ejo/cjz011
- 30- Choi JW, Lee JY, Yang SJ, Koh KS. The reliability of a surgery-first orthognathic approach without presurgical orthodontic treatment for skeletal class III dentofacial deformity. *Ann Plast Surg*. 2015;74:333-41. doi: 10.1097/sap.0b013e318295dcce
- 31- Kaba YN, Demirbas AE, Şirin Saribal G, Bilge S, Soylu E, Alkan A. Which factors affect the lingual fracture pattern in sagittal split ramus osteotomy? *J Craniomaxillofac Surg*. 2024;52:636-43. doi: 10.1016/j.jcms.2024.03.005.
- 32- Lim Y-K, Chu E-H, Lee D-Y, Yang I-H, Baek S-H. Three-dimensional evaluation of soft tissue change gradients after mandibular setback surgery in skeletal Class III malocclusion. *Angle Orthod*. 2010;80:896-903. doi: 10.2319/021210-90.1
- 33- Lin HH, Chiang WC, Lo LJ, Sheng-Pin Hsu S, Wang CH, et al. Artifact-resistant superimposition of digital dental models and cone-beam computed tomography images. *J Oral Maxillofac Surg*. 2013;71:1933-47. doi: 10.1016/j.joms.2013.06.199
- 34- Farkas LG. *Anthropometry of the head and face*. New York: Raven Pr Elsevier; 1995.
- 35- Farouk K, Eissa O, Ghoneima A, El-Bedwehi A, Abdel-Fattah E, Hussein F, et al. Treatment effects of Herbst appliance in skeletal Class II cases during pre-pubertal and post-pubertal periods: a cone-beam computed tomographic study. *Sci Rep*. 2023;13:11342. doi: 10.1038/s41598-023-37394-5
- 36- Lübbers HT, Medinger L, Kruse A, Grätz KW, Matthews F. Precision and accuracy of the 3dMD photogrammetric system in craniomaxillofacial application. *J Cranioc Surg*. 2010;21:763-7. doi: 10.1097/scs.0b013e3181d841f7
- 37- Day CJ, Lee RT. Three-dimensional assessment of the facial soft tissue changes that occur postoperatively in orthognathic patients. *World J Orthod*. 2006;7:15-26.
- 38- Soncul M, Bamber MA. Evaluation of facial soft tissue changes with optical surface scan after surgical correction of Class III deformities. *J Oral Maxillofac Surg*. 2004;62:1331-40. doi: 10.1016/j.joms.2004.04.019
- 39- McCance A, Moss J, Fright W, James D, Linney A. A three dimensional analysis of soft and hard tissue changes following bimaxillary orthognathic surgery in skeletal III patients. *Br J Oral Maxillofac Surg*. 1992;30(5):305-12. doi: 10.1016/0266-4356(92)90180-q
- 40- Baik H-S, Kim S-Y. Facial soft-tissue changes in skeletal Class III orthognathic surgery patients analyzed with 3-dimensional laser scanning. *Am J Orthod Dentofacial Orthop*. 2010;138:167-78. doi: 10.1016/j.ajodo.2010.02.022
- 41- Verhoeven T, Coppens C, Barkhuysen R, Bronkhorst E, Merckx M, Bergé S, et al. Three dimensional evaluation of facial asymmetry after mandibular reconstruction: validation of a new method using stereophotogrammetry. *Int J Oral Maxillofac Surg*. 2013;42:19-25. doi: 10.1016/j.ijom.2012.05.036
- 42- Purkayastha SK, Rabie AB, Wong R. Treatment of skeletal class II malocclusion in adults: stepwise vs single-step advancement with the Herbst appliance. *World J Orthod*. 2008;9:233-43.
- 43- Rabie AB, Al-Kalaly A. Does the degree of advancement during functional appliance therapy matter? *Eur J Orthod*. 2008;30:274-82. doi: 10.1093/ejo/cjm129
- 44- Hägg U, Rabie AB, Bendeus M, Wong RW, Wey MC, Du X, et al. Condylar growth and mandibular positioning with stepwise vs maximum advancement. *Am J Orthod Dentofacial Ortho*. 2008;134:525-36. doi: 10.1016/j.ijom.2012.05.036
- 45- Mariscal JF, Ravelli DB, Santos Pinto A, Dominguez GC, Arruda Aidar LA, Jacob HB. Comparisons of soft tissue changes after Herbst appliance treatment in three different growth phases. *J World Fed Orthod*. 2023;13:78-85. doi: 10.1016/j.ejwf.2023.11.002
- 46- Held AJ, Spirgi M, Cimasoni G. An orthopedically treated adult case of Class II malocclusion. *Am J Orthod*. 1963;49:761-5. Doi: 10.1016/0002-9416(63)90247-1
- 47- Manni A, Pasini M, Giuca MR, Morganti R, Cozzani M. A retrospective cephalometric study on pharyngeal airway space changes after rapid palatal expansion and Herbst appliance with or without skeletal anchorage. *Prog Orthod*. 2016;17:29. doi: 10.1186/s40510-016-0141-1
- 48- Jędrzejewski M, Smektała T, Sporniak-Tutak K, Olszewski R. Preoperative, intraoperative, and postoperative complications in orthognathic surgery: a systematic review. *Clin Oral Invest*. 2015;19:969-77. doi: 10.1007/s00784-015-1452-1.
- 49- Teerijoki-Oksa T, Jääskeläinen S, Forssell K, Forssell H, Vähätalo K, Tammisalo T, et al. Risk factors of nerve injury during mandibular sagittal split osteotomy. *Int J Oral Maxillofac Surg*. 2002;31:33-9. doi: 10.1054/ijom.2001.0157
- 50- Phillips C, Essick G, Blakey III G, Tucker M. Relationship between patients' perceptions of postsurgical sequelae and altered sensations after bilateral sagittal split osteotomy. *J Oral Maxillofac Surg*. 2007;65:597-607. doi: 10.1016/j.joms.2005.12.078
- 51- Niemi P, Kortelainen M, Harjunmaa U, Waltimo-Sirén J. Costs and duration of orthodontic-surgical treatment with mandibular advancement surgery. *Eur J Orthod*. 2023;45:558-64. doi: 10.1093/ejo/cjad051
- 52- Yang X, Zhu Y, Long H, Zhou Y, Jian F, Ye N, et al. The effectiveness of the Herbst appliance for patients with Class II malocclusion: a meta-analysis. *Eur J Orthod*. 2016;38:324-33. doi: 10.1093/ejo/cjv057
- 53- Almeida MR, Henriques JF, Almeida RR, Ursi W, McNamara Jr JA. Short-term treatment effects produced by the Herbst appliance in the mixed dentition. *Angle Orthod*. 2005;75:540-7. doi: 10.1043/0003-3219(2005)75[540:stepbt]2.0.co;2

- 54- Maal T, De Koning M, Plooi J, Verhamme LM, Rangel FA, Bergé SJ, et al. One year postoperative hard and soft tissue volumetric changes after a BSSO mandibular advancement. *Int J Oral Maxillofac Surg.* 2012;41:1137-45. doi: 10.1016/j.ijom.2012.04.004
- 55- Jakobsone G, Latkauskiene D, McNamara Jr JA. Mechanisms of Class II correction induced by the crown Herbst appliance as a single-phase Class II therapy: 1 year follow-up. *Prog Orthod.* 2013;14:27. doi: 10.1186/2196-1042-14-27
- 56- Panneerselvam E, Prasad PJ, Balasubramaniam S, Somasundaram S, Raja KV, Srinivasan D, et al. The influence of the mandibular gonial angle on the incidence of mandibular angle fracture — A radiomorphometric study. *J Oral Maxillofac Surg.* 2017;75:153-9. doi: 10.1016/j.joms.2016.08.016
- 57- Souki B, Vilefort P, Oliveira DD, Andrade I Jr, Ruellas AC, Yatabe MS, et al. Three-dimensional skeletal mandibular changes associated with Herbst appliance treatment. *Orthod Craniofac Res.* 2017;20:111-8. doi: 10.1111/ocr.12154
- 58- LeCornu M, Cevidanes LH, Zhu H, Wu CD, Larson B, Nguyen T. Three-dimensional treatment outcomes in Class II patients treated with the Herbst appliance: a pilot study. *Am J Orthod Dentofacial Orthop.* 2013;144:818-30. doi: 10.1016/j.ajodo.2013.07.014
- 59- Thygesen TH, Bardow A, Norholt SE, Jensen J, Svensson P. Surgical risk factors and maxillary nerve function after Le Fort I osteotomy. *J Oral Maxillofac Surg.* 2009;67:528-36. doi: 10.1016/j.joms.2008.07.004