

## THE HEALTH SCHOOL PROGRAMME: A HEALTH PROMOTION STRATEGY IN PRIMARY CARE IN BRAZIL

Maria de Fátima Antero Sousa Machado<sup>1</sup>, Fabiane do Amaral Gubert<sup>2</sup>,  
Anyá Pimentel Gomes Fernandes Vieira Meyer<sup>3</sup>, Yana Paula Coêlho Correia Sampaio<sup>4</sup>,  
Maria Socorro de Araújo Dias<sup>5</sup>, Ana Mattos Brito de Almeida<sup>6</sup>, Ana Patrícia Pereira Moraes<sup>7</sup>,  
Anamaria Cavalcante e Silva<sup>8</sup>, Jocileide Sales Campos<sup>8</sup>, Maristela Inês Osawa Chagas<sup>5</sup>,  
Emília Soares Chaves<sup>9</sup>

DOI: <http://dx.doi.org/10.7322/jhgd.96709>

### Abstract

**Introduction:** the Health School Programme (HSP) should be understood as a permanent development process. In this context, the actions of a policy aimed at children and adolescents are paramount in the HSP. **Objective:** to identify and describe the actions developed by the Family Health Group in the HSP, from the National Programme for Improving Access and Quality of Primary Care (PIPCAQ). **Methods:** this cross-sectional research used secondary data collected from the 17,202 groups who joined PIPCAQ in 2012. **Results:** all regions showed significant results concerning the execution of school activities. The Northern region was the one that performed most school actions (80.5%), followed by the Northeast, Midwest, South and Southeast, respectively. However, some items, such as professional training in education and health work need to be strengthened. **Conclusion:** HSP in Brazil has mobilised significant actions, even though it has not happened in homogeneously in all Brazilian regions.

**Keywords:** health evaluation, health school programme, primary health care.

### INTRODUCTION

In recent years in Brazil, numerous initiatives and evaluation experiences in primary care have been implemented to achieve improvements in health policies<sup>1,2</sup>.

The National Programme for Improving Access and Quality of Primary Care (PIPCAQ) has, as its main objective, stimulating the expansion of access to, and improvements in, primary health care, in order to allow greater transparency and effectiveness of government actions aimed at primary healthcare.

The PIPCAQ evaluated different health promotion actions carried out in primary care by the Health School Programme (HSP). For the Health Ministry,<sup>3</sup> the HSP is intended to provide a range of actions for prevention, promotion and attention in the health of children, adolescents and young students, from the basic level of public education, through to the communication between public

schools and teams from the Family Health Strategy (FHS), and the realisation of actions directed at students<sup>3</sup>.

The HSP is now major public policy for childhood and adolescence. Among its components are clinical, nutritional and healthy eating promotion, ophthalmologic evaluation, and continuing education actions in health, physical activity, the promotion of a prevention culture in the school context and the inclusion of health education topics in the school political pedagogical project<sup>4</sup>.

Health promotion in the school environment should be understood as a process in permanent development. In order to review and improve actions in the public management of health and professional work processes, and obtain more efficient preventive interventions, these ones have to be realised in places that present the national strategies and programmes in a public health context.

1 Universidade Regional do Cariri (URCA) – Crato (CE), Brasil.

2 Universidade Federal do Ceará (UFC) - Fortaleza (CE), Brasil.

3 Fundação Oswaldo Cruz, (FIOCRUZ) Fortaleza (CE), Brasil.

4 Faculdade de Medicina Estácio de Juazeiro do Norte, (Estácio FMJ) – Juazeiro do Norte (CE), Brasil.

5 Universidade Estadual Vale do Acaraú (UVA) - Sobral (CE), Brasil.

6 Escola de Saúde Pública do Ceará (ESPCE) – Fortaleza (CE), Brasil.

7 Universidade Estadual do Ceará (UECE) – Fortaleza (CE), Brasil.

8 Centro Universitário Christus (UNICHRISTUS), Fortaleza (CE), Brasil.

9 Universidade da Integração Internacional da Lusofonia Afro-Brasileira (UNILAB) – Redenção (CE), Brasil.

Regional project coordinators (EC, PI and RN) of the Department of Public Health, Federal University of Rio Grande do Norte/UFRN, Profs. Severina Alice Uchoa, Paulo Medeiro Rocha, Themis Xavier Pinheiro.

**Corresponding author:** Maria de Fátima Antero Sousa Machado. E-mail: [fatimaantero@uol.com.br](mailto:fatimaantero@uol.com.br)

**Suggested citation:** Machado MFAS, Gubert FA, Meyer APGFV, Sampaio YPCC, Dias MSA, Almeida AMB et al. The health school programme: a health promotion strategy in primary care in Brazil. *Journal of Human Growth and Development*. 25(3): 307-312. Doi: <http://dx.doi.org/10.7322/jhgd.96709>

Manuscript submitted Oct 22 2014, accepted for publication Dec 19 2014.

The purpose of the study is therefore to identify and describe the actions developed in Brazil by the Family Health Teams (FHT) in the HSP.

## METHODS

This is a cross-sectional study with a quantitative approach, carried out through secondary data provided by the Health Ministry, collected from 17,202 Family Health Teams (FHT) that joined the PIPCAQ in 2012 in Brazil.

The PIPCAQ aims to increase access to, and improve the quality of, primary health care, and is organised in four complementary and robust phases: (1) adherence and contracting; (2) development; (3) external evaluation, and (4) re-contracting. Data for this study were extracted from the third phase of PIPCAQ and it was conducted by teaching and/or research institutions contracted by the Health Ministry in the period June to October 2012.

The third phase, external evaluation, has been carried out by means of an instrument application that sought to determine the conditions of access to health services and the care quality of participating municipalities and primary care groups. The assessment tool comprised four information gathering modules. Module I asked about the infrastructure, materials, supplies, and medicine conditions of the basic health unit. Module II sought information about the group work process and about the organisation's care of the user and the actions taken by HSP. Module III checked user satisfaction with, and perception of, the health service, and Module IV represented a set of complementary information to the other modules, answered online by local management.

The results of this study are from Module II, specifically issues relating to health actions carried out within a school. The instrument contained questions relating to the activities of HSP conducted by groups of the FHS and was answered only with 'yes' or 'no', as follows: (Quadro 1).

### Quadro 1: Questions used in PIPCAQ for the evaluation of the HSP. Fortaleza, Ceará, Brazil, 2012.

Does the group conduct activities in school?

What types of services are provided by the group?

Does the group record the activities?

Does the group enter information in the Planning, Budget and Finance Integrated System of the Education Ministry: ISEM/HSP?

Has it identified the number of students with the health needs survey?

Does the team plan the activities?

What clinical evaluation activities are carried out (update immunisation schedule; early detection of hypertension; detection of neglected health disorders; anthropometric measurements; ophthalmologic evaluation; auditory evaluation; psychosocial assessment, nutritional assessment and evaluation of oral health)?

Which prevention and health promotion activities are carried out (educational activities on the promotion of nutrition and healthy lifestyles, promotion of corporal practices and physical activity in schools, education for sexual health, reproductive health and STD/AIDS; actions prevention of alcohol, tobacco and other drugs)?

Are training activities for education professionals to work with health education held, as well as discussions with school teachers?

The questionnaire was applied on location, conducted by an interviewer and only one professional group was selected to answer the questions, in a suitable place in the health unit and on a date agreed with the municipal management. The PIPCAQ participation of municipalities and teams was voluntary.

The data were automatically transferred to the national database of the Health Ministry. The study variables were the activities carried out by health professionals in school and are shown in Table 1. The analyses were performed by comparing the regions. The Statistical Package for Social Sciences software was used (SPSS Inc., Chicago, USA, version 19.0). Descriptive analyses were performed and categorical variables were analysed using non-parametric tests, Chi-square.

## RESULTS

A significant association between the regions was observed, according the value of  $p < 0.0001$  to all items of this study.

With regard to the activities that are performed in the schools, all regions showed impressive results, ranging from 80.5% to 69.4%. The Northern region performed more activities in this scenario, followed by the Northeast, Midwest, South and Southeast, respectively.

Children and adolescent care was observed in the study and the Northeast region achieved 47.7%. This same region showed better indicators when it came to entering information in the Planning, Budget and Finance Integrated System of the Ministry of Education (ISEM/HSP) with 29.5%

of health groups doing so. The Southeast region only had 12.6% of the groups entering information into the system.

With regard to the survey of the number of students and the need for health, 34.9% of health teams in the Northeast region performed it. The school actions planning item was considered more important than the school students survey, with 68.5% of the Northeast region. The Southeast region presented values of 28.2% and 56.8% to both items, respectively, considered the lowest among the regions of Brazil. Considering the importance of these two aspects (survey needs and planning of actions), the association was made between these variables, and the result was obtained that the regions who planned their activities also made a survey of health needs with a significance of  $p < 0.001$ .

On the issue of student clinical evaluation, the Midwest region presented at 10.4%, the region where most reported having performed this procedure. The update of the immunisation schedule is highlighted in the Northeast region (55.6%), followed by the North (52.6%). Northeastern Brazil

also achieved better results in the early detection of hypertension (40.4%), neglected diseases (30.2%), anthropometric assessment (57.9%), hearing evaluation (16.0%), psychosocial assessment (25.4%), nutritional assessment (52.6%), oral health assessment (59.6%), and food security actions and healthy eating (61.1%). The South region performed best in ophthalmic evaluation, with 24.6%.

For promotion and health education activities, the Northern region can be observed as the one that promoted activities in the theme of sexual and reproductive health and sexually transmitted diseases, with 65.1% and preventing the use of alcohol, tobacco and other drugs at 47.7%.

In the item related to training activities for education professionals to work with health education and have discussions with school teachers, the Northeast region was highlighted, at 23.2% and 33.2%, respectively. The Southeast region had the lowest percentages in these two items with 13.3% and 20.1%, respectively. The results are shown in Table 2 below: (Table 1).

**Table 1:** Distribution of activities executed by the Family Health teams that participate in the PIPCAQ, Brazil, 2013.

HSP Activities	North Region (n = 1045)				Northeast Region (n = 5559)				Southeast Region (n = 6569)				South Region (n = 2920)				Central-West Region (n = 1131)				p *
	S	%	N	%	S	%	N	%	S	%	N	%	S	%	N	%	S	%	N	%	
Executes school activities	841	<b>80,5</b>	196	18,8	4431	79,7	1124	20,2	4559	69,4	2010	30,6	2240	76,7	679	23,3	869	76,8	236	20,1	<b>0,001</b>
Attend ance routine	452	43,3	389	37,2	2652	<b>47,7</b>	1779	32,0	2475	37,7	2084	31,07	1273	43,6	967	33,1	395	34,9	474	43,0	<b>0,001</b>
ISEM/HSP informations	217	20,8	624	59,7	1639	<b>29,5</b>	2792	50,2	826	12,6	3733	56,8	428	14,7	1812	62,1	292	25,8	577	51,3	<b>0,001</b>
Scholars with health needs	281	26,9	560	53,3	1938	<b>34,9</b>	2493	44,8	1855	28,2	2704	41,2	980	33,6	1260	43,2	340	30,0	529	47,4	<b>0,001</b>
Planes activities	686	65,6	155	14,8	3806	<b>68,5</b>	625	11,2	3734	56,8	825	12,6	1866	63,9	374	12,8	670	59,2	199	17,7	<b>0,001</b>
Executes clinic evaluation	109	<b>10,4</b>	732	70	435	7,8	3996	71,9	614	9,4	3945	60	299	10,2	1941	66,5	91	8,2	778	85,0	<b>0,001</b>
Vaccine calendar update	550	52,6	291	27,8	3089	<b>55,6</b>	1342	24,1	2409	36,7	2150	32,7	1267	43,4	973	33,3	545	49,1	324	48,0	<b>0,001</b>
Preccocious systemic arterial hypertension detection	386	36,9	455	43,5	2248	<b>40,4</b>	2183	39,3	1583	24,1	2976	45,3	704	24,1	1536	52,6	348	31,4	521	58,0	<b>0,001</b>
Neglected grievances detection	284	27,2	557	53,3	1677	<b>30,2</b>	2754	49,5	1451	22,1	3108	47,3	825	28,3	1415	48,5	264	23,8	605	68,0	<b>0,001</b>
Anthropometric evaluation	525	50,2	316	30,2	3219	<b>57,9</b>	1212	21,8	2435	37,1	2124	32,3	1322	45,8	918	31,4	584	52,7	285	36,8	<b>0,001</b>
Ophthalmic evaluation	227	21,7	614	58,8	1155	<b>20,8</b>	3276	58,9	1197	18,2	3362	51,2	718	<b>24,6</b>	1522	52,1	195	17,3	674	86,7	<b>0,001</b>
Auditory evaluation	120	11,5	721	6,9	887	<b>16,0</b>	3544	63,8	709	10,8	3850	58,6	382	13,1	1858	63,7	159	14,3	710	64,4	<b>0,001</b>
Psychosocial evaluation	200	19,1	641	61,3	1410	<b>25,4</b>	3021	54,3	1159	17,6	3400	51,8	661	22,6	1579	54,1	247	22,3	622	56,9	<b>0,001</b>
Nutricional evaluation	446	42,7	395	37,8	2922	<b>52,6</b>	1509	27,1	2120	32,3	2439	37,1	1157	39,6	1083	37,1	499	45,0	370	33,9	<b>0,001</b>
Bucal health evaluation	551	52,7	290	27,8	3325	<b>59,6</b>	1106	19,9	2854	43,4	1705	26,0	1623	55,6	617	21,1	584	52,7	285	25,1	<b>0,001</b>
Feed security actions and healthy eating	588	56,3	253	24,2	3398	<b>61,1</b>	1033	18,6	2774	42,2	1785	27,2	1421	48,7	819	28,1	592	53,4	277	25,5	<b>0,001</b>
Corporal practices promotion and physical activity	362	34,6	479	45,8	1887	33,9	2544	<b>45,8</b>	1599	24,3	2960	45,1	750	25,7	1490	51	377	34,0	492	45,0	<b>0,001</b>
Sexual and reproductive health and STD/AIDS prevention	680	<b>65,1</b>	161	15,4	3256	58,6	1175	21,1	3038	46,2	1521	23,2	1679	57,5	561	19,2	662	59,7	207	18,5	<b>0,001</b>
Alcohol, tobacco and other drugs use prevention	491	<b>47,0</b>	350	33,5	2399	43,7	2032	36,6	2162	32,9	2397	36,5	1290	44,2	950	32,5	517	46,6	352	31,9	<b>0,001</b>
Professionals qualification to work education/health	215	20,6	626	59,9	1291	<b>23,2</b>	3140	56,5	1137	17,3	3422	52,1	646	22,1	1594	54,5	244	22,0	625	57,4	<b>0,001</b>
Debate with teachers	260	24,9	581	55,6	1848	<b>33,2</b>	2583	46,5	1322	20,1	3237	49,3	814	27,9	1426	48,6	224	20,3	645	60,0	<b>0,001</b>

**DISCUSSION**

The HS Pat the national level has mobilised significant actions, even if not evenly across all regions, as shown in the PIPCAQ evaluation results in 2012. Thus, this study contributes an overview of the actions developed in the HSP across the country and can certainly produce a reflection on the range of this programme and its contributions to the improvement in the care of the population. The importance of evaluation studies that allow an overview of reality by the studying the specific case of research is highlighted, as in the case of the HSP in Brazil.

With regard to the promotion and health education actions, the Northern region was the one that most promoted these activities<sup>5</sup>. It is worth mentioning that the Northern and Northeast regions are the ones with more social and economic inequalities than in the rest of the country, with the longest inclusion of Family Strategy activities, and with closer, more effective links.

Difficulties in implementing routine care delivered by health professionals still exist, given that the other community demands of the FHG end up needing a different look. It is necessary here to rethink the HSP from a broader perspective, discussing public policies in each territory, providing

appropriate environments, and reorienting health services beyond clinical and curative treatments. The health teams seem to hold few clinical actions with schools, since they have a high rate of activity, but still with a fragile routine<sup>6</sup>.

Studies<sup>7,8</sup>, describe the difficulty of establishing a routine in helping adolescents and that this may be linked to the need for some professionals to overcome prejudice before delivering educational programmes in school. Little exposure to the issues of this age group can mean a tendency to view adolescents in a negative and stereotypical manner, embarrassment in dealing with sexuality and violence situations (which may require interdisciplinary interventions), and work overload, making access to learning difficult. These and other situations are factors which can influence this point. For many teenagers, health is still linked to 'absence of disease' and their sense of invulnerability and immortality can justify the indifference and detachment that is observed in the behaviour of adolescents in relation to health services<sup>9</sup>.

The low data entry percentage in ISEM/HSP has hampered the monitoring and evaluation of these actions. Consistent monitoring allows managers and municipalities to demarcate their responsibilities and define the areas of coverage of the FHS teams, contributing to the situational diagnosis based on social determinants in the epidemiological scenario, including the transfer of appropriate incentives for the units.

Studies indicate there is a lack of regularity in the system, a lack of professional training to fill in the spreadsheets and, according to the professionals themselves, the vast number of information systems deployed by the Health Ministry in recent years has demanded too much effort from the municipalities. To overcome this situation, caregivers need support in the planning process and decision-making powers with regard to information systems<sup>9</sup>.

The actions planning at school was also evaluated and considered one of the most positive aspects. This shows that health groups are participating in the guidelines proposed by the Health Ministry which states that the execution time of each HSP action is planned by the FHG, taking into account the school year and the school political pedagogical project<sup>4</sup>.

Research conducted in Fortaleza showed the gradual strengthening of the integration of the education and health sectors, promoting the inter-sectoral approach proposed by the Unified Health System, despite the challenges such as difficulty maintaining these actions<sup>10</sup>.

The school environment has been recognised as a beneficial setting for promoting health practices, prevention and health education. In this context, it reinforces the prevention of health problems, contributing to the creation of conditions for student learning, and the construction of a social care system, focusing on the promotion of citizenship and human rights. It provides a strengthening of the capacity to cope with vulnerabilities in the health field, which can

compromise full school development<sup>11</sup>. Thus, it is within this approach, a health programme at school, inserted and integrated into daily life and school culture, is understood and justified. It is a very important reference space for children and teenagers, who increasingly develop their experiences within significant socialisation spaces and community life<sup>11</sup>.

It was found that the clinical evaluation of schools was used for some of the health units in Brazil. To improve care of children and adolescents, realising the clinical evaluation is necessary to provide adequate physical spaces<sup>12</sup> and transport for moving teams, as this becomes an essential resource for maintaining frequent contact between users and staff<sup>13</sup>, and providing innovative educational technologies that streamline and facilitate teaching and learning<sup>14</sup>.

The work with the schools regarding health education showed actions directed at promoting sexual and reproductive health were the most frequently implemented across the country, with high percentages in all regions. According to the Health Ministry<sup>4</sup> there has been an increase in the pregnancy rate in girls in recent decades. Also, teenagers have become more exposed to sexually transmitted diseases, such as AIDS, with the inappropriate use of contraceptive methods, coupled with lack of knowledge of, and access to, them. Another important topic discussed at school was the use of alcohol and drugs, and the Northern region was highlighted again. In this sense, the school's partnership with primary care is essential for dialogue, access to information and the re-organisation of practice is the main focus.

With regard to training for operations in the HSP, the study showed that the actions of this nature are not well performed. Ongoing training for professionals who work in schools is essential, and the need to hire professionals from diverse backgrounds, with skills and experience in health promotion actions<sup>15</sup>.

The implications of these findings, to identify the actions taken by HSP, reaffirms the commitment that public policy has to Brazilian children and adolescents, directly impacting their quality of life and their health. The transfer of the knowledge between the education and health areas enhances the development of actions that favour the educational dimension of healthcare, the care of oneself, of others and the environment, generating effects on healthy development, the role of education and the community in which they live, allowing them to have options that improve their quality of life<sup>16</sup>.

Thus, to know and handle risk factors and the vulnerabilities of children and adolescents in promoting and protecting health will impact positively on quality of life, on learning conditions and, consequently, on the construction of citizenship<sup>11</sup>.

Issues to be addressed in the school setting, such as sexuality, reveal the importance of developing an environment for health promotion through health education in this context, since it acts as a significant social space where



teenagers build knowledge and promote health in decision-making from their experiences and life expectancies, and trivia about sexuality, among other aspects relating to health<sup>17</sup>.

However, it is important to emphasise that it is not enough to have open space to promote health. Developing responsibility in the users in these practices, and student participation in the mobilisation, training and development of individual learning and social skills to deal with the processes of health and disease, is essential for implementation of these actions<sup>18</sup>.

Thus, it is necessary that these are consistent with the assumptions of health promotion; however, those actions are still a challenge, given the predominance of curative characteristics and individualistic practices<sup>19</sup>.

For the realisation of educational activities, the involvement of all stakeholders is a unique prerequisite for the full exercise of health<sup>20</sup>, however in the context involving the HSP, one of the challenges for implementation is the training of education professionals to work with education in health.

Research should be implemented to give visibility to situations such as these. However, while

in Brazil scientific production has been increasing significantly, this could increase even more with the proper training of health professionals who work with research in project management<sup>21</sup>.

Investments in permanent health education that contribute to the transformation of professional, educational and health practices, and also to the organisation of services, are essential strategies for improving these HSP actions.<sup>11</sup>

In this way, it has been shown that the HSP in Brazil has mobilised significant actions, even if it has not performed homogeneously in all Brazilian regions; although all the regions have had elements evaluated by PIPCAQ by presenting the implementation of actions, the regions that have performed more actions are the North and Northeast.

### Acknowledgement

Regional project coordinators (EC, PI and RN) of the Department of Public Health, Federal University of Rio Grande do Norte / UFRN, Profs. Severina Alice Uchoa, Paulo Medeiros Themis Rocha and Xavier Pinheiro.

### REFERENCES

- Miclos PV, Calvo MCM, Colussi CF. Avaliação do desempenho da atenção primária em saúde através da análise envoltória de dados. *Rev Eletr Gestão Saúde*. (2015; 6(2): 1749-63.
- Pinto Junior EP, Cavalcante JLM, Sousa RA, Moraes APP, Silva MGC. Análise da produção científica sobre avaliação, no contexto da saúde da família, em periódicos brasileiros. *Saúde Debate*. 2015; 39(104): 268-78. DOI: <http://dx.doi.org/10.1590/0103-110420151040226>
- Ministério da Saúde. Secretaria de Atenção à Saúde. Departamento de Atenção Básica. Instituto PSE. Brasília: Ministério da Saúde; 2011.
- Ministério da Saúde. Secretaria de Atenção à Saúde. Departamento de atenção básica. Cadernos de Atenção Básica. Saúde na escola. Brasília: Ministério da Saúde; 2009.
- Fausto MCR, Fonseca HS. Rotas de atenção básica no Brasil: experiências do trabalho de campo PMAQ/AB. Rio de Janeiro: Saberes; 2013.
- Gomes LC. O desafio da intersetorialidade: a experiência do Programa Saúde na Escola (PSE) em Manguinhos, no Município do Rio de Janeiro. Dissertação (Mestrado) - Escola Nacional de Saúde Pública Sergio Arouca. Rio de Janeiro: 2012.
- Vitalle MSS, Almeida RG, Silva FC. Capacitação na atenção à saúde do adolescente: experiência de ensino. *Rev Bras Educ Med*. 2010; 34(3): 459-468.
- Marques JF, Queiroz MV. Cuidado ao adolescente na atenção básica: necessidades dos usuários e sua relação com o serviço. *Rev Gaúcha Enferm*. 2012; 33(3): 65-72. DOI: <http://dx.doi.org/10.1590/S1983-14472012000300009>
- Raposo C. A Política de Atenção Integral à Saúde do Adolescente e Jovem: uma perspectiva de garantia de direito à saúde? *Rev Pauta*. Rio de Janeiro: 2009; 6(23): 117-38.
- Santiago LM, Rodrigues MTP, Oliveira Junior AD, Moreira TMM. Implantação do Programa Saúde na Escola em Fortaleza-CE: atuação de equipe da Estratégia Saúde da Família. *Rev Bras Enferm*. 2012; 65(6): 1026-9. DOI: <http://dx.doi.org/10.1590/S0034-71672012000600020>
- Ferraz DAS, Nemes MIB. Avaliação da implantação de atividades de prevenção das DST/AIDS na atenção básica: um estudo de caso na Região Metropolitana de São Paulo, Brasil. *Cad Saúde Pública*. 2009; 25(Suppl2): S240-50. DOI: <http://dx.doi.org/10.1590/S0102-311X2009001400006>
- Baratieri T, Marcon SS. Longitudinalidade no trabalho do enfermeiro: identificando dificuldades e perspectivas de transformação. *Texto Contexto Enferm*. 2012; 21(3): 549-57. DOI: <http://dx.doi.org/10.1590/S0104-07072012000300009>
- Roecker S, Budó MLD, Marcon SS. Trabalho educativo do enfermeiro na Estratégia Saúde da Família: dificuldades e perspectivas de mudanças. *Rev Esc Enferm USP*. 2012; 46(3): 641-9. DOI: <http://dx.doi.org/10.1590/S0080-62342012000300016>
- Silva Junior AJ. Programa saúde na escola: limites e possibilidades intersetoriais. *Interface*. 2014; 18(51): 799. DOI: <http://dx.doi.org/10.1590/1807-57622014.0583>
- Ferreira IRC, Moysés SJ, França BHS, Carvalho ML, Moysés ST. Percepções de gestores locais sobre a intersetorialidade no Programa Saúde na Escola. *Rev Bras Educ*. 2014; 19(56): 61-76.

DOI: <http://dx.doi.org/10.1590/S1413-24782014000100004>

16. Bezerra IMP, Machado MFAS, Souza OF, Antão JYFL, Dantas MNL, Reis AOA, et al. Professional activity in the context of health education: a systematic review. *J Hum Growth Dev.* 2014; 24(3): 255-262. DOI: <http://dx.doi.org/10.7322/jhdg.88909>
17. Arash HK, Carpentier R. The evolving role of public health in the delivery of health care. *J Hum Growth Dev.* 2012; 22(3): 396-399. systematic review. *J Hum Growth Dev.* 2014;24(3): 255-262.

## Resumo

**Introdução:** a promoção da saúde no cenário escolar deve ser entendida como um processo em permanente desenvolvimento. Nesse contexto, destacam-se as ações do Programa Saúde na Escola, como política voltada para crianças e adolescentes. **Objetivo:** identificar e descrever as ações desenvolvidas pelas Equipes de Saúde da Família (ESF) no Programa Saúde na Escola (PSE), participantes do Programa Nacional de Melhoria do Acesso e da Qualidade da Atenção Básica (PMAQ). **Método:** trata-se de uma pesquisa transversal que utilizou dados secundários coletados junto a 17.202 equipes que aderiram ao PMAQ em 2012. **Resultado:** todas as regiões apresentaram resultados expressivos acerca da realização de atividades na escola. A região Norte é a que mais executa ações ao escolar (80,5%), seguidas das regiões Nordeste, Centro-Oeste, Sul e Sudeste respectivamente. Contudo, alguns itens como a capacitação dos profissionais para trabalhar com educação e saúde precisam ser fortalecidos. **Conclusão:** o PSE no Brasil tem mobilizado ações relevantes, mesmo que isto não tenha se dado de forma homogênea em todas as regiões brasileiras.

**Palavras-chave:** avaliação em saúde, saúde escolar, atenção primária à saúde.