

Global panorama of studies about freshwater oligochaetes: main trends and gaps

Luciana Falci Theza Rodrigues^{1,2} & Roberto da Gama Alves^{1,3}

¹ Universidade Federal de Juiz de Fora (UFJF), Instituto de Ciências Biológicas (ICB), Departamento de Zoologia (DEPZOO), Laboratório de Invertebrados Bentônicos (LIB). Juiz de Fora, MG, Brasil.

² ORCID: 0000-0002-7610-870X. E-mail: biofalci@hotmail.com (corresponding author)

³ ORCID: 0000-0001-8392-2091. E-mail: gama.alves@ufjf.edu.br

Abstract. Freshwater oligochaetes have been studied over the years in a wide range of habitats around the world. To analyse the data published about freshwater oligochaetes in the 31 year period between 1985 and 2015, we searched for documents in the ISI Web of Science, Scopus and SciELO databases. A total of 979 works were evaluated from 184 periodicals. The United States is the country with the most publications about aquatic oligochaetes, followed by France, Germany and Italy. Works related to the ecology and ecotoxicology of these organisms are most abundant. Studies carried out in the laboratory are most frequent, consisting of investigations involving bioassays, morphology, genetics and molecular biology. The results also show that spring, cave water bodies and swamps have been studied less than any other aquatic habitat and that countries of the Southern Hemisphere are under-represented in the studies of aquatic oligochaetes. Hence, there is a need for increased efforts to fill in gaps in the knowledge about these oligochaetes.

Key-Words. Database; Literature review; Scientific production; Oligochaeta.

INTRODUCTION

Oligochaetes in freshwater ecosystems are composed of over 1,100 known species (Martin *et al.*, 2008). Despite their high richness and representation in various aquatic habitats (streams, rivers, lakes, reservoirs), the oligochaetes have not received proper attention in hydrobiology studies (Sambugar, 2007). Their identification in the majority of publications is at a relatively unrefined taxonomic level, such as class or family, causing a gap in knowledge about the species level. Their small body size, absence of peculiar morphological structures (Sambugar, 2007) and “poor” appearance, described by Righi (1984) as “lack of beauty and eye-catching behaviors”, are factors contributing to the relative lack of interest of researchers.

Studies of freshwater oligochaetes in the Neotropical realm started in the nineteenth century, carried out by vanguard researchers like Michaelsen, Stephenson, Beddard, Cognetti de Martiis and Cernosvitov, among others (Gavrilov, 1981). In Brazil, knowledge about limnic oligochaetes was pioneered by the taxonomic studies of the husband-wife team Eveline Du Bois-Reymond Marcus and Ernest Marcus starting in 1934, and by Gilberto Righi at the end of the 1960s. Subsequently (as of 2000), works about ecology such as Alves & Lucca (2000), Alves *et al.* (2006, 2008), Gorni & Alves (2006, 2008a,b, 2012, 2014), Behrend *et al.* (2009, 2012), Martins & Alves

(2008, 2010), Ragonha & Takeda (2014), Ragonha *et al.* (2014), Gorni *et al.* (2015), Oda *et al.* (2015), Petsch *et al.* (2015) and Rodrigues *et al.* (2013, 2015, 2016), and on aspects of the reproductive biology by Lobo & Alves (2011a,b), have contributed to the knowledge of these invertebrates in Brazil.

In other biogeographic regions, such as the Nearctic (Wetzel *et al.*, 2006, 2007) and the Australasian (Pinder *et al.*, 2006), information about worms is plentiful (Martin *et al.*, 2008). Although more studies about aquatic oligochaetes have been performed in these regions, not all habitats have been equally investigated. For example, little is known about the diversity of oligochaetes in groundwater (Giani *et al.*, 2001) and springs (Dumnicka, 2006).

In this context, the objective of this work was to analyze the publications on freshwater oligochaetes over time and in different geopolitical regions of the world, and based on the results to verify trends and provide information to help guide efforts to fill in the main knowledge gaps identified.

MATERIAL AND METHODS

To analyze the theme “freshwater oligochaetes”, we used the bibliographic production from 1985 to 2015. The search for data was performed using the following databases: ISI Web of Science (Thomson Reuters), Scopus (Elsevier) and SciELO (Scientific

Electronic Library Online). We used these three bases due to the different scopes regarding the number and quality of the scientific periodicals indexed. The keywords were: "aquatic oligochaet*" or "*water oligochaet*" or *clitellat** or *tubificid** or *naidid** or *lumbriculid** or *lumbriculus** or *tubifex**. Because the group is paraphyletic, many researchers avoid using the name "oligochaete" or "Oligochaeta" and instead use the term "clitellate" or "Clitellata". For this reason, we included this as a keyword. Furthermore, in many documents only the family names are given, not "oligochaete", so we also included the main families of freshwater oligochaetes, as well as some of their genera.

We selected the "Title" option for advance search, to filter only specific works about the theme of interest. We only considered articles, reviews, notes, books and book chapters. Proceedings papers, meeting abstracts and editorial pieces were excluded from the analysis. Often these types of writings wind up being redundant, such as when the results presented in proceeding abstracts are later included in scientific articles. All the documents found were analyzed individually and those found not to be related to the theme were removed. The databases were searched in August 2016.

The elements tallied in each article were: (i) year of publication; (ii) number of publications per country/continent (where the study was conducted); (iii) area of knowledge (genetics, taxonomy, ecology, morphology, among others); and (iv) type of habitat where the study was performed (river, stream, lake, reservoir, spring, laboratory, among others).

RESULTS

We found 2,325 results for publications containing the keywords in their titles. Of this number, 1,240 (53.33%) specifically investigated freshwater oligochaetes (the rest were related to terrestrial, marine, estuarine forms or clitellate other than oligochaetes, such as Hirudinea and Branchiobdellida). Of all the publications found referring to freshwater oligochaetes, 33 came from consulting SciELO, 621 from Scopus and 586 from Web of Science (Fig. 1). However, some of these works were found in more than one of the databases, so the final total was 979 publications analyzed (see Supporting Information).

The works analyzed were mainly scientific articles (97.34%). The others were reviews (1.63%), notes (0.82%) and book chapters (0.20%). These works were published in 184 different periodicals, but most of them (about 73%) contained fewer than five works. The most frequent periodicals were *Hydrobiologia* (24.56%), *Environmental Toxicology and Chemistry* (7.63%) and *Annales de Limnologie-International Journal of Limnology* (3.66%).

The number of publications on freshwater oligochaetes fluctuated during the years studied, without showing a gradual growth trend. The two years with the largest numbers (Fig. 2A) were 1987 (55 works) and 1999 (54 works). On the other hand, there was a rising trend in the sum of the articles in the 10-year periods (Fig. 2B).

Studies related to ecology, ecotoxicology and taxonomy were most abundant. "Others" in the topic category included works related to oligochaetes as food for fish and use in activated sludge reduction (Fig. 3A). The interest in ecotoxicology increased in the three decades, as did studies involving taxonomy and genetics/phylogeny (Fig. 3B). Among the oligochaetes species most cited and analyzed are *Tubifex tubifex* (Müller, 1774), *Lumbriculus variegatus* (Müller, 1774), *Branchiura sowerbyi* Beddard, 1892, *Limnodrilus hoffmeisteri* Claparède, 1862 and *Chaetogaster limnaei* Von Baer, 1827, (not necessarily in




Figure 1. Total number of exclusive and shared publications on oligochaetes in freshwater habitats found in each database entre 1985 e 2015.




Figure 2. Number of works on freshwater oligochaetes published between 1985 and 2015 (A) and 10-year totals (B).



**A****B**

Figure 3. Percentage of works on freshwater oligochaetes published between 1985 and 2015 by area of knowledge (A) and 10-year totals (B). The “Others” category contains works related to oligochaetes as food for fish and used in reduction of activated sludge.

this order), mainly in investigations related to toxicity, organic pollution and parasitism.

We found works on oligochaetes in various aquatic environments, such as groundwater, swamps, caves, springs, rivers, streams, lakes and reservoirs. In many cases the studies analyzed the interaction of these worms with other organisms (bryophytes, macrophytes, bromeliads, sponges, snails, insects, anurans and fish). We also found experimental laboratory studies (the latter being most frequent, accounting for about 50% of the total publications). Besides empirical studies in these various aquatic environments, we also found studies only using bibliographic data as primary source. Springs, cave water bodies and swamps were the least studied habitat in the past 31 years, representing under 2.5% of the published works (Fig. 4).

With respect to global distribution, studies of freshwater oligochaetes occurred in 51 countries, the five leading ones being the United States ($n = 157$), France ($n = 72$), Germany ($n = 60$), Italy ($n = 59$) and Canada ($n = 52$). Brazil and Argentina are the only Latin American countries with more than 30 published studies and appearing among the 15 leading countries (Fig. 5A). In the continental breakdown, Europe showed the highest percentage, with about 50% of the total, while Oceania and Africa each accounted for under 1%. Approximately 2.5% of the papers analyzed did not mention the place where the study was conducted, so they are included in the “Undefined” category (Fig. 5B).




Figure 4. Percentage of works on freshwater oligochaetes published between 1985 and 2015 by habitat/ecosystem where the study was conducted.



**A****B**

Figure 5. Percentage of works on freshwater oligochaetes published between 1985 and 2015 by continent (A) and by countries (15 leading ones in number of publications) (B).

DISCUSSION

Freshwater oligochaetes have been the target of studies involving biodiversity of aquatic habitats in preserved areas, biomonitoring, water quality analysis and toxicity monitoring in many countries. Therefore, researchers have published their findings in periodicals with different scopes, both widely distributed and limited, such as regional or local periodicals. This is shown by the survey of the three databases chosen here.

SciELO is a narrower database, because it only includes periodicals from countries in Latin America, along with South Africa, Spain and Portugal. For this reason, substantially fewer publications were found in this database. In contrast, Web of Science (WoS) and Scopus have global scope and presented a larger number of publications. The WoS database indexes more than 12,700 periodicals, in a wide range of scientific fields, and the majority of its titles are in English (Lopes *et al.*, 2012). Only about 2.5%

of the titles are from Latin American countries, explaining the small overlap of articles with SciELO. On the other hand, the Scopus base contains about 18,000 titles and has more European content than WoS, as well as including more languages besides English (about 60% of the coverage is countries other than the U.S.) (Lopes *et al.*, 2012). Many periodicals not indexed in WoS are indexed in Scopus, explaining the larger coverages of articles in the latter database and the greater overlap with SciELO.

According to Brinkhurst (1999), the majority of publications on oligochaetes up to 1970 were taxonomic, containing descriptions of new species and re-descriptions of known ones. In the 31 years examined (1985-2015), studies with ecological focus stood out over the other areas of knowledge, although the number of taxonomic works also continued to rise in those years. Besides this, there was a strong trend for increase of works related to toxicology (Rodriguez & Reynoldson, 2011). The increase in the number of publications in general with time and regarding determined academic fields, even if timid in some cases, is an indication of the increase in the number of researchers and/or interest in investigating certain themes. The increase in the total number of publications also indicates the scientific advances made, considering that the number of publications is one of the metrics most often used to quantify the evolution of science (Lima-Ribeiro *et al.*, 2007).

Bioassays and toxicity tests were largely responsible for the large number of laboratory studies found, and for the same reason, species such as *Tubifex tubifex*, *Branchiura sowerbyi* and *Lumbriculus variegatus* were frequently found in the titles. These species are typical models in ecotoxicological studies due to their high reproductive rate and ease cultivation (Lobo & Alves, 2011a). Works involving biology, morphology, genetics and molecular biology also accounted for a large number of publications about laboratory experiments.

Studies of oligochaetes in springs were less frequent, as were those focused on swamps and water bodies in caves. With respect to springs, for example Cantonati *et al.* (2012) also affirm that in general this type of habitat is less frequently studied than other aquatic ecosystems. Therefore, these three ecosystems deserve greater attention in future studies. This rarity of studies of oligochaetes in some aquatic environments makes it urgent to learn more about the fauna of these habitats, which are facing increasing degradation from human activity.

Nearly half of the works analyzed were European. The European continent has 61 countries, and of these, studies were found about freshwater oligochaetes in 30 of them, with a large number of overall publications. The American continent as a whole (North, Central and South America) was in second place, although the largest number of works were from the United States. Countries of Africa and Oceania were under-represented in studies of freshwater oligochaetes. This result indicates a need for efforts to fill in gaps in various areas of knowledge of these organisms, as well as the preservation of aquatic ecosystems in these regions. In particular, these regions contain many forests and aquatic ecosystems that under-

threat of human activity (Nicacio & Juen, 2015), accounting for nearly half (44%) of the global hotspots.

Among Latin American countries, considered emerging or developing, Brazil appeared in first place in the number of publications between 1985 and 2015. Most of these studies were performed in the states of São Paulo, Minas Gerais and Paraná. Many species of oligochaetes are endemic to tropical regions, mainly in South America (Christoffersen, 2010). Studies to take inventory of fauna and learn the ecological relations between species and the environment are very important to formulate policies to preserve biodiversity, especially in forest regions under strong threat such as the Atlantic Forest fragments in Brazil.

CONCLUSION

According to the literature review, some tendencies and gaps exist in knowledge about freshwater oligochaetes that deserve attention. First of all, although there was a good deal of yearly upward and downward fluctuation in the number of works published, the trend considering the totals in the three 10-year periods was increasing, albeit modestly. This increase in the general number of publications was accompanied by specific increases in certain areas of knowledge, such as ecotoxicology, genetics/phylogeny and taxonomy. Descriptions, redistributions and reclassifications of species became more frequent, spurred by advances in molecular techniques, applied alone or in combination with morphological investigation, through interactive taxonomy, an important tool in various areas of biology. Nevertheless, despite the modest increase in the number of publications, studies of freshwater oligochaetes are still scarce. There are few specialists devoted to studying freshwater oligochaetes, leading to many gaps regarding geographic region and habitat focus. The Southern Hemisphere has been under-represented in studies of these organisms, mainly in countries of Africa and Oceania, while springs, caves and swamps are relatively neglected habitats. Therefore, more varied investigation is necessary to deepen understanding of freshwater oligochaetes and provide insights about their global diversity, and to guide efforts to protect their habitats.

In closing, if readers want to start studying or deepen their studies of aquatic oligochaetes, we suggest Reynolds & Wetzel (2018). The site is very comprehensive, containing updated references, nomenclature and species counts as well as news about events and related sites.

ACKNOWLEDGEMENTS

We would like to thank the anonymous reviewers for their suggestions and contributions to this manuscript, the Brazilian Office to Improve University Personnel (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – CAPES) for the scholarship grant, and the

the National Council for Scientific and Technological Development (Conselho Nacional de Desenvolvimento Científico e Tecnológico – CNPq) for funding (Process: 304552/2015-1).

REFERENCES

- Alves, R.G. & Lucca, J.V. 2000. Oligochaeta (Annelida: Clitellata) como indicador de poluição orgânica em dois córregos pertencentes à Bacia do Ribeirão do Ouro Araraquara (São Paulo, Brasil). *Brazilian Journal of Ecology*, 1: 112-117.
- Alves, R.G.; Marchese, M.R. & Escarpinatti, S.C. 2006. Oligochaeta (Annelida, Clitellata) in lotic environments in the state of São Paulo, Brazil. *Iheringia Série Zoologia*, 96: 431-435.
- Alves, R.G.; Marchese, M.R. & Martins, R.T. 2008. Oligochaeta (Annelida, Clitellata) of lotic environments at Parque Estadual Intervales (São Paulo, Brazil). *Biota Neotropica*, 8: 21-25.
- Behrend, R.D.L.; Fernandes, S.E.P.; Fujita, D.S. & Takeda, A.M. 2009. Eight years of monitoring aquatic Oligochaeta from the Baía and Ivinhema Rivers. *Brazilian Journal of Biology*, 69: 559-571.
- Behrend, R.D.L.; Takeda, A.M.; Gomes, L.C. & Fernandes, S.E.P. 2012. Using oligochaeta assemblages as an indicator of environmental changes. *Brazilian Journal of Biology*, 72: 873-884.
- Brinkhurst, R.O. 1999. Retrospect and prospect: reflections on forty years of study of aquatic oligochaetes. *Hydrobiologia*, 406: 9-19.
- Cantonati, M.; Füreder, L.; Gerecke, R.; Jüttner, I. & Cox, E.J. 2012. Crenic habitats, hotspots for freshwater biodiversity conservation: toward an understanding of their ecology. *Freshwater Science*, 31: 463-480.
- Christoffersen, M.L. 2010. Continental biodiversity of South American oligochaetes: the importance of inventories. *Acta Zoologica Mexicana*, 2: 35-46.
- Dumnicka, E. 2006. Composition and abundance of oligochaetes (Annelida: Oligochaeta) in springs of Kraków-Częstochowa Upland (Southern Poland): effect of spring encasing and environmental factors. *Polish Journal of Ecology*, 54: 231-242.
- Gavrilov, K. 1981. Oligochaeta. In: Hurlbert, S.H.; Rodriguez, G. & Santos, N.D. (Eds.). *Aquatic biota of tropical South America*, 2: Anarthropoda. San Diego, San Diego State University. p. 170-190.
- Giani, N.; Sambugar, B.; Rodriguez, P. & Martínez-Ansemil, E. 2001. Oligochaetes in southern European groundwater: new records and an overview. *Hydrobiologia*, 463: 65-74.
- Gorni, G.R. & Alves, R.G. 2006. Naididae (Annelida, Oligochaeta) associated with *Pomacea bridgesii* (Reeve) (Gastropoda, Ampullariidae). *Revista Brasileira de Zoologia*, 23: 1059-1061.
- Gorni, G.R. & Alves, R.G. 2008a. Naididae species (Annelida: Oligochaeta) associated with the sponge *Metania spinata* (Carter, 1881) (Porifera: Metaniidae) from a southeastern Brazilian reservoir. *Acta Limnologica Brasiliensis*, 20: 261-263.
- Gorni, G.R. & Alves, R.G. 2008b. Oligochaeta (Annelida: Clitellata) em córregos de baixa ordem do Parque Estadual de Campos do Jordão (São Paulo, Brasil). *Biota Neotropica*, 8: 161-165.
- Gorni, G.R. & Alves, R.G. 2012. Oligochaetes (Annelida, Clitellata) in a neotropical stream: a mesohabitat approach. *Iheringia Série Zoologia*, 102: 106-110.
- Gorni, G.R. & Alves, R.G. 2014. Influência de variáveis ambientais sobre a comunidade de oligoquetos (Annelida: Clitellata) em um córrego neotropical. *Biota Neotropica*, 28: 59-66.
- Gorni, G.R.; Peiró, D.F. & Sanches, N. 2015. Aquatic Oligochaeta (Annelida: Clitellata) from State of São Paulo, Brazil: diversity and occurrence review. *Biota Neotropica*, 15: 1-8.
- Lima-Ribeiro, M.S.; Nabout, J.C.; Pinto, M.P.; Moura, I.O.; Melo, T.L.; Costa, S.S. & Rangel, T.F.L.V.B. 2007. Análise cirométrica em ecologia de populações: importância e tendências dos últimos 60 anos. *Acta Scientiarum, Biological Sciences*, 29: 39-47.
- Lobo, H. & Alves, R.G. 2011a. Reproductive cycle of *Branchiura sowerbyi* (Oligochaeta: Naididae: Tubificinae) cultivated under laboratory conditions. *Zoologia*, Curitiba, 28: 427-431.
- Lobo, H. & Alves, R.G. 2011b. Influence of body weight and substrate granulometry on the reproduction of *Limnodrilus hoffmeisteri* (Oligochaeta: Naididae: Tubificinae). *Zoologia*, Curitiba, 28: 558-564.
- Lopes, S.; Costa, M.T.; Fernández-Llimós, F.; Amante, M.J. & Lopes, P.F. 2012. A Bibliometria e a Avaliação da Produção Científica: indicadores e ferramentas. In: Congresso Nacional de Bibliotecários, Arquivistas e Documentalistas, 11º. Actas. Lisboa, B.A.D.
- Martin, P.; Martínez-Ansemil, E.; Pinder, A.; Timm, T. & Wetzel, M.J. 2008. Global diversity of *Oligochaetous clitellates* ("Oligochaeta"; Clitellata) in freshwater. *Hydrobiologia*, 595: 117-127.
- Martins, R.T. & Alves, R.G. 2008. Occurrence of Naididae (Annelida: Oligochaeta) from three gastropod species in irrigation fields in southeastern Brazil. *Biota Neotropica*, 8: 255-257.
- Martins, R.T. & Alves, R.G. 2010. Occurrence of *Chaetogaster limnaei* K. von Baer, 1927 (Oligochaeta, Naididae) associated with Gastropoda mollusks in horticultural channels in Southeastern Brazil. *Brazilian Journal of Biology*, 70: 1055-1057.
- Nicacio, G. & Juen, L. 2015. Chironomids as indicators in freshwater ecosystems: an assessment of the literature. *Insect Conservation and Diversity*, 8(5): 393-403.
- Oda, F.H.; Petsch, D.K.; Ragonha, F.H.; Batista, V.G.; Takeda A.M. & Takemoto, R.M. 2015. *Dero (Allodero) lutzi* Michaelsen, 1926 (Oligochaeta: Naididae) associated with *Scinax fuscovarius* (Lutz, 1925) (Anura: Hylidae) from Semi-deciduous Atlantic Rain Forest, southern Brazil. *Brazilian Journal of Biology*, 75: 86-90.
- Petsch, D.K.; Ragonha, F.H.; Gimenez, B.C.G.; Barboza, L.G.A. & Takeda, A.M. 2015. Partitioning beta diversity of aquatic Oligochaeta in different environments of a Neotropical floodplain. *Acta Scientiarum. Biological Sciences*, 37: 41-49.
- Pinder, A.M.; Eberhard, S.M. & Humphreys, W.F. 2006. New phalodrilines (Annelida: Clitellata: Tubificidae) from Western Australian groundwater. *Zootaxa*, 1304: 31-48.
- Ragonha, F.H. & Takeda, A.M. 2014. Does richness of Oligochaeta (Annelida) follows a linear distribution with habitat structural heterogeneity in aquatic sediments? *Journal of Limnology*, 73: 146-156.
- Ragonha, F.H.; Petsch, D.K.; Alves, G.H.Z.; Santana, H.S.; Michelan, T.S. & Takeda, A.M. 2014. Tributaries as richness source for Oligochaeta assemblage (Annelida) of Neotropical dammed river. *Brazilian Journal of Biology*, 74: 861-869.
- Reynolds, J.W. & Wetzel, M.J. 2018. *Nomenclatura Oligochaetologica – A catalogue of names, descriptions and type specimens*. 2.ed. <http://www.inhs.illinois.edu/people/mjwetzel/nomenoligo>.
- Righi, G. 1984. *Manual de identificação de invertebrados límnicos do Brasil*. Brasília, CNPq/Coordenação Editorial.
- Rodrigues, L.F.T.; Leite, F.S.L. & Alves, R.G. 2013. Inventory and distribution of Oligochaeta (Annelida, Clitellata) in first-order streams in preserved areas of the state of Minas Gerais, Brazil. *Biota Neotropica*, 13: 245-254.
- Rodrigues, L.F.T.; Leite, F.S.L. & Alves, R.G. 2016. Influence of bryophyte biomass and organic matter quantity on the abundance and richness of oligochaetes in forest streams with different phytophysiognomies in southeastern Brazil. *Journal of Limnology*, 75: 340-346.
- Rodrigues, L.F.T.; Rosa, B.F.J.V.; Lobo, H.; Divino, A.C. & Alves, R.G. 2015. Diversity and distribution of oligochaetes in tropical forested streams, southeastern Brazil. *Journal of Limnology*, 74: 433-443.

- Rodriguez, P. & Reynoldson, T.B. 2011. *The pollution biology of aquatic oligochaetes*. New York, Springer Science & Business Media.
- Sambugar, B. 2007. Oligochetes from Alpine springs: a review. In: Cantonati M.; Bertuzzi, E. & Spitale, D. (Eds.). *The spring habitat: biota and sampling methods*. Trento, Museo Tridentino di Scienze Naturali. p. 185-192.
- Wetzel, M.J.; Fend, S.V.; Coates, K.A.; Kathman, R.D. & Gelder, S.R. 2006. *Taxonomy, systematics, and ecology of the aquatic Oligochaeta and Branchiobdellidae (Annelida, Clitellata) of North America, with emphasis*

on the fauna occurring in Florida. A workbook. 10 September. Illinois Natural History Survey, Champaign.

Wetzel, M.J.; Kathman, R.D.; Fend, S.V. & Coates, K.A. 2007. *Classification and Checklist of the Freshwater Oligochaeta occurring in North America North of Mexico*. Available at: <http://www.inhs.illinois.edu/~mjwetzel/FWOligoNAChlst.html>. Access in: 15/02/2017.

SUPPLEMENTARY MATERIAL

Additional Supporting Information may be found in the online version of this article:

This list contains all the references consulted and utilized to obtain the data in the article “Global panorama of studies about freshwater oligochaetes: main trends and gaps”.

- The list of SciELO contains references found only in this database (7 references).
- The list of Web of Science contains references found only in this database (351 references).
- The list of Scopus contains references found in this database and those that were common with the SciELO or Web of Science bases (621 references).



SCIELO

- Armendáriz, L.C. 2007. Efectos de la temperatura sobre el crecimiento individual y formación de zooides en *Stylaria lacustris* (Linnaeus, 1767) (Oligochaeta, Naididae). *Biota Neotropica*, 7: 77-81.
- Christoffersen, M.L. 2010. Continental biodiversity of South American oligochaetes: The importance of inventories. *Acta Zoológica Mexicana*, 26: 35-46.
- Corbi, J.J.; Jancso, M.A.; Strixino, S.T. & Fragoso, E.N. 2004. Occurrence of Oligochaeta living on larvae of Odonata from Ipeúna (São Paulo state, Brazil). *Biota Neotropica*, 4: 1-3.
- Cortelezzi, A.; Armendáriz, L.C.; López van Oosterom, M.V.; Cepeda, R. & Capítulo, A.R. 2012. Different levels of taxonomic resolution in bioassessment: a case study of Oligochaeta in lowland streams. *Acta Limnologica Brasiliensis*, 23: 412-425.
- Fujita, D.S.; Takeda, A.M.; Coutinho, R. & Fernandes, F.C. 2015. Influence of antifouling paint on freshwater invertebrates (Mytilidae, Chironomidae and Naididae): density, richness and composition. *Brazilian Journal of Biology*, 75: 70-78.
- Pamplin, P.A.Z.; Rocha, O. & Marchese, M. 2005. Riqueza de espécies de Oligochaeta (Annelida, Clitellata) em duas represas do Rio Tietê (São Paulo). *Biota Neotropica*, 5: 63-70.
- Righi, G. & Hamoui, V. 2002. Oligochaeta, Naididae of the West Indies and adjacent regions. *Papéis Avulsos de Zoologia*, 42: 119-167.

WEB OF SCIENCE

- Achurra, A., Rodriguez, P. & Erseus, C. 2015. Pseudo-cryptic speciation in the subterranean medium: A new species of *Stylodrilus Claparede*, 1862, with a revision of the status of *Bichaeta Bretscher*, 1900 (Annelida, Clitellata, Lumbriculidae). *Zoologischer Anzeiger*, 257, 71-86.
- Agbo, S.O., Keinanen, M., Keski-Saari, S., Lemmettyinen, J., Akkanen, J., Leppanen, M.T., Mayer, P. & Kukkonen, J.V.K. 2013a. Changes in *Lumbriculus variegatus* metabolites under hypoxic exposure to benzo(a)pyrene, chlorpyrifos and pentachlorophenol: Consequences on biotransformation. *Chemosphere*, 93(2), 302-310.
- Agbo, S.O., Lemmettyinen, J., Keinanen, M., Keski-Saari, S., Akkanen, J., Leppanen, M.T., Wang, Z.X., Wang, H.L., Price, D.A. & Kukkonen, J.V.K. 2013b. Response of *Lumbriculus variegatus* transcriptome and metabolites to model chemical contaminants. *Comparative Biochemistry and Physiology – Part C: Toxicology and Pharmacology*, 157(2), 183-191.
- Ahamed, M.T. & Mollah, M.F.A. 1992. Effects of various levels of wheat bran and mustard oil cake in the culture media on tubificid production. *Aquaculture*, 107(1), 107-113.
- Akinschina, T.V. & Snimschikova, L.N. 1991. Isochaetides-werestschagini sp. nov. (Oligochaeta, Tubificidae). *Zoologichesky Zhurnal*, 70(7), 130-132.
- Anlauf, A. 1990. Cyst formation of *Tubifex tubifex* (Müller) – an adaptation to survive food deficiency and drought. *Hydrobiologia*, 190(1), 79-82.
- Anlauf, A. & Neumann, D. 1997. The genetic variability of *Tubifex tubifex* (Müller) in 20 populations and its relation to habitat type. *Archiv Fur Hydrobiologie*, 139(2), 145-162.
- Anlauf, K.J. & Moffitt, C.M. 2008. Models of stream habitat characteristics associated with tubificid populations in an intermountain watershed. *Hydrobiologia*, 603, 147-158.
- Anlauf, K.J. & Moffitt, C.M. 2010. Modelling of landscape variables at multiple extents to predict fine sediments and suitable habitat for *Tubifex tubifex* in a stream system. *Freshwater Biology*, 55(4), 794-805.
- Anschutz, P., Ciutat, A., Lecroart, P., Gerino, M. & Boudou, A. 2012. Effects of Tubificid Worm Bioturbation on Freshwater Sediment Biogeochemistry. *Aquatic Geochemistry*, 18(6), 475-497.
- Arrendarczyk, A., Jakubowska, A., Zgorska, A. & Grabinska-Sota, E. 2014. Toxic effects of cadmium-spiked sediments in *Tubifex tubifex*: enzyme biomarkers measurements. *Desalination and Water Treatment*, 52(19-21), 3798-3803.
- Armendariz, L.C. 2000. Population dynamics of *Stylaria lacustris* (Linnaeus, 1767) (Oligochaeta, Naididae) in Los Talas, Argentina. *Hydrobiologia*, 438(1-3), 217-226.
- Arsan, E.L., Hallett, S.L. & Bartholomew, J.L. 2007. *Tubifex tubifex* from Alaska and their susceptibility to *Myxobolus cerebralis*. *Journal of Parasitology*, 93(6), 1332-1342.
- Artola-Garicano, E., Sinnige, T.L., van Holsteijn, I., Vaes, W.H.J. & Hermens, J.L.M. 2003. Bioconcentration and acute toxicity of polycyclic musks in two benthic organisms (*Chironomus riparius* and *Lumbriculus variegatus*). *Environmental Toxicology and Chemistry*, 22(5), 1086-1092.
- Atanackovic, A., Sporka, F., Tomovic, J., Vasiljevic, B., Markovic, V., Simic, V. & Paunovic, M. 2012. First record of *Bothrioneurum vejdovskyanum Štolc*, 1886 (Oligochaeta, Tubificidae) in Serbia. *Archives of Biological Sciences*, 64(3), 1123-1126.
- Back, H. 1990. Epidermal uptake of PB, CD, and ZN in Tubificid worms. *Oecologia*, 85(2), 226-232.
- Back, H. & Prosi, F. 1985. Distribution of inorganic cations in *Limnodrilus udekemianus* (Oligochaeta, Tubificidae) using laser-induced microprobe mass analysis, with special emphasis on heavy-metals. *Micron and Microscopica Acta*, 16(3), 145-150.
- Basim, Y., Farzadkia, M., Jaafarzadeh, N. & Hendrickx, T. 2012. Sludge reduction by *Lumbriculus variegatus* in Ahvas wastewater treatment plant. *Iranian Journal of Environmental Health Science & Engineering*, 9.
- Baturina, M., Timm, T. & Loskutova, O. 2014. Oligochaete (Annelida, Clitellata) communities in lakes of the Ural Mountains (Russia), in Pinder, A., Arslan, N. & Wetzel, M. (Eds.). *Proceedings of the 12th International Symposium on Aquatic Oligochaeta, Zoosymposia*, 77-94.
- Beauchamp, K.A., Kelley, G.O., Nehring, R.B. & Hedrick, R.P. 2005. The severity of whirling disease among wild trout corresponds to the differences in genetic composition of *Tubifex tubifex* populations in central Colorado. *Journal of Parasitology*, 91(1), 53-60.
- Beaudet, J.F., Prevost, M., Niquette, P., Clement, B., Coallier, J. & Arcouette, N. 2000. Simple methods for evaluating the population of naidids in drinking water treatment plants. *Canadian Journal of Civil Engineering*, 27(4), 601-609.
- Beckett, D.C., Aartila, T.P. & Miller, A.C. 1992. Seasonal change in plant-dwelling Chironomidae and Naididae in a Wisconsin Lake. *Journal of Freshwater Ecology*, 7(1), 45-57.
- Beckingham, B. & Ghosh, U. 2010. Comparison of field and laboratory exposures of *Lumbriculus variegatus* to polychlorinated biphenyl-impacted river sediments. *Environmental Toxicology and Chemistry*, 29(12), 2851-2858.
- Belden, J.B., Lotufo, G.R. & Lydy, M.J. 2005. Accumulation of hexahydro-1,3,5-trinitro-1,3,5-triazine in channel catfish (*Ictalurus punctatus*) and aquatic oligochaetes (*Lumbriculus variegatus*). *Environmental Toxicology and Chemistry*, 24(8), 1962-1967.
- Bely, A.E. & Wray, G.A. 2004. Molecular phylogeny of naidid worms (Annelida: Clitellata) based on cytochrome oxidase I. *Molecular Phylogenetics and Evolution*, 30(1), 50-63.
- Bervoets, L., Blust, R., deWit, M. & Verheyen, R. 1997. Relationships between river sediment characteristics and trace metal concentrations in Tubificid worms and chironomid larvae. *Environmental Pollution*, 95(3), 345-356.

- Bettinetti, R. & Provini, A. 2002. Toxicity of 4-nonylphenol to *Tubifex tubifex* and *Chironomus riparius* in 28-day whole-sediment tests. *Ecotoxicology and Environmental Safety*, 53(1), 113-121.
- Blazer, V.S., Waldrop, T.B., Schill, W.B., Densmore, C.L. & Smith, D. 2003. Effects of water temperature and substrate type on spore production and release in eastern *Tubifex tubifex* worms infected with *Myxobolus cerebralis*. *Journal of Parasitology*, 89(1), 21-26.
- Bonacina, C., Bonomi, G. & Monti, C. 1989a. Density-dependent processes in cohorts of *Tubifex tubifex*, with special emphasis on the control of fecundity. *Hydrobiologia*, 180, 135-141.
- Bonacina, C., Bonomi, G. & Monti, C. 1989b. Population analysis in mass-cultures of *Tubifex tubifex*. *Hydrobiologia*, 180, 127-134.
- Bonacina, C., Pasteris, A., Bonomi, G. & Marzuoli, D. 1994. Quantitative observations on the population ecology of *Branchiura sowerbyi* (Oligochaeta, Tubificidae). *Hydrobiologia*, 278(1-3), 267-274.
- Bonacina, C., Pasteris, A., DiCola, G. & Bonomi, G. 1996. Production and population dynamics of *Tubifex tubifex* in the profundal zone of a freshwater reservoir in N Italy. *Hydrobiologia*, 334(1-3), 141-146.
- Bouche, M.L., Arnoult, F. & Vernet, G. 2003. Caudal regeneration in *Tubifex tubifex* (Oligochaeta, Tubificidae) following copper exposure. *Invertebrate Biology*, 122(1), 42-51.
- Bouche, M.L., Biagiotti-Risbourg, S. & Vernet, G. 1999. A light and scanning electron microscope study of the morphology of the chaetae of *Tubifex tubifex* in a non-polluted medium. *Hydrobiologia*, 411, 39-44.
- Bouche, M.L., Habets, F., Biagiotti-Risbourg, S. & Vernet, G. 2000. Toxic effects and bioaccumulation of cadmium in the aquatic oligochaete *Tubifex tubifex*. *Ecotoxicology and Environmental Safety*, 46(3), 246-251.
- Brinkhurst, R.O. 1985a. The generic and subfamilial classification of the Naididae (Annelida, Oligochaeta). *Proceedings of the Biological Society of Washington*, 98(2), 470-475.
- Brinkhurst, R.O. 1985b. *Varichaetadrilus-minutus* (Brinkhurst, 1965) new combination for psammoryctides, minutus (Oligochaeta, Tubificidae). *Proceedings of the Biological Society of Washington*, 98(3), 655-656.
- Brinkhurst, R.O. 1987. Notes on *Varichaetadrilus* Brinkhurst and Kathman, 1983 (Oligochaeta, Tubificidae). *Proceedings of the Biological Society of Washington*, 100(3), 515-517.
- Brinkhurst, R.O. 1989a. A phylogenetic analysis of the Lumbriculidae (Annelida, Oligochaeta). *Canadian Journal of Zoology*, 67(11), 2731-2739.
- Brinkhurst, R.O. 1989b. *Varichaetadrilus-angustipenis* Brinkhurst & Cook, 1966, new combination for *Limnodrilus angustipenis* (Oligochaeta, Tubificidae). *Proceedings of the Biological Society of Washington*, 102(2), 311-312.
- Brinkhurst, R.O. 1991. A phylogenetic analysis of the Tubificinae (Oligochaeta, Tubificidae). *Canadian Journal of Zoology*, 69(2), 392-397.
- Brinkhurst, R.O. & Barbour, M.T. 1985. A new species of *Aulodrilus* Bretscher (Oligochaeta, Tubificidae) from North America. *Proceedings of the Biological Society of Washington*, 98(4), 931-934.
- Brinkhurst, R.O. & Coates, K.A. 1985. The genus *paranais* (Oligochaeta, Naididae) in North America. *Proceedings of the Biological Society of Washington*, 98(2), 303-313.
- Brinkhurst, R.O. & Diaz, R.J. 1985. *Isochaetides columbiensis*, new species (Oligochaeta, Tubificidae) from the Columbia River, Oregon. *Proceedings of the Biological Society of Washington*, 98(4), 949-953.
- Brinkhurst, R.O. & Marchese, M. 1987. A contribution to the taxonomy of the aquatic Oligochaeta (Haplotaenidae, Phreodrilidae, Tubificidae) of South America. *Canadian Journal of Zoology – Revue Canadienne de Zoologie*, 65(12), 3154-3165.
- Brinkhurst, R.O., Rodriguez, P., Chon, T.S. & Kwon, T.S. 1994. A new genus of Lumbriculidae (Oligochaeta) from Korea. *Canadian Journal of Zoology – Revue Canadienne de Zoologie*, 72(11), 1960-1966.
- Bunke, D. 2003. Early development of metanephridia in the caudal budding zone of a clitellate annelid, *Dero digitata* (Naidida): an electron-microscopical study. *Acta Zoologica*, 84(2), 87-97.
- Burkhard, L.P., Hubin-Barrows, D., Billa, N., Highland, T.L., Hockett, J.R., Mount, D.R., Norberg-King, T.J., Hawthorne, S., Miller, D.J. & Grabanski, C.B. 2015. Sediment Bioaccumulation Test with *Lumbriculus variegatus*: Effects of Feeding. *Archives of Environmental Contamination and Toxicology*, 68(4), 696-706.
- Burkhard, L.P., Mount, D.R., Highland, T.L., Hockett, J.R., Norberg-King, T., Billa, N., Hawthorne, S.B., Miller, D.J. & Grabanski, C.B. 2013. Evaluation of PCB bioaccumulation by *Lumbriculus variegatus* in field-collected sediments. *Environmental Toxicology and Chemistry*, 32(7), 1495-1503.
- Camusso, M., Polesello, S., Valsecchi, S. & Vignati, D.A.L. 2012a. Importance of dietary uptake of trace elements in the benthic deposit-feeding *Lumbriculus variegatus*. *TRAC – Trends in Analytical Chemistry*, 36, 103-112.
- Camusso, M., Polesello, S., Valsecchi, S. & Vignati, D.A.L. 2012b. Importance of dietary uptake of trace elements in the benthic deposit-feeding *Lumbriculus variegatus*. *TRAC – Trends in Analytical Chemistry*, 36, 103-112.
- Casellato, S., Covre, V. & Del Piero, S. 2014. Effects of sodium fluoride on the gametogenesis of the tubificid oligochaete *Branchiura sowerbyi* Beddard, in Pinder, A., Arslan, N. & Wetzel, M. (Eds.). *Proceedings of the 12th International Symposium on Aquatic Oligochaeta, Zoosymposia*, 51-58.
- Casellato, S., Martinucci, G. & Zoja, E. 1987. Ultrastructural features of gametogenesis during the life-cycle in *Branchiura sowerbyi* Beddard (Oligochaeta, Tubificidae). *Hydrobiologia*, 155, 145-154.
- Casellato, S. & Negrisolo, P. 1989. Acute and chronic effects of an anionic surfactant on some fresh-water tubificid species. *Hydrobiologia*, 180, 243-252.
- Chen, T., Furst, A. & Chien, P.K. 1994. The effects of cadmium and iron on catalase activities in *Tubifex*. *Journal of the American College of Toxicology*, 13(2), 112-120.
- Chuko, G.M., Lapkina, L.N. & Podgornaya, V.A. 2011. Substrate-inhibitory analysis of the cholinesterase in the freshwater oligochaete *Lumbriculus variegatus* (of Müller, 1773; Lumbriculidae, Oligochaeta). *Biochemical Systematics and Ecology*, 39(3), 169-174.
- Ciutat, A., Gerino, M. & Boudou, A. 2007. Remobilization and bioavailability of cadmium from historically contaminated sediments: Influence of bioturbation by Tubificids. *Ecotoxicology and Environmental Safety*, 68(1), 108-117.
- Ciutat, A., Gerino, M., Mesmer-Dudons, N., Anschutz, P. & Boudou, A. 2005. Cadmium bioaccumulation in Tubificidae from the overlying water source and effects on bioturbation. *Ecotoxicology and Environmental Safety*, 60(3), 237-246.
- Ciutat, A., Weber, O., Gerino, M. & Boudou, A. 2006. Stratigraphic effects of Tubificids in freshwater sediments: a kinetic study based on X-Ray images and grain-size analysis. *Acta Oecologica – International Journal of Ecology*, 30(2), 228-237.
- Coleman, J.G., Kennedy, A.J., Bednar, A.J., Ranville, J.F., Laird, J.G., Harmon, A.R., Hayes, C.A., Gray, E.P., Higgins, C.P., Lotufo, G. & Steevens, J.A. 2013. Comparing the effects of nanosilver size and coating variations on bioavailability, internalization, and elimination, using *Lumbriculus variegatus*. *Environmental Toxicology and Chemistry*, 32(9), 2069-2077.
- Coler, R.A., Coler, M.S. & Kostecki, P.T. 1988. Tubificid behavior as a stress indicator. *Water Research*, 22(2), 263-267.
- Contardo-Jara, V., Klingelmann, E. & Wiegand, C. 2009. Bioaccumulation of glyphosate and its formulation Roundup Ultra in *Lumbriculus variegatus* and its effects on biotransformation and antioxidant enzymes. *Environmental Pollution*, 157(1), 57-63.

- Crisp, K.M., Grupe, R.E., Lobsang, T.T. & Yang, X. 2010. Biogenic amines modulate pulse rate in the dorsal blood vessel of *Lumbriculus variegatus*. *Comparative Biochemistry and Physiology – Part C: Toxicology and Pharmacology*, 151(4), 467-472.
- Crottini, A., Marotta, R., Barbuto, M., Casiraghi, M. & Ferraguti, M. 2008. The world in a river? A preliminary analysis of the 16S rDNA variability of *Tubifex* species (Clitellata: Tubificidae) from the Lambro River. *Molecular Phylogenetics and Evolution*, 48(3), 1189-1203.
- Cuadrado, S. & Martinez-Ansemil, E. 2001. External structures used during attachment and sperm transfer in Tubificids (Annelida, Oligochaeta). *Hydrobiologia*, 463, 107-113.
- Cui, Y.D., He, X.B., Peng, Y. & Wang, H.Z. 2015. Records of Naididae and Lumbriculidae (Clitellata) from Tibet, China, with description of a new species of *Nais*. *Zootaxa*, 3956(4), 513-530.
- Dawson, T.D., Lott, K.G., Leonard, E.N. & Mount, D.R. 2003. Time course of metal loss in *Lumbriculus variegatus* following sediment exposure. *Environmental Toxicology and Chemistry*, 22(4), 886-889.
- Deeguileor, M., Lanzavecchia, G., Valvassori, R. & Lanzavecchia, P. 1987. Unusual model of lumbriculids helical muscles – comparison with body wall muscles in other microdriles. *Hydrobiologia*, 155, 135-144.
- Del Piero, S., Masiero, L. & Casellato, S. 2014. Toxicity and bioaccumulation of fluoride ion on *Branchiura sowerbyi*, Beddard, (Oligochaeta, Tubificidae), in Pinder, A., Arslan, N. & Wetzel, M. (Eds.). *Proceedings of the 12th International Symposium on Aquatic Oligochaeta, Zoosymposia*, 44-50.
- Delmotte, S., Meysman, F.J.R., Ciutat, A., Boudou, A., Sauvage, S. & Gerino, M. 2007. Cadmium transport in sediments by tubificid bioturbation: An assessment of model complexity. *Geochimica et Cosmochimica Acta*, 71(4), 844-862.
- Dermott, R. & Munawar, M. 1992. A simple and sensitive assay for evaluation of sediment toxicity using *Lumbriculus variegatus* (Müller). *Hydrobiologia*, 235, 407-414.
- Di Chiara, A.P., Pagani, M. & Leoni, B. 1999. Enzymatic variability in sympatric species of *Tubifex* (Clitellata: Tubificidae). *Hydrobiologia*, 406, 83-86.
- Di, S.S., Liu, T.T., Diao, J.L. & Zhou, Z.Q. 2013. Enantioselective bioaccumulation and degradation of sediment-associated metalaxyl enantiomers in *Tubifex tubifex*. *Journal of Agricultural and Food Chemistry*, 61(21), 4997-5002.
- Dierickx, P.J. 1985. Anionic and neutral glutathione s-transferase isoenzymes in the fresh-water worm *Tubifex tubifex* (OFM). *Archives Internationales de Physiologie de Biochimie et de Biophysique*, 93(3), 193-198.
- Dolidze, T.M. 1994. Biology of *Limnodrilus claparedeanus ratzeli* (Oligochaeta, Tubificidae) in the Tsimlyansk Reservoir. *Hydrobiologia*, 278(1-3), 275-279.
- Drewes, C.D. 1999. Helical swimming and body reversal behaviors in *Lumbriculus variegatus* (Annelida: Clitellata: Lumbriculidae). *Hydrobiologia*, 406, 263-269.
- Drewes, C.D. & Brinkhurst, R.O. 1990. Giant nerve-fibers and rapid escape reflexes in newly hatched aquatic oligochaetes, *Lumbriculus variegatus* (family Lumbriculidae). *Invertebrate Reproduction & Development*, 17(2), 91-95.
- Drewes, C.D. & Fourtner, C.R. 1989. Hindsight and rapid escape in a fresh-water oligochaete. *Biological Bulletin*, 177(3), 363-371.
- Drewes, C.D. & Fourtner, C.R. 1990. Morphallaxis in an aquatic oligochaete, *Lumbriculus variegatus* – reorganization of escape reflexes in regenerating body fragments. *Developmental Biology*, 138(1), 94-103.
- Drewes, C.D. & Fourtner, C.R. 1993. Helical swimming in a fresh-water oligochaete. *Biological Bulletin*, 185(1), 1-9.
- Drewes, C.D. & Zoran, M.J. 1989. Neuro-behavioral specializations for respiratory movements and rapid escape from predators in posterior segments of the tubificid *Branchiura sowerbyi*. *Hydrobiologia*, 180, 65-71.
- DuBey, R. & Caldwell, C. 2004. Distribution of *Tubifex tubifex* lineages and *Myxobolus cerebralis* infection in the tailwater of the San Juan River, New Mexico. *Journal of Aquatic Animal Health*, 16(4), 179-185.
- DuBey, R., Caldwell, C. & Gould, W.R. 2005. Effects of temperature, photoperiod, and *Myxobolus cerebralis* infection on growth, reproduction, and survival of *Tubifex tubifex* lineages. *Journal of Aquatic Animal Health*, 17(4), 338-344.
- DuBey, R.J. 2008. Genetic differentiation of lineages of *Tubifex tubifex* from the San Juan River, New Mexico. *Southwestern Naturalist*, 53(2), 268-272.
- Ducrot, V., Pery, A.R.R., Queau, H., Mons, R., Lafont, M. & Garric, J. 2007. Rearing and estimation of life-cycle parameters of the tubificid worm *Branchiura sowerbyi*: Application to ecotoxicity testing. *Science of the Total Environment*, 384(1-3), 252-263.
- Dunnicka, E. 1986. Amsterdam expeditions to the west-indian-islands. 50. Naididae (Oligochaeta) from subterranean waters of west-indian-islands – distribution, taxonomic remarks and description of a new species. *Bijdragen Tot De Dierkunde*, 56(2), 267-281.
- Eckroth, M.C. & Brinkhurst, R.O. 1996. *Tenagodrilus musculus*, a new genus and species of Lumbriculidae (Clitellata) from a temporary pond in Alabama, USA. *Hydrobiologia*, 334(1-3), 1-9.
- Edwards, P.A. & Simkiss, K. 2005. Dietary influences on the bioaccumulation of pollutants by the annelid, *Lumbriculus variegatus*: Experiments comparing artificial particles and natural sediments. *Bulletin of Environmental Contamination and Toxicology*, 74(2), 328-334.
- Egeler, P., Rombke, J., Meller, M., Knacker, T., Franke, C., Studinger, G. & Nagel, R. 1997. Bioaccumulation of lindane and hexachlorobenzene by tubificid sludge worms (Oligochaeta) under standardised laboratory conditions. *Chemosphere*, 35(4), 835-852.
- Egeler, P., Rombke, J., Meller, M., Knacker, T. & Nagel, R. 1999. Bioaccumulation test with *Tubificid Sludgeworms* in artificial media – development of a standardisable method. *Hydrobiologia*, 406, 271-280.
- Elwell, L.C.S., Kerans, B.L. & Zickovich, J. 2009. Host-parasite interactions and competition between tubificid species in a benthic community. *Freshwater Biology*, 54(8), 1616-1628.
- Envall, I., Erseus, C. & Gustavsson, L.M. 2008. Ultrastructural investigation of coelomocytes in representatives of Naidinae and Rhyacodrilinae (Annelida, Clitellata, Tubificidae). *Journal of Morphology*, 269(9), 1157-1167.
- Envall, I., Gustavsson, L.M. & Erseus, C. 2012. Genetic and chaetal variation in *Nais* worms (Annelida, Clitellata, Naididae). *Zoological Journal of the Linnean Society*, 165(3), 495-520.
- Envall, I., Kallersjo, M. & Erseus, C. 2006. Molecular evidence for the non-monophyletic status of Naidinae (Annelida, Clitellata, Tubificidae). *Molecular Phylogenetics and Evolution*, 40(2), 570-584.
- Erseus, C. 1999. *Parvidrilus strayeri*, a new genus and species, an enigmatic interstitial clitellate from underground waters in Alabama. *Proceedings of the Biological Society of Washington*, 112(2), 327-337.
- Erseus, C. & Dunnicka, E. 1988. A new *Phallodrilus* (Oligochaeta, Tubificidae) from subterranean waters in central Italy. *Stygologia*, 4, 116-120.
- Erseus, C. & Gustavsson, L. 2002. A proposal to regard the former family Naididae as a subfamily within Tubificidae (Annelida, Clitellata). *Hydrobiologia*, 485(1-3), 253-256.
- Erseus, C., Kallersjo, M., Ekman, M. & Hovmoller, R. 2002. 18S rDNA phylogeny of the Tubificidae (Clitellata) and its constituent taxa: Dismissal of the Naididae. *Molecular Phylogenetics and Evolution*, 22(3), 414-422.
- Erseus, C., Prestegaard, T. & Kallersjo, M. 2000. Phylogenetic analysis of Tubificidae (Annelida, Clitellata) based on 18S rDNA sequences. *Molecular Phylogenetics and Evolution*, 15(3), 381-389.
- Famme, P. & Knudsen, J. 1985. Anoxic survival, growth and reproduction by the fresh-water annelid, *Tubifex* sp., demonstrated using a new simple

- anoxic chemostat. *Comparative Biochemistry and Physiology – Part A: Physiology*, 81(2), 251-253.
- Fara, D.G. & Erseus, C. 1991. *Phallodrilus-aquaedulcis hrabe*, 1960, a meiobenthic fresh-water oligochaete (Tubificidae) previously known only from Europe, recorded from the Niagara River, North America. *Canadian Journal of Zoology – Revue Canadienne de Zoologie*, 69(2), 291-294.
- Fargasova, A. 1994. Toxicity of metals on *Daphnia magna* and *Tubifex tubifex*. *Ecotoxicology and Environmental Safety*, 27(2), 210-213.
- Fargasova, A. 1997. Determination of toxicity of plant growth regulators in selected aquatic organisms *Daphnia magna* and *Tubifex tubifex*. *Biology*, 52(3), 405-408.
- Fargasova, A. 1998. Comparative acute toxicity of Cu²⁺, Cu⁺, Mn²⁺, Mo⁶⁺, Ni²⁺ and V⁵⁺ to *Chironomus plumosus* larvae and *Tubifex tubifex* worms. *Biology*, 53(3), 315-319.
- Fargasova, A. & Kizlink, J. 1996. Acute toxic effects of organotin compounds on benthic organisms: *Tubifex tubifex* and *Chironomus plumosus*. *Biology*, 51(6), 677-681.
- Fend, S.V. 2005. A review of the genus *Eclipidrilus* (Annelida: Clitellata: Lumbriculidae), with description of a new species from western North America. *Zootaxa*, (969), 1-42.
- Fend, S.V. & Gustafson, D.L. 2001. A new Lumbriculid genus and species from North America (Clitellata, Lumbriculidae). *Hydrobiologia*, 463, 13-22.
- Fend, S.V. & Lenat, D.R. 2007. Two new genera of Lumbriculidae (Annelida: Clitellata) from North Carolina, USA. *Zootaxa*, (1666), 1-22.
- Fend, S.V. & Ohtaka, A. 2004. *Yamaguchia toyensis* sp. nov., gen. nov. (Annelida, Clitellata, Lumbriculidae) from profundal lake habitat in Japan. *Zoological Science*, 21(6), 677-683.
- Ferraguti, M., Bondi, M. & Lanzavecchia, S. 1991. Ultrastructural analysis of sperm flagella in 2 clitellates (Annelida), plasma-membrane and periaxonemal area. *Tissue & Cell*, 23(4), 537-546.
- Ferraguti, M., Grassi, G. & Erseus, C. 1989. Different models of tubificid spermatozeugmata. *Hydrobiologia*, 180, 73-82.
- Ferraguti, M. & Jamieson, B.G.M. 1987. Spermiogenesis in bythonomus lemani and the phylogenetic position of the Lumbriculidae (Oligochaeta, Annelida). *Hydrobiologia*, 155, 123-134.
- Ferraguti, M. & Ruprecht, D. 1992. The double sperm line in the tubificid *Clitellio arenarius* (Annelida, Oligochaeta). *Bollettino di Zoologia*, 59(4), 349-362.
- Ferraguti, M., Ruprecht, D., Erseus, C. & Giere, O. 1994. An ultrastructural overview of tubificid spermatozoa. *Hydrobiologia*, 278(1-3), 165-178.
- Finogenova, N.P. 1986. Diagnostic characters of subfamilies in the family Tubificidae (Oligochaeta). *Zoologichesky Zhurnal*, 65(2), 194-202.
- Fischer, E. & Molnar, L. 1986. 3 Discrete stages of the nuclear volume alterations of the chloragocytes in experimentally treated *Limnodrilus hoffmeisteri* (Oligochaeta, Tubificidae). *Zoologischer Anzeiger*, 216(1-2), 99-108.
- Fischer, E. & Molnar, L. 1987. Karyometrical evidences of the role of chloragocytes in the salt balance mechanisms of the tubificid worm *Limnodrilus hoffmeisteri*. *Zoologischer Anzeiger*, 218(5-6), 374-378.
- Fischer, E., Molnar, L. & Szucs, I. 1986. Possible reasons for chloragocyte nuclear swelling under hypoxic conditions and under saline load in the tubificid worm *Limnodrilus hoffmeisteri*. *Zoologischer Anzeiger*, 217(3-4), 283-288.
- Fukuhara, H., Ohtaka, A., Isobe, Y. & Sakamoto, M. 1987. Seasonal-changes in vertical-distribution, biomass and fecal production of Tubificids in the profundal region of a shallow Japanese lake. *Holarctic Ecology*, 10(1), 8-13.
- Fukuhara, H. & Sakamoto, M. 1987. Enhancement of inorganic nitrogen and phosphate release from lake sediment by Tubificid worms and chironomid larvae. *Oikos*, 48(3), 312-320.
- Gainutdinov, M.K., Yargunov, V.G., Kalinnikova, T.B., Gainutdinov, T.M. & Varlamov, V.E. 2000. On the role of a decentralized control system in adaptation to high ambient temperature in a population of the oligochaete *Tubifex tubifex*. *Russian Journal of Ecology*, 31(6), 444-446.
- Galluba, S., Oetken, M. & Oehlmann, J. 2012. Comprehensive sediment toxicity assessment of Hessian surface waters using *Lumbriculus variegatus* and *Chironomus riparius*. *Journal of Environmental Science and Health – Part A: Toxic/Hazardous Substances & Environmental Engineering*, 47(4), 507-521.
- Geta, R., Postolache, C. & Vadineanu, A. 2004. Ecological significance of nitrogen cycling by tubificid communities in shallow eutrophic lakes of the Danube Delta. *Hydrobiologia*, 524(1), 193-202.
- Giani, N., Erseus, C. & Martinezsemil, E. 1990. Redefinition of the subterranean genus *Krenedrilus Dunnicka* (Oligochaeta, Tubificidae) with a redescription of *Krenedrilus papillatus Dunnicka*, 1983 and the description of 2 new species. *Stylogorgia*, 5(1), 55-65.
- Giani, N. & Rodriguez, P. 1988. Description of some new species of Tubificidae (Oligochaeta) from caves and karstic springs of the Iberian Peninsula. *Stylogorgia*, 4, 121-137.
- Gillis, P.L., Dixon, D.G., Borgmann, U. & Reynoldson, T.B. 2004a. Uptake and depuration of cadmium, nickel, and lead in laboratory-exposed *Tubifex tubifex* and corresponding changes in the concentration of a metallothionein-like protein. *Environmental Toxicology and Chemistry*, 23(1), 76-85.
- Gillis, P.L., Reynoldson, T.B. & Dixon, D.G. 2004b. Natural variation in a metallothionein-like protein in *Tubifex tubifex* in the absence of metal exposure. *Ecotoxicology and Environmental Safety*, 58(1), 22-28.
- Goto, A., Kitamura, K., Arai, A. & Shimizu, T. 1999a. Cell fate analysis of teloblasts in the *Tubifex* embryo by intracellular injection of HRP. *Development Growth & Differentiation*, 41(6), 703-713.
- Goto, A., Kitamura, K. & Shimizu, T. 1999b. Cell lineage analysis of pattern formation in the *Tubifex* embryo. I. Segmentation in the mesoderm. *International Journal of Developmental Biology*, 43(4), 317-327.
- Guerin, C. & Giani, N. 1996. Analytical study of the locomotor and respiratory movements of Tubificid worms by means of video recording. *Hydrobiologia*, 333(1), 63-69.
- Gunn, A.M., Hunt, D.T.E. & Winnard, D.A. 1989. The effect of heavy-metal speciation in sediment on bioavailability to Tubificid worms. *Hydrobiologia*, 188, 487-496.
- Guo, X.S., Liu, J.X., Wei, Y.S. & Li, L. 2007. Sludge reduction with Tubificidae and the impact on the performance of the wastewater treatment process. *Journal of Environmental Sciences*, 19(3), 257-263.
- Gustafsson, D.R., Price, D.A. & Erseus, C. 2009. Genetic variation in the popular lab worm *Lumbriculus variegatus* (Annelida: Clitellata: Lumbriculidae) reveals cryptic speciation. *Molecular Phylogenetics and Evolution*, 51(2), 182-189.
- Gutierrez, R.M.P. & Reyes, I.C. 2006. Sterols isolated from *Tubifex tubifex*. *Natural Product Research*, 20(13), 1271-1276.
- Hall, S., Lockwood, R. & Harrass, M.C. 2014. Application of a unique test design to determine the chronic toxicity of boron to the aquatic worm *Lumbriculus variegatus* and Fatmucket mussel *Lampsilis siliquoidea*. *Archives of Environmental Contamination and Toxicology*, 66(1), 58-68.
- Hamilton, A.J. & Canning, E.U. 1987. Studies on the proposed role of *Tubifex tubifex* (Müller) as an intermediate host in the life-cycle of *Myxosoma-cerebralis* (Hofer, 1903). *Journal of Fish Diseases*, 10(3), 145-151.
- Harman, W.J., Brinkhurst, R.O. & Marchese, M. 1988. A contribution to the taxonomy of the aquatic Oligochaeta (Naididae) of South America. *Canadian Journal of Zoology – Revue Canadienne de Zoologie*, 66(10), 2233-2242.

- Hartmann, N.B., Legros, S., Von der Kammer, F., Hofmann, T. & Baun, A. 2012. The potential of TiO₂ nanoparticles as carriers for cadmium uptake in *Lumbriculus variegatus* and *Daphnia magna*. *Aquatic Toxicology*, 118, 1-8.
- He, X.B., Cui, Y.D. & Wang, H.Z. 2012. Two new species of Tubificinae (Annelida: Clitellata: Naididae) from Tibet, China. *Zootaxa*, (3458), 159-165.
- Hessling, R., Müller, M.C. & Westheide, W. 1999. CLSM analysis of serotonin-immunoreactive neurons in the central nervous system of *Nais variabilis*, *Slavina appendiculata* and *Stylaria lacustris* (Oligochaeta: Naididae). *Hydrobiologia*, 406, 223-233.
- Higgins, C.P., Paesani, Z.J., Chalew, T.E.A. & Halden, R.U. 2009. Bioaccumulation of triclocarban in *Lumbriculus variegatus*. *Environmental Toxicology and Chemistry*, 28(12), 2580-2586.
- Hirabayashi, K., Oga, K. & Yamamoto, M. 2014. Bathymetric distribution of aquatic Oligochaeta in Lake Kizaki, Central Japan, in Pinder, A., Arslan, N. & Wetzel, M. (Eds.). *Proceedings of the 12th International Symposium on Aquatic Oligochaeta, Zoosymposia*, 36-43.
- Hirsch, M.P. 1998. Bioaccumulation of silver from laboratory-spiked sediments in the oligochaete (*Lumbriculus variegatus*). *Environmental Toxicology and Chemistry*, 17(4), 605-609.
- Hoover, J.J. & Lodes, M.J. 1986. Chaetogaster limnæi (Oligochaeta, Naididae) in Oklahoma. *Southwestern Naturalist*, 31(4), 542-544.
- Huang, X., Liang, P. & Qian, Y. 2007. Excess sludge reduction induced by *Tubifex tubifex* in a recycled sludge reactor. *Journal of Biotechnology*, 127(3), 443-451.
- Ingersoll, C.G., Brunson, E.L., Wang, N., Dwyer, E.J., Ankley, G.T., Mount, D.R., Huckins, J., Petty, J. & Landrum, P.E. 2003. Uptake and depuration of nonionic organic contaminants from sediment by the oligochaete, *Lumbriculus variegatus*. *Environmental Toxicology and Chemistry*, 22(4), 872-885.
- Ishii, R. & Shimizu, T. 1995. Unequal first cleavage in the *Tubifex* egg – involvement of a monastral mitotic apparatus. *Development Growth & Differentiation*, 37(6), 687-701.
- Ishii, R. & Shimizu, T. 1997a. Equalization of unequal first cleavage in the *Tubifex* egg by introduction of an additional centrosome: Implications for the absence of cortical mechanisms for mitotic spindle asymmetry. *Developmental Biology*, 189(1), 49-56.
- Ishii, R. & Shimizu, T. 1997b. Reproductive capability of maternal centrosomes in the *Tubifex* egg. *Zoological Science*, 14(6), 961-968.
- Jantunen, A.P.K., Tuikka, A., Akkanen, J. & Kukkonen, J.V.K. 2008. Bioaccumulation of atrazine and chlorpyrifos to *Lumbriculus variegatus* from lake sediments. *Ecotoxicology and Environmental Safety*, 71(3), 860-868.
- Jordan, M. 1986. *Tubifex* bioassay test in studies on the pyrethroid decis 2.5 EC. *Folia Biologica-Krakow*, 34(4), 393-397.
- Juget, J., Chatelliers, M.C.D. & Rodriguez, P. 2006. *Troglodrilus* (Annelida, Oligochaeta, Tubificidae), a new genus from subterranean habitats in southwestern Europe. *Hydrobiologia*, 564(1), 7-17.
- Kaeser, A.J. & Sharpe, W.E. 2008. The ecology of *Tubifex tubifex* in two *Myxobolus cerebralis* enzootic streams in Pennsylvania. *Journal of Freshwater Ecology*, 23(4), 575-588.
- Kaiser, M., Irmer, U. & Weiler, K. 1989. Monitoring of water-quality – seasonal-variations of heavy-metals in sediment, suspended particulate matter and Tubificids of the Elbe River. *Environmental Technology Letters*, 10(9), 845-854.
- Kajak, Z. & Prus, P. 2004. Influence of the population density and the amount of food on *Chironomus plumosus* (L.) and Tubificidae. Laboratory experiments. *Polish Journal of Ecology*, 52(1), 47-53.
- Kalininkova, T.B., Gainutdinov, T.M., Seraziev, R.K. & Gainutdinov, M.K. 2003. The effect of ambient temperature on intraspecific interactions controlling heat tolerance in the oligochaetes *Enchytraeus albidus* and *Tubifex tubifex*. *Russian Journal of Ecology*, 34(3), 198-201.
- Kaonga, C.C., Kumwenda, J. & Mapoma, H.T. 2010. Accumulation of lead, cadmium, manganese, copper and zinc by sludge worms; *Tubifex tubifex* in sewage sludge. *International Journal of Environmental Science & Technology*, 7(1), 119-126.
- Kathman, R.D. 1985. Synonymy of *Pristinella jenkinsae* (Oligochaeta, Naididae). *Proceedings of the Biological Society of Washington*, 98(4), 1022-1027.
- Kaygorodova, I.A., Sherbakov, D.Y. & Martin, P. 2007. Molecular phylogeny of Baikalian Lumbriculidae (Oligochaeta): evidence for recent explosive speciation. *Comparative Cytogenetics*, 1(1), 71-84.
- Keilty, T.J., White, D.S. & Landrum, P.F. 1988. Sublethal responses to endrin in sediment by *Stylodrilus heringianus* (Lumbriculidae) as measured by a 137 cesium marker layer technique. *Aquatic Toxicology*, 13(3), 251-270.
- Kerans, B.L., Rasmussen, C., Stevens, R., Colwell, A.E.L. & Winton, J.R. 2004. Differential propagation of the metazoan parasite *Myxobolus cerebralis* by *Limnodrilus hoffmeisteri*, *Ilyodrilus templetoni*, and genetically distinct strains of *Tubifex tubifex*. *Journal of Parasitology*, 90(6), 1366-1373.
- Kerans, B.L., Stevens, R.I. & Lemmon, J.C. 2005. Water temperature affects a host-parasite interaction: *Tubifex tubifex* and *Myxobolus cerebralis*. *Journal of Aquatic Animal Health*, 17(3), 216-221.
- Khan, F.R., Paul, K.B., Dybowska, A.D., Valsami-Jones, E., Lead, J.R., Stone, V. & Fernandes, T.F. 2015. Accumulation dynamics and acute toxicity of silver nanoparticles to *Daphnia magna* and *Lumbriculus variegatus*: Implications for metal modeling approaches. *Environmental Science & Technology*, 49(7), 4389-4397.
- Khangarot, B.S. 1991. Toxicity of metals to a fresh-water tubificid worm, *Tubifex tubifex* (Müller). *Bulletin of Environmental Contamination and Toxicology*, 46(6), 906-912.
- Khangarot, B.S. & Rathore, R.S. 2004. Protective action of 24 amino acids on the toxicity of copper to a freshwater tubificid worm *Tubifex tubifex* Müller. *Water Air and Soil Pollution*, 157(1-4), 53-63.
- Khangarot, B.S., Rathore, R.S. & Singh, B.B. 2003. pH-Dependent toxicity of heavy metals to a freshwater sludgeworm *Tubifex tubifex* Müller. *Bulletin of Environmental Contamination and Toxicology*, 71(2), 283-289.
- Kitamura, K. & Shimizu, T. 2000a. Analyses of segment-specific expression of alkaline phosphatase activity in the mesoderm of the oligochaete annelid *Tubifex*: Implications for specification of segmental identity. *Developmental Biology*, 219(2), 214-223.
- Kitamura, K. & Shimizu, T. 2000b. Embryonic expression of alkaline phosphatase activity in the oligochaete annelid *Tubifex*. *Invertebrate Reproduction & Development*, 37(1), 69-73.
- Kitamura, K. & Shimizu, T. 2002. Segment-specific expression of alkaline phosphatase in the *Tubifex* embryo requires DNA replication and mRNA synthesis. *Journal of Experimental Zoology*, 294(1), 68-76.
- Komala, Z. 1992. Toxicity of fastac 10-EC, a pyrethroid insecticide, to *Paramecium primaurelia* and *Tubifex* sp. *Folia Biologica-Krakow*, 40(3-4), 109-112.
- Kraaij, R.H., Tollens, J., Sijm, D., Cornelissen, G., Heikens, A. & Belfroid, A. 2002. Effects of contact time on the sequestration and bioavailability of different classes of hydrophobic organic chemicals to benthic oligochaetes (Tubificidae). *Environmental Toxicology and Chemistry*, 21(4), 752-759.
- Kracun-Kolarevic, M., Kolarevic, S., Atanackovic, A., Markovic, V., Gacic, Z., Paunovic, M. & Vukovic-Gacic, B. 2015. Effects of 5-Fluorouracil, Etoposide and CdCl₂ in aquatic oligochaeta *Limnodrilus udekemianus claparede* (Tubificidae) measured by comet assay. *Water Air and Soil Pollution*, 226(8).

- Kukkonen, J. & Landrum, P.F. 1995. Effects of sediment-bound polydimethylsiloxane on the bioavailability and distribution of benzo a pyrene in lake sediment to *Lumbriculus variegatus*. *Environmental Toxicology and Chemistry*, 14(3), 523-531.
- Kukkonen, J.V.K., Mitra, S., Landrum, P.F., Gossiaux, D.C., Gunnarsson, J. & Weston, D. 2005. The contrasting roles of sedimentary plant-derived carbon and black carbon on sediment-spiked hydrophobic organic contaminant bioavailability to *Diporeia* species and *Lumbriculus variegatus*. *Environmental Toxicology and Chemistry*, 24(4), 877-885.
- Kyselkova, I. & Marsalek, B. 2000. Using of *Daphnia pulex*, *Artemia salina* and *Tubifex tubifex* for cyanobacterial microcystins toxicity detection. *Biologia*, 55(6), 637-643.
- Lafont, M. 1987. Production of *Tubificidae* in the littoral-zone of Lake Leman near thonon-les-bains – a methodological approach. *Hydrobiologia*, 155, 179-187.
- Lafont, M. & Juget, J. 1993. *Rhyacodrilus-ardierae*, sp. nov., a new species of *Tubificidae* (Oligochaeta) from the River Ardieres (France, tributary of the River Saone). *Bulletin de la Societe Zoologique de France – Evolution et Zoologie*, 118(2), 115-123.
- Lagauzere, S., Boyer, P., Stora, G. & Bonzom, J.M. 2009a. Effects of uranium-contaminated sediments on the bioturbation activity of *Chironomus riparius* larvae (Insecta, Diptera) and *Tubifex tubifex* worms (Annelida, Tubificidae). *Chemosphere*, 76(3), 324-334.
- Lagauzere, S., Terrail, R. & Bonzom, J.M. 2009b. Ecotoxicity of uranium to *Tubifex tubifex* worms (Annelida, Clitellata, Tubificidae) exposed to contaminated sediment. *Ecotoxicology and Environmental Safety*, 72(2), 527-537.
- Landrum, P.E., Leppanen, M., Robinson, S.D., Gossiaux, D.C., Burton, G.A., Greenberg, M., Kukkonen, J.V.K., Eadie, B.J. & Lansing, M.B. 2004a. Comparing behavioral and chronic endpoints to evaluate the response of *Lumbriculus variegatus* to 3,4,3',4'-tetrachlorobiphenyl sediment exposures. *Environmental Toxicology and Chemistry*, 23(1), 187-194.
- Landrum, P.F., Gedeon, M.L., Burton, G.A., Greenberg, M.S. & Rowland, C.D. 2002. Biological responses of *Lumbriculus variegatus* exposed to fluoranthene-spiked sediment. *Archives of Environmental Contamination and Toxicology*, 42(3), 292-302.
- Landrum, P.F., Leppanen, M., Robinson, S.D., Gossiaux, D.C., Burton, G.A., Greenberg, M., Kukkonen, J.V.K., Eadie, B.J. & Lansing, M.B. 2004b. Effect of 3,4,3',4'-tetrachlorobiphenyl on the reworking behavior of *Lumbriculus variegatus* exposed to contaminated sediment. *Environmental Toxicology and Chemistry*, 23(1), 178-186.
- Lang, C. 1989. Effects of small-scale sedimentary patchiness on the distribution of tubificid and Lumbriculid worms in Lake Geneva. *Freshwater Biology*, 21(3), 477-481.
- Lang, C. 2000. Response of oligochaete (Tubificidae and Lumbriculidae) and diptera Chironomidae communities to the decrease of phosphorus concentrations in Lake Geneva (Little Lake). *Annales de Limnologie – International Journal of Limnology*, 36(1), 13-20.
- Lanzavecchia, G., Deeguileor, M., Valvassori, R. & Lanzavecchia, P. 1987. Analysis and reconstruction of unusual obliquely striated fibers in Lumbriculids (Annelida, Oligochaeta). *Journal of Muscle Research and Cell Motility*, 8(3), 209-219.
- Lasier, P.J., Washington, J.W., Hassan, S.M. & Jenkins, T.M. 2011. Perfluorinated chemicals in surface waters and sediments from northwest Georgia, USA, and their bioaccumulation in *Lumbriculus variegatus*. *Environmental Toxicology and Chemistry*, 30(10), 2194-2201.
- Leonard, E.M. & Wood, C.M. 2013. Acute toxicity, critical body residues, Michaelis-Menten analysis of bioaccumulation, and ionoregulatory disturbance in response to waterborne nickel in four invertebrates: *Chironomus riparius*, *Lymnaea stagnalis*, *Lumbriculus variegatus* and *Daphnia pulex*. *Comparative Biochemistry and Physiology – Part C: Toxicology and Pharmacology*, 158(1), 10-21.
- Leppanen, M.T. & Kukkonen, J.V.K. 1998. Relationship between reproduction, sediment type, and feeding activity of *Lumbriculus variegatus* (Müller): Implications for sediment toxicity testing. *Environmental Toxicology and Chemistry*, 17(11), 2196-2202.
- Leppanen, M.T. & Kukkonen, J.V.K. 2000. Fate of sediment-associated pyrene and benzo a pyrene in the freshwater oligochaete *Lumbriculus variegatus* (Müller). *Aquatic Toxicology*, 49(3), 199-212.
- Lestochova, E.I. 1994. Influence of small river conditions on the abundance of *Tubificidae*. *Hydrobiologia*, 278(1-3), 129-131.
- Leynen, M., Van den Berckt, T., Aerts, J.M., Castlein, B., Berckmans, D. & Ollevier, F. 1999. The use of *Tubificidae* in a biological early warning system. *Environmental Pollution*, 105(1), 151-154.
- Liu, T.T., Diao, J.L., Di, S.S. & Zhou, Z.Q. 2014. Stereoselective bioaccumulation and metabolite formation of triadimefon in *Tubifex tubifex*. *Environmental Science & Technology*, 48(12), 6687-6693.
- Liu, T.T., Diao, J.L., Di, S.S. & Zhou, Z.Q. 2015. Bioaccumulation of isocarbophos enantiomers from laboratory-contaminated aquatic environment by *Tubificid* worms. *Chemosphere*, 124, 77-82.
- Lobo, H. & Espindola, E.L.G. 2014. *Branchiura sowerbyi* Beddard, 1892 (Oligochaeta: Naididae) as a test species in ecotoxicology bioassays: a review, in Pinder, A., Arslan, N. & Wetzel, M. (Eds.). *Proceedings of the 12th International Symposium on Aquatic Oligochaeta, Zoosymposia*, 59-69.
- Lodh, N., Rizzo, D.M., Kerans, B.L., McGinnis, S., Fytalis, N. & Stevens, L. 2015. If you've seen one worm, have you seen them all? Spatial, community, and genetic variability of tubificid communities in Montana. *Freshwater Science*, 34(3), 909-917.
- Lohlein, B. 1996. Seasonal dynamics of aufwuchs Naididae (Oligochaeta) on *Phragmites australis* in a eutrophic lake. *Hydrobiologia*, 334(1-3), 115-123.
- Lopez, L.C.S., Rodrigues, P. & Rios, R.I. 1999. Frogs and snakes as phoretic dispersal agents of bromeliad ostracods (Limnocytheridae: Elpidium) and annelids (Naididae: Dero). *Biotropica*, 31(4), 705-708.
- Lou, J.Q., Sun, P.D., Guo, M.X., Wu, G. & Song, Y.Q. 2011. Simultaneous sludge reduction and nutrient removal (SSRNR) with interaction between *Tubificidae* and microorganisms: A full-scale study. *Bioresource Technology*, 102(24), 11132-11136.
- Luszczek-Trojnar, E., Sroka, K., Klaczak, A., Nowak, M. & Popek, W. 2014. Bioaccumulation and purification of cadmium in *Tubifex tubifex*. *Turkish Journal of Fisheries and Aquatic Sciences*, 14(4), 939-946.
- Lytykainen, M., Pehkonen, S., Akkanen, J., Leppanen, M. & Kukkonen, J.V.K. 2007. Bioaccumulation and biotransformation of polycyclic aromatic hydrocarbons during sediment tests with oligochaetes (*Lumbriculus variegatus*). *Environmental Toxicology and Chemistry*, 26(12), 2660-2666.
- Madill, J., Coates, K.A., Wetzel, M.J. & Gelder, S.R. 1992. Common and scientific names of Aphanoneuran and clitellate annelids of the United States of America and Canada. *Soil Biology & Biochemistry*, 24(12), 1259-1262.
- Maenpaa, K., Leppanen, M.T. & Kukkonen, J.V.K. 2009. Sublethal toxicity and biotransformation of pyrene in *Lumbriculus variegatus* (Oligochaeta). *Science of the Total Environment*, 407(8), 2666-2672.
- Maestre, Z., Martinez-Madrid, M. & Rodriguez, P. 2009. Monitoring the sensitivity of the oligochaete *Tubifex tubifex* in laboratory cultures using three toxicants. *Ecotoxicology and Environmental Safety*, 72(8), 2083-2089.
- Marchese, M.R. & Brinkhurst, R.O. 1996. A comparison of two tubificid oligochaete species as candidates for sublethal bioassay tests relevant to subtropical and tropical regions. *Hydrobiologia*, 334(1-3), 163-168.
- Marian, M.P. & Pandian, T.J. 1985. Interference of chironomus in an open culture system for *Tubifex tubifex*. *Aquaculture*, 44(3), 249-251.

- Marotta, R., Crottini, A., Prada, V. & Ferraguti, M. 2009. A morphological reappraisal of *Tubifex blanchardi* Vejdovsky, 1891 (Clitellata: Tubificidae). *Acta Zoologica*, 90(2), 179-188.
- Marotta, R., Crottini, A., Raimondi, E., Fondello, C. & Ferraguti, M. 2014. A like but different: the evolution of the *Tubifex tubifex* species complex (Annelida, Clitellata) through polyploidization. *BMC Evolutionary Biology*, 14.
- Martin, P., Ferraguti, M. & Kaygorodova, I. 1998. Description of two new species of *Rhynchelmis* (Oligochaeta: Lumbriculidae) from Lake Baikal (Russia), using classical morphology and ultrastructure of spermatozoa. *Annales de Limnologie – International Journal of Limnology*, 34(3), 283-293.
- Martin, P. & Giani, N. 1995. 2 New species of *eptiordilus* (Oligochaeta, Tubificidae) from Lake Nyasa and Tanganyika (East-Africa), with redescriptions of *E. slovenicus* and *E. michaelensi*. *Zoologica Scripta*, 24(1), 13-19.
- Martin, P. & Kaygorodova, I. 2008. A new species of *Pseudorhynchelmis* Hrabe, 1982 (Clitellata: Lumbriculidae) from Lake Baikal, with re-descriptions of *P. parva* and *P. olchonensis*. *Zootaxa*, (1938), 23-39.
- Martin, P., Martinez-Ansemil, E. & Sambugar, B. 2010. The Baikalian genus *Rhyacodriloides* in Europe: phylogenetic assessment of *Rhyacodriloidinae* subfam. nov., within the Naididae (Annelida). *Zoologica Scripta*, 39(5), 462-482.
- Martinez, V.G., Mengler, G.J. & Zoran, M.J. 2005. Regeneration and asexual reproduction share common molecular changes: upregulation of a neural glycoepitope during morphallaxis in *Lumbriculus*. *Mechanisms of Development*, 122(5), 721-732.
- Martinez, V.G., Reddy, P.K. & Zoran, M.J. 2006. Asexual reproduction and segmental regeneration, but not morphallaxis, are inhibited by boric acid in *Lumbriculus variegatus* (Annelida: Clitellata: Lumbriculidae). *Hydrobiologia*, 564(1), 73-86.
- Martinez-Madrid, M., Rodriguez, P., Perez-Iglesias, J.I. & Navarro, E. 1999. Sediment toxicity bioassays for assessment of contaminated sites in the Nervion River (Northern Spain). 2. *Tubifex tubifex* reproduction sediment bioassay. *Ecotoxicology*, 8(2), 111-124.
- Martinsson, S., Achurra, A., Svensson, M. & Erseus, C. 2013. Integrative taxonomy of the freshwater worm *Rhyacodrilus falciformis* s.l. (Clitellata: Naididae), with the description of a new species. *Zoologica Scripta*, 42(6), 612-622.
- Matisoff, G., Wang, X.S. & McCall, P.L. 1999. Biological redistribution of lake sediments by tubificid oligochaetes: Branchiura sowerbyi and *Limnodrilus hoffmeisteri/Tubifex tubifex*. *Journal of Great Lakes Research*, 25(1), 205-219.
- Matsuo, K., Yoshida, H. & Shimizu, T. 2005. Differential expression of caudal and dorsal genes in the teloblast lineages of the oligochaete annelid *Tubifex tubifex*. *Development Genes and Evolution*, 215(5), 238-247.
- McKeyfender, D. & Fender, W.M. 1988. *Phagodrilus* gen. nov. (Lumbriculidae) – systematics and biology of a predaceous oligochaete from western North America. *Canadian Journal of Zoology – Revue Canadienne de Zoologie*, 66(10), 2304-2311.
- Meller, M., Egeler, P., Rombke, J., Schallnass, H., Nagel, R. & Streit, B. 1998. Short-term toxicity of lindane, hexachlorobenzene, and copper sulfate to tubificid sludge worms (Oligochaeta) in artificial media. *Ecotoxicology and Environmental Safety*, 39(1), 10-20.
- Mermilliod-Blondin, F., des Chatelliers, M.C. & Gerino, M. 2003. Effects of the interaction between Tubificid worms on the functioning of hyporheic sediments: an experimental study in sediment columns. *Archiv Fur Hydrobiologie*, 156(2), 203-223.
- Mermilliod-Blondin, F., Foulquier, A., Gilbert, F., Navel, S., Montuelle, B., Bellvert, F., Comte, G., Grossi, V., Fourel, F., Lecuyer, C. & Simon, L. 2013. Benzo(a)pyrene inhibits the role of the bioturbator *Tubifex tubifex* in river sediment biogeochemistry. *Science of the Total Environment*, 450, 230-241.
- Mermilliod-Blondin, F., Nogaro, G., Datry, T., Malard, F. & Gibert, J. 2005. Do Tubificid worms influence the fate of organic matter and pollutants in stormwater sediments? *Environmental Pollution*, 134(1), 57-69.
- Mermilliod-Blondin, F., Nogaro, G., Vallier, F. & Gibert, J. 2008. Laboratory study highlights the key influences of stormwater sediment thickness and bioturbation by Tubificid worms on dynamics of nutrients and pollutants in stormwater retention systems. *Chemosphere*, 72(2), 213-223.
- Milbrink, G. 1987. Biological characterization of sediments by standardized tubificid bioassays. *Hydrobiologia*, 155, 267-275.
- Mino, L.A., Folco, S., de D'Angelo, A.M.P. & Guerrero, N.R.V. 2006. Modeling lead bioavailability and bioaccumulation by *Lumbriculus variegatus* using artificial particles. Potential use in chemical remediation processes. *Chemosphere*, 63(2), 261-268.
- Misra, R.B., Babu, G.S., Ray, R.S. & Hans, R.K. 2002. *Tubifex*: A sensitive model for UV-B-induced phototoxicity. *Ecotoxicology and Environmental Safety*, 52(3), 288-295.
- Molnar, L., Kiszler, G. & Pollak, E. 2006a. Identification and pattern of primary sensory cells in the body wall epithelium of the tubificid worm, *Limnodrilus hoffmeisteri*. *Hydrobiologia*, 564(1), 45-50.
- Molnar, L., Kiszler, G., Pollak, E. & Deres, L. 2006b. Distribution pattern of gamma-amino butyric acid immunoreactive neural structures in the central and peripheral nervous system of the tubificid worm, *Limnodrilus hoffmeisteri*. *Hydrobiologia*, 564(1), 33-43.
- Morev, A.P. 1992. On the taxonomical status of *Alexandrovia ringulata* and *A. onegensis* (Oligochaeta, Tubificidae). *Zoologichesky Zhurnal*, 71(5), 128-133.
- Mosleh, Y.Y., Mofeed, J., Afifi, M. & Almaghrabi, O.A. 2014. Biological effects of pyrimethinal on aquatic worms (*Tubifex tubifex*) under laboratory conditions. *Bulletin of Environmental Contamination and Toxicology*, 92(1), 85-89.
- Mosleh, Y.Y., Paris-Palacios, S., Ahmed, M.T., Mahmoud, F.M., Osman, M.A. & Biagianti-Risbourg, S. 2007. Effects of chitosan on oxidative stress and metallothioneins in aquatic worm *Tubifex tubifex* (Oligochaeta, Tubificidae). *Chemosphere*, 67(1), 167-175.
- Mosleh, Y.Y., Paris-Palacios, S., Couderchet, M., Biagianti-Risbourg, S. & Vernet, G. 2005. Effects of the herbicide isoproturon on metallothioneins, growth, and antioxidative Defenses in the aquatic worm *Tubifex tubifex* (Oligochaeta, Tubificidae). *Ecotoxicology*, 14(5), 559-571.
- Nakamoto, A., Arai, A. & Shimizu, T. 2000. Cell lineage analysis of pattern formation in the *Tubifex* embryo. II. Segmentation in the ectoderm. *International Journal of Developmental Biology*, 44(7), 797-805.
- Nakamoto, A., Arai, A. & Shimizu, T. 2004. Specification of polarity of teloblastogenesis in the oligochaete annelid *Tubifex*: cellular basis for bilateral symmetry in the ectoderm. *Developmental Biology*, 272(1), 248-261.
- Nakamoto, A., Arai, A. & Shimizu, T. 2006. Generation of bilateral symmetry in the ectoderm of the *Tubifex* embryo: Involvement of cell-cell interactions. *Hydrobiologia*, 564(1), 19-32.
- Narita, T. 2001. Cocoon deposition of *Rhyacodrilus hiemalis* Ohtaka (Tubificidae) in Lake Biwa, Japan. *Hydrobiologia*, 463, 141-148.
- Narita, T. 2006. Seasonal vertical migration and aestivation of *Rhyacodrilus hiemalis* (Tubificidae, Clitellata) in the sediment of Lake Biwa, Japan. *Hydrobiologia*, 564(1), 87-93.
- Naveed, M.I. 2012. Preliminary studies on aquatic Oligochaeta in and around Chennai, Tamil Nadu, India. *Turkish Journal of Zoology*, 36(1), 25-37.
- Naveed, M.I., Thulasiraja, S., Karunakaran, S., Kulandaivel, C. & Selvan, R.T. 2014. Preliminary survey of aquatic oligochaetes in Eastern Tamil Nadu,

- India, in Pinder, A., Arslan, N. & Wetzel, M. (Eds.). *Proceedings of the 12th International Symposium on Aquatic Oligochaeta, Zoosymposia*, 14-23.
- Nemec, A.F.L. & Brinkhurst, R.O. 1987. A comparison of methodological approaches to the subfamilial classification of the Naididae (Oligochaeta). *Canadian Journal of Zoology*, 65(3), 691-707.
- Nybom, I., Waissi-Leinonen, G., Maenpaa, K., Leppanen, M.T., Kukkonen, J.V.K., Werner, D. & Akkanen, J. 2015. Effects of activated carbon ageing in three PCB contaminated sediments: Sorption efficiency and secondary effects on *Lumbriculus variegatus*. *Water Research*, 85, 413-421.
- Nybom, I., Werner, D., Leppanen, M.T., Siavalas, G., Christanis, K., Karapanagioti, H.K., Kukkonen, J.V.K. & Akkanen, J. 2012. Responses of *Lumbriculus variegatus* to Activated Carbon Amendments in Uncontaminated Sediments. *Environmental Science & Technology*, 46(23), 12895-12903.
- O'Gara, B.A., Murray, P.M., Hoyt, E.M., Leigh-Logan, T. & Smeaton, M.B. 2006. The Vitamin E analog Trolox reduces copper toxicity in the annelid *Lumbriculus variegatus* but is also toxic on its own. *Neurotoxicology*, 27(4), 604-614.
- O'Rourke, S., Stone, V., Stolpe, B. & Fernandes, T. 2015. Assessing the acute hazards of zinc oxide nanomaterials to *Lumbriculus variegatus*. *Ecotoxicology*, 24(6), 1372-1384.
- Ohtaka, A. 1995. A new species of the genus *Rhyacodrilus* Bretscher (Oligochaeta, Tubificidae) from Japanese lakes. *Zoological Science*, 12(4), 491-498.
- Ohtaka, A. 2014. Profundal oligochaete faunas (Annelida, Clitellata) in Japanese lakes, in Pinder, A., Arslan, N. & Wetzel, M. (Eds.). *Proceedings of the 12th International Symposium on Aquatic Oligochaeta, Zoosymposia*, 24-35.
- Omodeo, P. & Coates, K.A. 2001. New alluroidids (Annelida, Clitellata) from Guyana. *Hydrobiologia*, 463, 39-47.
- Oppinger, R.W., Bartley, M. & Wagner, E.J. 2011. Culture of *Tubifex tubifex*: Effect of feed type, ration, temperature, and density on juvenile recruitment, production, and adult survival. *North American Journal of Aquaculture*, 73(1), 68-75.
- Otubu, J.E., Hunter, J.V., Francisco, K.L. & Uchrin, C.G. 2006. Temperature effects on Tubificid worms and their relation to sediment oxygen demand. *Journal of Environmental Science and Health – Part A: Toxic/Hazardous Substances & Environmental Engineering*, 41(8), 1607-1613.
- Ozdemir, A., Duran, M., Akyildiz, G.K. & Sen, A. 2011. EROD and metallothionein in *Limnodrilus profundicola* (Oligochaeta: Tubificidae) as an indicator of pollution exposure in the Curuksu stream of Menderes River, Denizli, Turkey. *Desalination and Water Treatment*, 26(1-3), 98-103.
- Pakarinen, K., Petersen, E.J., Leppanen, M.T., Akkanen, J. & Kukkonen, J.V.K. 2011. Adverse effects of fullerenes (nC(60)) spiked to sediments on *Lumbriculus variegatus* (Oligochaeta). *Environmental Pollution*, 159(12), 3750-3756.
- Paoletti, A. 1989. Cohort cultures of *Tubifex tubifex* forms. *Hydrobiologia*, 180, 143-150.
- Paoletti, A.D. & Sambugar, B. 1996. Aquatic Oligochaeta in Italy, with special reference to Naididae. *Hydrobiologia*, 334(1-3), 37-49.
- Paris-Palacios, S., Mosleh, Y.Y., Almohamad, M., Delahaut, L., Conrad, A., Arnoult, F. & Biagioli-Risbourg, S. 2010. Toxic effects and bioaccumulation of the herbicide isoproturon in *Tubifex tubifex* (Oligochaeta, Tubificidae): A study of significance of autotomy and its utility as a biomarker. *Aquatic Toxicology*, 98(1), 8-14.
- Pasteris, A., Bonacina, C. & Bonomi, G. 1994. Observations on cohorts of *Tubifex tubifex* cultured at different food levels, using cellulose substrate. *Hydrobiologia*, 278(1-3), 315-320.
- Pasteris, A., Bonomi, G. & Bonacina, C. 1996. Age, stage and size structure as population state variables for *Tubifex tubifex* (Oligochaeta, Tubificidae). *Hydrobiologia*, 334(1-3), 125-132.
- Peng, Y., Wang, H.Z. & Cui, Y.D. 2014. Two species of Naididae (Annelida, Clitellata) from southern Tibet, China. *ZooKeys*, (444), 59-68.
- Penttinen, O.P. & Kukkonen, J.V.K. 2000. Metabolic response of *Lumbriculus variegatus* to respiratory uncoupler in cold and anoxic water. *Environmental Toxicology and Chemistry*, 19(8), 2073-2075.
- Petersen, S., Arlt, G., Faubel, A. & Carman, K.R. 1998. On the nutritive significance of dissolved free amino acid uptake for the cosmopolitan oligochaete *Nais elinguis* Müller (Naididae). *Estuarine, Coastal and Shelf Science*, 46(1), 85-91.
- Pinder, A. 2001. Notes on the diversity and distribution of Australian Naididae and Phreodrilidae (Oligochaeta: Annelida). *Hydrobiologia*, 463, 49-64.
- Pinder, A.M., Eberhard, S.M. & Humphreys, W.F. 2006. New phallodrilines (Annelida: Clitellata: Tubificidae) from Western Australian groundwater. *Zootaxa*, (1304), 31-48.
- Postolache, C., Risnoveanu, G. & Vadineanu, A. 2006. Nitrogen and phosphorous excretion rates by Tubificids from the Prahova River (Romania). *Hydrobiologia*, 553, 121-127.
- Puigagut, J., Chazarenc, F. & Comeau, Y. 2014. Influence of Tubificid worms on nutrient fluxes across water-sediment interface in fish farm settling ponds. *Knowledge and Management of Aquatic Ecosystems*, (413).
- Purschke, G. 2003. Ultrastructure of phaemosome photoreceptors in *Stylaria lacustris* (Naididae, 'Oligochaeta', Clitellata) and their importance for the position of the Clitellata in the phylogenetic system of the Annelida. *Journal of Zoological Systematics and Evolutionary Research*, 41(2), 100-108.
- Rathore, R.S. & Khangarot, B.S. 2002. Effects of temperature on the sensitivity of sludge worm *Tubifex tubifex* Müller to selected heavy metals. *Ecotoxicology and Environmental Safety*, 53(1), 27-36.
- Rathore, R.S. & Khangarot, B.S. 2003. Effects of water hardness and metal concentration on a freshwater *Tubifex tubifex* Müller. *Water Air and Soil Pollution*, 142(1-4), 341-356.
- Reynoldson, T.B., Rodriguez, P. & Madrid, M.M. 1996. A comparison of reproduction, growth and acute toxicity in two populations of *Tubifex tubifex* (Müller, 1774) from the North American Great Lakes and Northern Spain. *Hydrobiologia*, 334(1-3), 199-206.
- Robbins, J.A., Keilty, T., White, D.S. & Edgington, D.N. 1989. Relationships among tubificid abundances, sediment composition, and accumulation rates in Lake Erie. *Canadian Journal of Fisheries and Aquatic Sciences*, 46(2), 223-231.
- Rodriguez, P., Fend, S.V. & Lenat, D.R. 2014. *Sylphella pucoo* gen. nov., sp. nov., and two additional new species of aquatic oligochaetes (*Lumbriculidae*, Clitellata) from poorly-known lotic habitats in North Carolina (USA). *ZooKeys*, (451), 1-32.
- Rodriguez, P., Martinez-Madrid, M., Arrate, J.A. & Navarro, E. 2001. Selective feeding by the aquatic oligochaete *Tubifex tubifex* (Tubificidae, Clitellata). *Hydrobiologia*, 463, 133-140.
- Rogge, R.W. & Drewes, C.D. 1993. Assessing sublethal neurotoxicity effects in the fresh-water oligochaete, *Lumbriculus variegatus*. *Aquatic Toxicology*, 26(1-2), 73-89.
- Romano Spica, V., Giampaoli, S., Buggiotti, L., Vitali, M., Gianfranceschi, G. & Soldati, R. 2014. Environment health and intraspecific biodiversity in *T. tubifex*: a preliminary analysis of a population from Apennines springs. *International Journal of Environmental Science & Technology*, 11(2), 461-468.
- Ruidisch, S., Elmatbouli, M. & Hoffmann, R.W. 1991. The role of *Tubifex* worms as an intermediate host in the life-cycle of *Myxobolus pavlovskii* (Akhmerov, 1954). *Parasitology Research*, 77(8), 663-667.
- Sardo, A.M., Pereira, L., Gerhardt, A. & Soares, A. 2011. Effect of the exposure to metal lead on the regenerative ability of *Lumbriculus variegatus* (Oligochaeta). *Environmental Toxicology and Pharmacology*, 31(1), 205-211.

- Sardo, A.M. & Soares, A. 2010. Assessment of the effects of the pesticide imidacloprid on the behaviour of the aquatic oligochaete *Lumbriculus variegatus*. *Archives of Environmental Contamination and Toxicology*, 58(3), 648-656.
- Sato, C. & Ohtaka, A. 2012. Redescription of *Piguetiella denticulata* (Annelida, Clitellata, Naididae) from Japan and the Russian Far East, with a description of the genital organs in the species. *Limnology*, 13(1), 143-148.
- Sauter, G. & Gude, H. 1996. Influence of grain size on the distribution of tubificid oligochaete species. *Hydrobiologia*, 334(1-3), 97-101.
- Schenkova, J. & Cermak, V. 2013. Description of *Pristina armata* sp. nov. (Clitellata: Naididae: Pristininae) from a carnivorous plant (*Nepenthes* sp.) in Borneo, Indonesia. *Zootaxa*, 3686(5), 587-592.
- Schubauerberigan, M.K., Dierkes, J.R., Monson, P.D. & Ankley, G.T. 1993. PH-dependent toxicity of CD, CU, NI, PB and ZN to *Ceriodaphnia dubia*, *Pimephales promelas*, *Hyalaea azteca* and *Lumbriculus variegatus*. *Environmental Toxicology and Chemistry*, 12(7), 1261-1266.
- Schubauerberigan, M.K., Monson, P.D., West, C.W. & Ankley, G.T. 1995. Influence of pH on the toxicity of ammonia to *Chironomus tentans* and *Lumbriculus variegatus*. *Environmental Toxicology and Chemistry*, 14(4), 713-717.
- Schuler, L.J., Wheeler, M., Bailer, A.J. & Lydy, M.J. 2003. Toxicokinetics of sediment-sorbed benzo a pyrene and hexachlorobiphenyl using the freshwater invertebrates *Hyalaea azteca*, *Chironomus tentans*, and *Lumbriculus variegatus*. *Environmental Toxicology and Chemistry*, 22(2), 439-449.
- Shimizu, T. 1986. Bipolar segregation of mitochondria, actin network, and surface in the *Tubifex* egg – role of cortical polarity. *Developmental Biology*, 116(1), 241-251.
- Shimizu, T. 1988. Localization of actin networks during early development of *Tubifex* embryos. *Developmental Biology*, 125(2), 321-331.
- Shimizu, T. 1989. Asymmetric segregation and polarized redistribution of pole plasm during early cleavages in the *Tubifex* embryo – role of actin networks and mitotic apparatus. *Development Growth & Differentiation*, 31(3), 283-297.
- Shimizu, T. 1990. Polar body formation in *Tubifex* eggs. *Annals of the New York Academy of Sciences*, 582, 260-272.
- Shimizu, T. 1994. The prevention of smaller blastomeres of early *Tubifex* embryos from entering mitosis by unreplicated DNA. *Developmental Biology*, 161(1), 274-284.
- Shimizu, T. 1995a. Lineage-specific alteration in cell-cycle structure in early *Tubifex* embryos. *Development Growth & Differentiation*, 37(3), 263-272.
- Shimizu, T. 1995b. Role of the cytoskeleton in the generation of spatial patterns in *Tubifex* eggs, in Capco, D.G. (Ed.). *Cytoskeletal Mechanisms During Animal Development*. Current Topics in *Developmental Biology*, 197-235.
- Shimizu, T. 1996a. Behaviour of centrosomes in early *Tubifex* embryos: Asymmetric segregation and mitotic cycle-dependent duplication. *Roux's Archives of Developmental Biology*, 205(5-6), 290-299.
- Shimizu, T. 1996b. Ooplasmic redistribution in *Tubifex* eggs with selectively impaired cortical actin cytoskeleton. *Developmental Biology*, 180(1), 54-62.
- Shimizu, T. 1996c. The first two cleavages in *Tubifex* involve distinct mechanisms to generate asymmetry in mitotic apparatus. *Hydrobiologia*, 334(1-3), 269-276.
- Shimizu, T. 1997a. Reorganization of the cortical actin cytoskeleton during maturation division in the *Tubifex* egg: Possible involvement of protein kinase C. *Developmental Biology*, 188(1), 110-121.
- Shimizu, T. 1997b. Separation of microvilli from *Tubifex* eggs upon activation: Its inhibition by concanavalin A hinders ooplasmic segregation and cleavage. *Development Growth & Differentiation*, 39(6), 787-794.
- Shimizu, T., Ishii, R. & Takahashi, H. 1998. Unequal cleavage in the early *Tubifex* embryo. *Development Growth & Differentiation*, 40(3), 257-266.
- Shimizu, T., Kitamura, K., Arai, A. & Nakamoto, A. 2001. Pattern formation in embryos of the oligochaete annelid *Tubifex*: cellular basis for segmentation and specification of segmental identity. *Hydrobiologia*, 463, 123-131.
- Sibley, T.H. & Stohr, J.S. 1990. Accumulation of AM-241 and CM-244 from water and sediments by *hyalella* sp., and *Tubifex* sp. *Bulletin of Environmental Contamination and Toxicology*, 44(1), 165-172.
- Singh, B.B., Chandra, R. & Sharma, Y.K. 2008. Effect of pyridine and formaldehyde on a macrophyte (*Lemna minor* L.) and a sludge worm (*Tubifex tubifex* Müller) in fresh water microcosms. *Applied Ecology and Environmental Research*, 6(2), 21-35.
- Singh, R.K., Chavan, S.L. & Sapkale, P.H. 2007. Heavy metal concentrations in water, sediments and body tissues of red worm (*Tubifex* sp.) collected from natural habitats in Mumbai, India. *Environmental Monitoring and Assessment*, 129(1-3), 471-481.
- Sjolin, E., Erseus, C. & Kallersjo, M. 2005. Phylogeny of Tubificidae (Annelida, Clitellata) based on mitochondrial and nuclear sequence data. *Molecular Phylogenetics and Evolution*, 35(2), 431-441.
- Slootweg, T., Alvinerie, M., Egeler, P., Gilberg, D., Kukkonen, J.V.K., Oehlmann, J., Prasse, C., Sormunen, A.J. & Liebig, M. 2010. Bioaccumulation of ivermectin from natural and artificial sediments in the benthic organism *Lumbriculus variegatus*. *Journal of Soils and Sediments*, 10(8), 1611-1622.
- Smith, M.E. 1985a. Naididae (Oligochaeta) as hosts for mermithid nematodes (Enoploida, Mermithidae). *Canadian Journal of Zoology – Revue Canadienne de Zoologie*, 63(6), 1459-1462.
- Smith, M.E. 1985b. Setal morphology and its intraspecific variation in *Dero-digitata* and *Dero-nivea* (Oligochaeta, Naididae). *Transactions of the American Microscopical Society*, 104(1), 45-51.
- Smith, M.E. 1985c. Tubificid worms – important organisms in aquatic ecosystems. *American Biology Teacher*, 47(7), 412-415.
- Smutna, M., Hilscherova, K., Paskova, V. & Marsalek, B. 2008. Biochemical parameters in *Tubifex* tubifex as an integral part of complex sediment toxicity assessment. *Journal of Soils and Sediments*, 8(3), 154-164.
- Snimschikova, L.N. 1989. New species and subspecies of the genus *Baikalodrilus* (Oligochaeta, Tubificidae). *Zoologichesky Zhurnal*, 68(9), 23-34.
- Snimschikova, L.N. 1991. A revision of *Lycodrilus phreodrioloides* (Oligochaeta, Tubificidae). *Zoologica Scripta*, 20(3), 221-223.
- Snimschikova, L.N. & Akinschina, T.V. 1993. New species of the genus *Tasserkidrilus* (Oligochaeta, Tubificidae) from Lake Baikal. *Zoologichesky Zhurnal*, 72(10), 5-13.
- Snimschikova, L.N. & Akinschina, T.V. 1994a. *Isochaetides-resorptus* sp. nov. (Oligochaeta, Tubificidae). *Zoologichesky Zhurnal*, 73(9), 158-161.
- Snimschikova, L.N. & Akinschina, T.W. 1994b. *Rhyacodrilus-stephensonii* (Oligochaeta, Tubificidae) from Lake Baikal. *Zoologichesky Zhurnal*, 73(6), 22-27.
- Snimshchikova, L.N. 1989. *Baikalodrilus multicrystallifer* sp. nov. (Oligochaeta, Tubificidae). *Zoologichesky Zhurnal*, 68(2), 300-303.
- Snimshchikova, L.N. 1998. Revision of the genus *Lycodrilus* and description of the evolution of *Limnodrilus* (Oligochaeta, Tubificidae) species in Lake Baikal. I. Revision of *Lycodrilus dybowskii* Grube species and *Tubipenifer semernoyi* genus. *Zoologichesky Zhurnal*, 77(5), 540-547.
- Snimstschikova, L.N. 1985. The taxonomy of the genus *svetlovia* (Oligochaeta, Tubificidae) from the Baikal Lake. *Zoologichesky Zhurnal*, 64(8), 1148-1153.
- Song, Z.H. 2007. Effects of Pentachlorophenol on *Galba pertia*, *Tubifex sinicus* and *Chironomus plumosus* Larvae. *Bulletin of Environmental Contamination and Toxicology*, 79(3), 278-282.

- Soni, A.K. & Joshi, P.C. 1997. High sensitivity of *Tubifex* for ultraviolet-B. *Biochemical and Biophysical Research Communications*, 231(3), 818-819.
- Soors, J., van Haaren, T., Timm, T. & Speybroeck, J. 2013. Bratislavia dadayi (Michaelsen, 1905) (Annelida: Clitellata: Naididae): a new non-indigenous species for Europe, and other non-native annelids in the Schelde estuary. *Aquatic Invasions*, 8(1), 37-44.
- Spencer, D.R. & Wissman, R.E. 1993. Some new records of Naididae and Tubificidae (Annelida, Oligochaeta) from Washington. *Great Basin Naturalist*, 53(4), 395-401.
- Sporka, F. 1996. First record of *Aulodrilus piguetti* (Oligochaeta, Tubificidae) from the River Morava in Slovakia. *Biologia*, 51(5), 494-494.
- Stacey, D.F. & Coates, K.A. 1996. Oligochaetes (Naididae, Tubificidae, Opistocystidae, Enchytraeidae, Sparganophilidae and Alluroididae) of Guyana. *Hydrobiologia*, 334(1-3), 17-29.
- Stacey, D.F. & Hubley, B.R. 1994. New records of a sludge worm *Teneridrilus flexus* erseus and *hiltunen* (Oligochaeta, Tubificidae) from Lake Huron. *Canadian Field-Naturalist*, 108(2), 182-185.
- Steen Redeker, E., van Campenhout, K., Bervoets, L., Reijnders, H. & Blust, R. 2007. Subcellular distribution of Cd in the aquatic oligochaete *Tubifex tubifex*, implications for trophic availability and toxicity. *Environmental Pollution*, 148(1), 166-175.
- Takahashi, H. & Shimizu, T. 1997. Role of intercellular contacts in generating an asymmetric mitotic apparatus in the *Tubifex* embryo. *Development Growth & Differentiation*, 39(3), 351-362.
- Theuer, M., Pilz, I., Schwarz, E., Wilhelm, P., Mainwaring, M.G. & Vinogradov, S.N. 1985. Molecular-size and shape of the native and reassociated forms of the extracellular hemoglobin of *Tubifex tubifex*. *International Journal of Biological Macromolecules*, 7(1), 25-29.
- Tian, Y. & Lu, Y.B. 2010. Simultaneous nitrification and denitrification process in a new *Tubificidae* reactor for minimizing nutrient release during sludge reduction. *Water Research*, 44(20), 6031-6040.
- Tian, Y., Lu, Y.B., Chen, L. & Lin, H.L. 2010. Optimization of process conditions with attention to the sludge reduction and stable immobilization in a novel *Tubificidae* reactor. *Bioresource Technology*, 101(15), 6069-6076.
- Tichy, M., Rucki, M., Hanzlikova, I. & Roth, Z. 2007. The *Tubifex tubifex* assay for the determination of acute toxicity. *Atla-Alternatives to Laboratory Animals*, 35(2), 229-237.
- Timm, T. 1998. *Lumbriculidae* (Oligochaeta) of Lake Taimyr. *Journal of Natural History*, 32(9), 1291-1301.
- Tolgyessy, P. 1988. The degradation of bentazone and chlorotoluron in aqueous-solutions by gamma-radiation – biodegradability and toxicity to *Tubifex tubifex* of radiolysis products. *Journal of Radioanalytical and Nuclear Chemistry-Letters*, 128(4), 321-329.
- Tweeten, K.A. & Abitz, A. 2012. Patterns of cleavage and gastrulation in embryos of freshwater oligochaetes from the *Lumbriculus* complex. *Integrative and Comparative Biology*, 52, E341-E341.
- Tweeten, K.A. & Reiner, A. 2012. Characterization of serine proteases of *Lumbriculus variegatus* and their role in regeneration. *Invertebrate Biology*, 131(4), 322-332.
- Urbisz, A.Z., Chajec, L. & Swiatek, P. 2015. The ovary of *Tubifex tubifex* (Clitellata, Naididae, Tubificinae) is composed of one, huge germ-line cyst that is enriched with cytoskeletal components. *Plos One*, 10(5).
- Varela, M.E. 1990. Taxonomic and ecological notes about some oligochaetes of fresh-water of northeastern Argentina. 1. Naididae. *Studies on Neotropical Fauna and Environment*, 25(4), 223-233.
- Verma, S., Das, S. & Khangarot, B.S. 2011. Toxicity of metallic oxides nanoparticle suspensions to a freshwater sludge worm *Tubifex tubifex* Müller. *Journal of Biomedical Nanotechnology*, 7(1), 216-217.
- Vivien, R., Wyler, S., Lafont, M. & Pawłowski, J. 2015. Molecular barcoding of aquatic oligochaetes: Implications for biomonitoring. *Plos One*, 10(4).
- Volpers, M. & Neumann, D. 2005. Tolerance of two tubificid species (*Tubifex tubifex* and *Limnodrilus hoffmeisteri*) to hypoxic and sulfidic conditions in novel, long-term experiments. *Archiv Fur Hydrobiologie*, 164(1), 13-38.
- Wang, J.F., Maul, J.D., Mayer, G.D. & Cobb, G.P. 2012. Uptake of fullerene (C60) induced catalase (CAT) activity in *Lumbriculus variegatus*. *Abstracts of Papers of the American Chemical Society*, 244.
- Wang, J.F., Wages, M., Yu, S.Y., Maul, J.D., Mayer, G., Hope-Weeks, L. & Cobb, G.P. 2014. Bioaccumulation of fullerene (C-60) and corresponding catalase elevation in *Lumbriculus variegatus*. *Environmental Toxicology and Chemistry*, 33(5), 1135-1141.
- West, C.W., Ankley, G.T., Nichols, J.W., Elonen, G.E. & Nessa, D.E. 1997. Toxicity and bioaccumulation of 2,3,7,8-tetrachlorodibenzo-p-dioxin in long-term tests with the freshwater benthic invertebrates *Chironomus tentans* and *Lumbriculus variegatus*. *Environmental Toxicology and Chemistry*, 16(6), 1287-1294.
- Wetzel, M.J. 1987. *Limnodrilus tortilipenis*, a new north american species of fresh-water *Tubificidae* (Annelida: Clitellata: Oligochaeta). *Proceedings of the Biological Society of Washington*, 100(1), 182-185.
- Wiederholm, T. & Dave, G. 1989. Toxicity of metal polluted sediments to *Daphnia magna* and *Tubifex tubifex*. *Hydrobiologia*, 176, 411-417.
- Wiederholm, T., Wiederholm, A.M. & Milbrink, G. 1987. Bulk sediment bioassays with 5 species of fresh-water oligochaetes. *Water Air and Soil Pollution*, 36(1-2), 131-154.
- Wiegand, C., Pehkonen, S., Akkanen, J., Penttinen, O.P. & Kukkonen, J.V.K. 2007. Bioaccumulation of paraquat by *Lumbriculus variegatus* in the presence of dissolved natural organic matter and impact on energy costs, biotransformation and antioxidative enzymes. *Chemosphere*, 66(3), 558-566.
- Wiramanaden, C.I.E., Orr, P.L. & Russel, C.K. 2015. Assessment of radium-226 bioavailability and bioaccumulation downstream of decommissioned uranium operations, using the caged oligochaete (*Lumbriculus variegatus*). *Environmental Toxicology and Chemistry*, 34(3), 507-517.
- Yanez, E., Cuadrado, S. & Martinez-Ansemil, E. 2006. External sense organs in freshwater oligochaetes (Annelida, Clitellata) revealed by scanning electron microscopy. *Journal of Morphology*, 267(2), 198-207.
- Yoshioka, Y., Ose, Y. & Sato, T. 1986. Testing and evaluation of chemical toxicity on *Tubifex*. *Eisei Kagaku – Japanese Journal of Toxicology and Environmental Health*, 32(4), 308-311.
- You, J., Brennan, A. & Lydy, M.J. 2009. Bioavailability and biotransformation of sediment-associated pyrethroid insecticides in *Lumbriculus variegatus*. *Chemosphere*, 75(11), 1477-1482.
- Zattara, E.E. & Bely, A.E. 2015. Fine taxonomic sampling of nervous systems within Naididae (Annelida: Clitellata) reveals evolutionary lability and revised homologies of annelid neural components. *Frontiers in Zoology*, 12.
- Zhang, L., Liao, Q.J.H., Gu, X.Z., He, W., Zhang, Z. & Fan, C.X. 2014a. Oxygen and phosphorus dynamics in freshwater sediment after the deposition of flocculated cyanobacteria and the role of Tubificid worms. *Journal of Hazardous Materials*, 266, 1-9.
- Zhang, L., Shang, J.G., He, W., You, B.S. & Fan, C.X. 2014b. The role of Tubificid worms (*Limnodrilus hoffmeisteri*) in sediment resuspension: a microcosm study. *Annales de Limnologie – International Journal of Limnology*, 50(3), 253-260.
- Zhang, X.F., Liu, Z.W., Jeppesen, E. & Taylor, W.D. 2014c. Effects of deposit-feeding Tubificid worms and filter-feeding bivalves on benthic-pelagic coupling: Implications for the restoration of eutrophic shallow lakes. *Water Research*, 50, 135-146.
- Zoran, M.J. & Drewes, C.D. 1988. The rapid tail withdrawal reflex of the tubificid worm, *Branchiura sowerbyi*. *Journal of Experimental Biology*, 137, 487-500.

Zoran, M.J., Drewes, C.D., Fourtner, C.R. & Siegel, A.J. 1988. The lateral giant fibers of the tubificid worm, *Branchiura sowerbyi* – structural and functional asymmetry in a paired interneuronal system. *Journal of Comparative Neurology*, 275(1), 76-86.

SCOPUS

- Achurra, A., Des Chatelliers, M.C. & Rodriguez, P. 2012. *Troglodrilus jugeti* sp. nov. (Annelida, Clitellata, Tubificinae), a new stygobiont oligochaete species from south-western europe. *Zootaxa*, (3229), 35-46. Retrieved from: www.scopus.com
- Achurra, A., Elejalde, M.A. & Rodriguez, P. 2011. Phylogenetic analysis of oligochaete Tubificinae (Annelida: Clitellata) based on mitochondrial sequence data. *Invertebrate Systematics*, 25(3), 208-218. doi: [10.1071/IS10040](https://doi.org/10.1071/IS10040)
- Achurra, A. & Rodriguez, P. 2008. Biodiversity of groundwater oligochaetes from a karst unit in northern Iberian Peninsula: Ranking subterranean sites for conservation management. *Hydrobiologia*, 605(1), 159-171. doi: [10.1007/s10750-008-9331-2](https://doi.org/10.1007/s10750-008-9331-2)
- Achurra, A., Rodriguez, P. & Reynoldson, T.B. 2015. Is the cantabrian region of northern Spain a biodiversity hotspot for obligate groundwater fauna? The case of oligochaetes (Annelida, Clitellata). *Hydrobiologia*, 745(1), 151-166. doi: [10.1007/s10750-014-2101-4](https://doi.org/10.1007/s10750-014-2101-4)
- Alejandra Pujals, M. 1985. Especies de los géneros *Pristina Ehrenberg, 1828* y *Bratislavia Kosel, 1976* (Oligochaeta: Naididae) en la provincia de Buenos Aires, Argentina. *Studies on Neotropical Fauna and Environment*, 20(4), 203-210. doi: [10.1080/01650528509360690](https://doi.org/10.1080/01650528509360690)
- Alexander, A.C., Culp, J.M., Liber, K. & Cessna, A.J. 2007. Effects of insecticide exposure on feeding inhibition in mayflies and oligochaetes. *Environmental Toxicology and Chemistry*, 26(8), 1726-1732. doi: [10.1897/07-015R.1](https://doi.org/10.1897/07-015R.1)
- Alves, R.D.G., Marchese, M.R. & Escarpinatti, S.C. 2006. Oligochaeta (Annelida, Clitellata) in lotic environments in the state of São Paulo, Brazil. *Iheringia – Série Zoologia*, 96(4), 431-435. doi: [10.1590/S0073-47212006000400007](https://doi.org/10.1590/S0073-47212006000400007)
- Alves, R.D.G., Marchese, M.R. & Martins, R.T. 2008. Oligochaeta (Annelida, Clitellata) of lotic environments at Parque Estadual Intervales (São Paulo, Brazil). *Biota Neotropica*, 8(1), 69-72. doi: [10.1590/S1676-06032008000100009](https://doi.org/10.1590/S1676-06032008000100009)
- Anderson, R.V. & Holm, D.J. 1987. *Chaetogaster limnaei* (Oligochaeta: Naididae) infesting unionid mollusks (Pelecypoda: Unionidae) and corbicula fluminea (Pelecypoda: Corbiculidae) in pool 19, Mississippi River. *Journal of Freshwater Ecology*, 4(1), 61-64. doi: [10.1080/02705060.1987.9665161](https://doi.org/10.1080/02705060.1987.9665161)
- Andrews, J.M., Childress, J.N., Iakovidis, T.J. & Langford, G.J. 2015. Elucidating the life history and ecological aspects of *Allodero hylae* (Annelida: Clitellata: Naididae), a parasitic oligochaete of invasive cuban tree frogs in Florida. *Journal of Parasitology*, 101(3), 275-281. doi: [10.1645/14-608.1](https://doi.org/10.1645/14-608.1)
- Ankley, G.T., Leonard, E.N. & Mattson, V.R. 1994. Prediction of bioaccumulation of metals from contaminated sediments by the oligochaete, *Lumbriculus variegatus*. *Water Research*, 28(5), 1071-1076. doi: [10.1016/0043-1354\(94\)90192-9](https://doi.org/10.1016/0043-1354(94)90192-9)
- Anlauf, A. 1997. Enzyme variability of *Tubifex tubifex* (Müller) (Oligochaeta, Tubificidae) and seven other tubificid species. *Archiv Fur Hydrobiologie*, 139(1), 83-100. Retrieved from: www.scopus.com
- Anlauf, A. 1994. Some characteristics of genetic variants of *Tubifex tubifex* (Müller, 1774) (Oligochaeta: Tubificidae) in laboratory cultures. *Hydrobiologia*, 278(1-3), 1-6. doi: [10.1007/BF00142306](https://doi.org/10.1007/BF00142306)
- Antonio, D.B., Andree, K.B., McDowell, T.S. & Hedrick, R.P. 1998. Detection of *Myxobolus cerebralis* in rainbow trout and oligochaete tissues by using a nonradioactive in situ hybridization (ISH) protocol. *Journal of Aquatic Animal Health*, 10(4), 338-347. doi: [10.1577/1548-8667\(1998\)010<0338:DOMCIR>2.0.CO;2](https://doi.org/10.1577/1548-8667(1998)010<0338:DOMCIR>2.0.CO;2)
- Arimoro, F.O., Ikomi, R.B. & Iwegbue, C.M.A. 2007. Ecology and abundance of oligochaetes as indicators of organic pollution in an urban stream in southern Nigeria. *Pakistan Journal of Biological Sciences*, 10(3), 446-453. Retrieved from: www.scopus.com
- Arkhipova, N.R. 1996. Morphology of pectinate setae in Tubificids (Tubificidae, Oligochaeta). *Zoologicheskii Zhurnal*, 75(2), 186-187. Retrieved from: www.scopus.com
- Armendáriz, L., Ocón, C. & Rodrigues Capítulo, A. 2012. Potential responses of oligochaetes (Annelida, Clitellata) to global changes: Experimental fertilization in a lowland stream of Argentina (South America). *Limnologica*, 42(2), 118-126. doi: [10.1016/j.limno.2011.09.005](https://doi.org/10.1016/j.limno.2011.09.005)
- Armendariz, L.C. 1999. Population dynamics of *Allonais lairdi* (Oligochaeta, Naididae) from Los Talas, Buenos Aires Province. [Dinamica poblacional de *Allonais lairdi* (Oligochaeta, Naididae) en Los Talas, Provincia de Buenos Aires]. *Ecología Austral*, 9(1-2), 20-27. Retrieved from: www.scopus.com
- Armendariz, L.C. 2008. Life cycle of *Dero* (*Aulophorus*) *costatus* Marcus, 1944 (Tubificidae, Oligochaeta) in a vegetated pond at Los Talas, Argentina. [Ciclo de vida de *Dero* (*Aulophorus*) *costatus* Marcus, 1944 (Tubificidae, Oligochaeta) en un cuerpo de agua con vegetacion flotante en Los Talas, Argentina]. *Gayana*, 72(1), 23-30. Retrieved from: www.scopus.com
- Armendariz, L.C. 2000. Population dynamics of *Stylaria lacustris* (Linnaeus, 1767) (Oligochaeta, Naididae) in Los Talas, Argentina. *Hydrobiologia*, 438, 217-226. doi: [10.1023/A:1004139622036](https://doi.org/10.1023/A:1004139622036)
- Armendariz, L.C. & César, I.I. 2001. The distribution and ecology of littoral Oligochaeta and Aphanoneura (Annelida) of the natural and historical reserve of Isla Martín García, Río de la Plata River, Argentina. *Hydrobiologia*, 463, 207-216. doi: [10.1023/A:1013120128722](https://doi.org/10.1023/A:1013120128722)
- Arslan, N. 2006. Records of Aphanoneura and aquatic oligochaetes from Turkey. *Fresenius Environmental Bulletin*, 15(4), 249-254. Retrieved from: www.scopus.com
- Arslan, N., Koç, B. & Çiçek, A. 2010. Metal contents in water, sediment, and Oligochaeta Chironomidae of Lake Ulubat, a ramsar site of Turkey. *The Scientific World Journal*, 10, 1269-1281. doi: [10.1100/tsw.2010.117](https://doi.org/10.1100/tsw.2010.117)
- Arslan, N. & Şahin, Y. 2006. A preliminary study on the identification of the littoral oligochaete (Annelida) and Chironomidae (Diptera) fauna of Lake Kovada, a national park in Turkey. *Turkish Journal of Zoology*, 30(1), 67-72. Retrieved from: www.scopus.com
- Arslan, N. & Şahin, Y. 2004. First records of some Naididae (Oligochaeta) species for Turkey. [Türkiye İçin bazı Naididae (Oligochaeta) Türlerinin ilk Kayıtları]. *Turkish Journal of Zoology*, 28(1), 7-18. Retrieved from: www.scopus.com
- Arslan, N. & Şahin, Y. 2003. Two new records of *Aulodrilus Bretscher, 1899* (Oligochaeta, Tubificidae) for the turkish fauna. [Türkiye Faunası İçin İki Yeni *Aulodrilus Bretscher, 1899* (Oligochaeta, Tubificidae) Türü]. *Turkish Journal of Zoology*, 27(4), 275-280. Retrieved from: www.scopus.com
- Arslan, N., Timm, T. & Erséus, C. 2007. Aquatic Oligochaeta (Annelida) of Ballıdamlı wetland (Turkey), with description of two new species of Phalodrilinae. *Biology*, 62(3), 323-334. doi: [10.2478/s11756-007-0055-y](https://doi.org/10.2478/s11756-007-0055-y)
- Atanacković, A., Šporka, F., Tomović, J., Vasiljević, B., Marković, V., Simić, V. & Paunović, M. 2012. First record of *Bothrioneurum vejdovskyanum* Štolc, 1886 (Oligochaeta, Tubificidae) in Serbia. *Archives of Biological Sciences*, 64(3), 1123-1126. doi: [10.2298/ABS1203123A](https://doi.org/10.2298/ABS1203123A)
- Atanacković, A.D., Šporka, F., Csányi, B., Vasiljević, B.M., Tomović, J.M. & Paunović, M.M. 2013. Oligochaeta of the Danube River – a faunistical

- review. *Biologia (Poland)*, 68(2), 269-277. doi: [10.2478/s11756-013-0155-9](https://doi.org/10.2478/s11756-013-0155-9)
- Back, H. & Prosi, F. 1985. Distribution of inorganic cations in Limnodrilus udekemianus (Oligochaeta, Tubificidae) using laser induced microprobe mass analysis, with special emphasis on heavy metals. *Micron and Microscopica Acta*, 16(3), 145-150. doi: [10.1016/0739-6260\(85\)90058-9](https://doi.org/10.1016/0739-6260(85)90058-9)
- Balatre-Veltz, I., Biagianti-Risbourg, S. & Vernet, G. 1999. On the cephalic regeneration of Lumbriculus variegatus Müller 1774 (Annelida, Oligochaeta). [Sur la régénération céphalique de Lumbriculus variegatus Müller, 1774 (Annelida, Oligochaeta)]. *Bulletin de la Societe Zoologique de France*, 124(1), 101-109. Retrieved from: www.scopus.com
- Baldo, L. & Ferraguti, M. 2005. Mixed reproductive strategy in *Tubifex tubifex* (Oligochaeta, Tubificidae)? *Journal of Experimental Zoology – Part A: Comparative Experimental Biology*, 303(2), 168-177. doi: [10.1002/jeza.144](https://doi.org/10.1002/jeza.144)
- Balık, S., Ustaoglu, M.R. & Yildiz, S. 2004. Oligochaeta and Aphanoneura (Annelida) fauna of the gediz delta (Menemen-Izmir). *Turkish Journal of Zoology*, 28(3), 183-197. Retrieved from: www.scopus.com
- Baturina, M. 2012. Distribution and diversity of aquatic Oligochaeta in small streams of the middle taiga. *Turkish Journal of Zoology*, 36(1), 75-84. doi: [10.3906/zoo-1002-64](https://doi.org/10.3906/zoo-1002-64)
- Bauer-Hilty, A., Dallinger, R. & Berger, B. 1989. Isolation and partial characterization of a cadmium-binding protein from *Lumbriculus variegatus* (Oligochaeta, Annelida). *Comparative Biochemistry and Physiology – Part C: Toxicology and Pharmacology*, 94(2), 373-379. doi: [10.1016/0742-8413\(89\)90085-6](https://doi.org/10.1016/0742-8413(89)90085-6)
- Bazzanti, M. & Lafont, M. 1985. Variability in the spermathecal setae of *Potamothrix heuscheri* (Bretschner) (Oligochaeta, Tubificidae) in Lake Nemi and notes on the relationship between this species and the trophic level of lakes. [Variabilité des soies spermathécales de *Potamothrix heuscheri* (Bretschner) (Oligochaeta, Tubificidae) dans le lac de Nemi et remarques sur la relation entre cette espèce et le degré de trophie des lacs]. *Annales de Limnologie*, 21(2), 107-115. doi: [10.1051/limn/1985010](https://doi.org/10.1051/limn/1985010)
- Beauchamp, K.A., Kathman, R.D., McDowell, T.S. & Hedrick, R.P. 2001. Molecular phylogeny of tubificid oligochaetes with special emphasis on *Tubifex tubifex* (Tubificidae). *Molecular Phylogenetics and Evolution*, 19(2), 216-224. doi: [10.1006/mpev.2001.0923](https://doi.org/10.1006/mpev.2001.0923)
- Behrend, R.D.L., Fernandes, S.E.P., Fujita, D.S. & Takeda, A.M. 2009. Eight years of monitoring aquatic Oligochaeta from the Baía and Ivinhema Rivers. [Oito anos de monitoramento de Oligochaeta aquáticas nos Rios Baía e Ivinhema]. *Brazilian Journal of Biology*, 69 (suppl. 2), 559-571. Retrieved from: www.scopus.com
- Behrend, R.D.L., Takeda, A.M., Gomes, L.C. & Fernandes, S.E.P. 2012. Using Oligochaeta assemblages as an indicator of environmental changes. [Usando assembléias de Oligochaeta como um indicador de mudanças ambientais]. *Brazilian Journal of Biology*, 72(4), 873-884. doi: [10.1590/S1519-69842012000500014](https://doi.org/10.1590/S1519-69842012000500014)
- Behrend, R.D.L., Teixeira, M.C., Fernandes, S.E.P., Camargo, J.C., Rosin, G.C. & Takeda, A.M. 2013. Effects of a native and a non-native macrophyte species of hydrocharitaceae on CHIRONOMIDAE and Oligochaeta assemblages structure. [Efeitos de uma espécie de macrófita nativa e uma invasora de Hydrocharitaceae sobre a estrutura das assembleias de Chironomidae e Oligochaeta]. *Acta Scientiarum – Biological Sciences*, 35(3), 351-358. doi: [10.4025/actascibiolsci.v35i3.10855](https://doi.org/10.4025/actascibiolsci.v35i3.10855)
- Bely, A.E. 1999. Decoupling of fission and regenerative capabilities in an asexual oligochaete. *Hydrobiologia*, 406, 243-251. doi: [10.1023/A:1003763915697](https://doi.org/10.1023/A:1003763915697)
- Bergter, A., Hunnekuhl, V.S., Schniederjans, M. & Paululat, A. 2007. Evolutionary aspects of pattern formation during clitellate muscle development. *Evolution and Development*, 9(6), 602-617. doi: [10.1111/j.1525-142X.2007.00184.x](https://doi.org/10.1111/j.1525-142X.2007.00184.x)
- Bernoud, S., Lafont, M. & Durbec, A. 2003. Ecological diagnostic methods based on benthic Oligochaeta communities; technical and economic viewpoints. [Diagnostic écologique des sédiments à partir des peuplements d'oligochètes; aspects techniques et économiques]. *Techniques – Sciences – Methodes*, (5), 69-76. Retrieved from: www.scopus.com
- Bingham, C.R. & Miller, A.C. 1989. Colonization of a man-made gravel bar by Oligochaeta. *Hydrobiologia*, 180(1), 229-234. doi: [10.1007/BF00027555](https://doi.org/10.1007/BF00027555)
- Blettler, M., Amsler, M., De Drago, I.E. & Marchese, M. 2008. Effects of stream hydraulics and other environmental variables on density of Narapa bonettoi (Oligochaeta) in the Paraná River system. *River Research and Applications*, 24(8), 1124-1140. doi: [10.1002/rra.1115](https://doi.org/10.1002/rra.1115)
- Bock, S., Sedlmeier, U.A. & Hoffmann, K.H. 1989. Metabolism of absorbed short-chain carboxylic acids by the freshwater oligochaete *Tubifex tubifex*. *Comparative Biochemistry and Physiology – Part B: Biochemistry and Molecular Biology*, 92(1), 35-40. doi: [10.1016/0305-0491\(89\)90309-X](https://doi.org/10.1016/0305-0491(89)90309-X)
- Boi, S. & Ferraguti, M. 2001. Temporal pattern of the double sperm line production in *Tubifex tubifex* (Annelida, Oligochaeta). *Hydrobiologia*, 463, 103-106. doi: [10.1023/A:1013191205524](https://doi.org/10.1023/A:1013191205524)
- Bojková, J., Schenková, J., Horská, M. & Hájek, M. 2011. Species richness and composition patterns of clitellate (Annelida) assemblages in the treeless spring fens: The effect of water chemistry and substrate. *Hydrobiologia*, 667(1), 159-171. doi: [10.1007/s10750-011-0634-3](https://doi.org/10.1007/s10750-011-0634-3)
- Bonacina, C., Pasteris, A., Bonomi, G. & Marzuoli, D. 1994. Quantitative observations on the population ecology of Branchiura sowerbyi (Oligochaeta, Tubificidae). *Hydrobiologia*, 278(1-3), 267-274. doi: [10.1007/BF00142334](https://doi.org/10.1007/BF00142334)
- Bonacinal, C., Bonomi, G. & Monti, C. 1987. Progress in cohort cultures of aquatic Oligochaeta. *Hydrobiologia*, 155(1), 163-169. doi: [10.1007/BF00025645](https://doi.org/10.1007/BF00025645)
- Bonomi, G. & Pasteris, A. 2006. From demographic strategies to mathematical models: Trends in population dynamics studies of aquatic Oligochaeta. *Hydrobiologia*, 564 (1 spec. iss.), 61-71. doi: [10.1007/s10750-005-1708-x](https://doi.org/10.1007/s10750-005-1708-x)
- Bouguenec, V. & Giani, N. 1989. Aquatic Oligochaeta as prey for invertebrates and vertebrates: A review. [Les Oligochètes aquatiques en tant que proies des Invertébrés et des Vertébrés: une revue]. *Acta Oecologica – Oecologia Applicata*, 10(3), 177-196. Retrieved from: www.scopus.com
- Boumaiza, M., Giani, N. & Martinez-Ansemil, E. 1986. Oligochaeta and Aphanoneura in the running waters of Tunisia. II. A key for determining the species presently tallied. [Les Oligochètes et Aphanoneura des eaux courantes de Tunisie. II. Clé pour la détermination des espèces actuellement recensées]. *Archives de l’Institut Pasteur de Tunis*, 63(2-3), 299-323. Retrieved from: www.scopus.com
- Boumaiza, M., Martinez-Ansemil, E. & Giani, N. 1986. Oligochaetes and Aphanoneura in streams of Tunisia. I. Faunistic data. [Les Oligochètes et Aphanoneura des eaux courantes de Tunisie. I – Données faunistiques]. *Annales de Limnologie*, 22(3), 231-237. doi: [10.1051/limn/1986021](https://doi.org/10.1051/limn/1986021)
- Bowker, D.W., Wareham, M.T. & Learner, M.A. 1985. A choice chamber experiment on the selection of algae as food and substrata by *Nais elinguis* (Oligochaeta: Naididae). *Freshwater Biology*, 15(5), 547-557. doi: [10.1111/j.1365-2427.1985.tb00225.x](https://doi.org/10.1111/j.1365-2427.1985.tb00225.x)
- Bowker, D.W., Wareham, M.T. & Learner, M.A. 1985. The ingestion and assimilation of algae by *Nais elinguis* (Oligochaeta: Naididae). *Oecologia*, 67(2), 282-285. doi: [10.1007/BF00384300](https://doi.org/10.1007/BF00384300)
- Brinkhurst, R.O. & Nemec, A.F.L. 1987. A comparison of phenetic and phylogenetic methods applied to the systematics of Oligochaeta. *Hydrobiologia*, 155(1), 65-74. doi: [10.1007/BF00025632](https://doi.org/10.1007/BF00025632)

- Brinkhurst, R.O., Qi Sang & Liang Yanling. 1990. The aquatic Oligochaeta from the people's Republic of China. *Canadian Journal of Zoology*, 68(5), 901-916. doi: [10.1139/z90-131](https://doi.org/10.1139/z90-131)
- Brown, L., Finston, T., Humphreys, G., Eberhard, S. & Pinder, A. 2015. Groundwater oligochaetes show complex genetic patterns of distribution in the pilbara region of western Australia. *Invertebrate Systematics*, 29(5), 405-420. doi: [10.1071/IS14037](https://doi.org/10.1071/IS14037)
- Brunson, E.L., Canfield, T.J., Dwyer, F.J., Ingersoll, C.G. & Kemble, N.E. 1998. Assessing the bioaccumulation of contaminants from sediments of the upper Mississippi River using field-collected oligochaetes and laboratory exposed *Lumbriculus variegatus*. *Archives of Environmental Contamination and Toxicology*, 35(2), 191-201. doi: [10.1007/s002449900367](https://doi.org/10.1007/s002449900367)
- Bunke, D. 1998. Ultrastructure of the nephrostome in *Enchytraeus albidus* (Annelida, Clitellata). *Zoomorphology*, 118(3), 177-182. doi: [10.1007/s004350050067](https://doi.org/10.1007/s004350050067)
- Bunke, D. 2000. Ultrastructure of the preseptal part of metanephridia in *Nais variabilis* and *Dero digitata* (Annelida, Clitellata). *Zoomorphology*, 120(1), 39-46. doi: [10.1007/s004350000023](https://doi.org/10.1007/s004350000023)
- Cakić, P., Petrović, Z., Kataranovski, D., Jakovčev, D. & Lenhardt, M. 2000. The first record and description of *Chaetogaster limnaei* von Baer, 1827 (Naididae, Oligochaeta) on huso huso fry in Serbia. *Helminthologia*, 37(3), 162-164. Retrieved from: www.scopus.com
- Campitelli-Ramos, R., Lucca, J.V., Oliveira, L.L.D., Marchese, M.R. & Rocha, O. 2014. First record of *Dero* (*Aulophorus*) *Bimagnasetus harman* (Oligochaeta) from Brazil and habitat characteristics. [Primeiro registro de *Dero* (*Aulophorus*) *Bimagnasetus harman* (Oligochaeta) do Brasil e notas ecológicas]. *Brazilian Journal of Biology*, 74(2), 483-488. doi: [10.1590/1519-6984.14512](https://doi.org/10.1590/1519-6984.14512)
- Çapraz, S. & Arslan, N. 2005. The Oligochaeta (Annelida) fauna of Aksu stream (Antalya). *Turkish Journal of Zoology*, 29(3), 229-236. Retrieved from: www.scopus.com
- Carrasco Navarro, V., Brozinski, J.-., Leppänen, M.T., Honkanen, J.O., Kronberg, L. & Kukkonen, J.V. 2011. Inhibition of pyrene biotransformation by piperonyl butoxide and identification of two pyrene derivatives in *Lumbriculus variegatus* (Oligochaeta). *Environmental Toxicology and Chemistry*, 30(5), 1069-1078. doi: [10.1002/etc.485](https://doi.org/10.1002/etc.485)
- Carrasco Navarro, V., Leppänen, M.T., Honkanen, J.O. & Kukkonen, J.V.K. 2012. Trophic transfer of pyrene metabolites and nonextractable fraction from oligochaete (*Lumbriculus variegatus*) to juvenile brown trout (*Salmo Trutta*). *Chemosphere*, 88(1), 55-61. doi: [10.1016/j.chemosphere.2012.02.060](https://doi.org/10.1016/j.chemosphere.2012.02.060)
- Casellato, S. 1999. Oligochaetes of Karavasta Lagoon (Albania). Preliminary results. *Hydrobiologia*, 406, 175-182. doi: [10.1023/A:1003766527084](https://doi.org/10.1023/A:1003766527084)
- Casellato, S., Aiello, R. & Negrisolo, P.A. 1992. Long-term experiment on Branchiura sowerbyi Beddard (Oligochaeta, Tubificidae) using sediment treated with LAS (Linear Alkylbenzene Sulphonate). *Hydrobiologia*, 232(2), 169-173. doi: [10.1007/BF00017476](https://doi.org/10.1007/BF00017476)
- Casellato, S. & Caneva, F. 1994. Composition and distribution of bottom oligochaete fauna of a north Italian eutrophic lake (Lake Ledro). *Hydrobiologia*, 278(1-3), 87-92. doi: [10.1007/BF00142315](https://doi.org/10.1007/BF00142315)
- Casellato, S., Del Piero, S., Masiero, L. & Covre, V. 2013. Fluoride toxicity and its effects on gametogenesis in the aquatic oligochaete Branchiura sowerbyi Beddard. *Fluoride*, 46(1), 7-18. Retrieved from: www.scopus.com
- Cellot, B. & Juget, J. 1998. Oligochaete drift in a large river (French Upper Rhône): The effect of life cycle and discharge. *Hydrobiologia*, 389(1-3), 183-191. Retrieved from: www.scopus.com
- Ceretti, G. & Nocentini, A.M. 1996. Notes on the distribution of some macrobenthonic populations (Oligochaeta and diptera Chironomidae) in the littoral of a few small lakes in northern Italy. *Memorie – Istituto Italiano Di Idrobiologia*, 54, 109-124. Retrieved from: www.scopus.com
- César, I.I. 2014. Annelida (Oligochaeta and Aphanoneura) from the natural reserve of Isla Martín García (upper Río de la Plata Estuary, Argentina): Biodiversity and response to environmental variables. [Annelida (Oligochaeta e Aphanoneura) da reserva natural de usos múltiplos Ilha Martín García, Río de la Plata: Biodiversidade e resposta às variáveis ambientais]. *Brazilian Journal of Biology*, 74(1), 128-136. doi: [10.1590/1519-6984.21412](https://doi.org/10.1590/1519-6984.21412)
- Chapman, K.K., Benton, M.J., Brinkhurst, R.O. & Scheuerman, P.R. 1999. Use of the aquatic oligochaetes *Lumbriculus variegatus* and *Tubifex tubifex* for assessing the toxicity of copper and cadmium in a spiked-artificial-sediment toxicity test. *Environmental Toxicology*, 14(2), 271-278. doi: [10.1002/\(SICI\)1522-7278\(199905\)14:2<271::AID-TOX8>3.0.CO;2-F](https://doi.org/10.1002/(SICI)1522-7278(199905)14:2<271::AID-TOX8>3.0.CO;2-F)
- Chapman, P.M. & Brinkhurst, R.O. 1987. Hair today, gone tomorrow: Induced chaetal changes in tubificid oligochaetes. *Hydrobiologia*, 155(1), 45-55. doi: [10.1007/BF00025630](https://doi.org/10.1007/BF00025630)
- Chapman, P.M. & Mitchell, D.G. 1986. Acute tolerance tests with the oligochaetes *Nais communis* (Naididae) and *ilyodrilus frantzi* (Tubificidae). *Hydrobiologia*, 137(1), 61-64. doi: [10.1007/BF00004173](https://doi.org/10.1007/BF00004173)
- Chauvet, E., Giani, N. & Gessner, M.O. 1993. Breakdown and invertebrate colonization of leaf litter in two contrasting streams: Significance of oligochaetes in a large river. *Canadian Journal of Fisheries and Aquatic Sciences*, 50(3), 488-495. doi: [10.1139/f93-057](https://doi.org/10.1139/f93-057)
- Chen, J. & Xie, P. 2008. Accumulation of hepatotoxic microcystes in freshwater mussels, aquatic insect larvae and oligochaetes in a large, shallow eutrophic lake (Lake Chaohu) of subtropical China. *Fresenius Environmental Bulletin*, 17 (7 A), 849-854. Retrieved from: www.scopus.com
- Christensen, B. & Theisen, B.F. 1998. Phylogenetic status of the family Naididae (Oligochaeta, Annelida) as inferred from DNA analyses. *Journal of Zoological Systematics and Evolutionary Research*, 36(4), 169-172. Retrieved from: www.scopus.com
- Christoffersen, M.L. 2012. Checklist of questidae (Clitellata) and aeolosomatidae (Oligochaeta) from South America. *Turkish Journal of Zoology*, 36(1), 133-138. doi: [10.3906/zoo-1002-26](https://doi.org/10.3906/zoo-1002-26)
- Ciparis, S. & Hale, R.C. 2005. Bioavailability of polybrominated diphenyl ether flame retardants in biosolids and spiked sediment to the aquatic oligochaete, *Lumbriculus variegatus*. *Environmental Toxicology and Chemistry*, 24(4), 916-925. doi: [10.1897/04-179R.1](https://doi.org/10.1897/04-179R.1)
- Coastes, K.A. 1987. Phylogenetics of some Enchytraeidae (Annelida: Oligochaeta): A preliminary investigation of relationships to the Haplotaenidae. *Hydrobiologia*, 155(1), 91-106. doi: [10.1007/BF00025634](https://doi.org/10.1007/BF00025634)
- Coates, K.A. & Stacey, D.F. 1994. Oligochaetes (Naididae, Tubificidae, Enchytraeidae and Alluroididae) of Guyana, Peru and Ecuador. *Hydrobiologia*, 278(1-3), 79-84. doi: [10.1007/BF00142313](https://doi.org/10.1007/BF00142313)
- Collado, R. & De Mendoza, G. 2009. *Environmental factors and distribution of littoral oligochaetes in pyrenean lakes*. doi: [10.1127/adylim/62/2009/215](https://doi.org/10.1127/adylim/62/2009/215). Retrieved from: www.scopus.com
- Collado, R., Kasprzak, P. & Schmelz, R.M. 1999. Oligochaeta and Aphanoneura in two northern german hardwater lakes of different trophic state. *Hydrobiologia*, 406, 143-148. doi: [10.1023/A:1003700604272](https://doi.org/10.1023/A:1003700604272)
- Collado, R., Martínez-Ansemil, E. & Giani, N. 1993. The aquatic oligochaetes of the iberian peninsula: Description of *Stylodrilus curvithecus* sp. nov. (Lumbriculidae) and *Mesenchytraeus lusitanicus* sp. nov., with a redescription of *cognettia hibernica* healy (Enchytraeidae). [Les Oligochètes aquatiques de la Péninsule Ibérique: Description de *Stylodrilus curvithecus* sp. nov. (Lumbriculidae) et de *Mesenchytraeus lusitanicus* sp. nov. et redescription de *Cognettia hibernica* Healy (Enchytraeidae)]. *Annales de Limnologie*, 29(2), 129-138. doi: [10.1051/limn/1993012](https://doi.org/10.1051/limn/1993012)
- Collado, R. & Schmelz, R.M. 2001. Oligochaete distribution patterns in two german hardwater lakes of different trophic state. *Limnologica*, 31(4), 317-328. Retrieved from: www.scopus.com

- Collado, R. & Schmelz, R.M. 2000. Pedonais crassifaucis gen. nov., sp. nov. (Naididae) and Bothrioneurum righii sp. nov. (Tubificidae), two new tropical soil-dwelling species of “aquatic” oligochaetes (Clitellata, Annelida) from central Amazonia. *Amazoniana*, 16(1-2), 223-235. Retrieved from: www.scopus.com
- Collado, R. & Schmelz, R.M. 2000. Pristina silvicola and Pristina terrena sp. nov., two new soil-dwelling species of Naididae (Oligochaeta, Annelida) from the tropical rain forest near Manaus, Brazil, with comments on the genus Pristinella. *Journal of Zoology*, 251(4), 509-516. doi: [10.1017/S0952836900008104](https://doi.org/10.1017/S0952836900008104)
- Conn, D.B., Ricciardi, A., Babapulle, M.N., Klein, K.A. & Rosen, D.A. 1996. Chaetogaster limnaei (Annelida: Oligochaeta) as a parasite of the zebra mussel *dreissena polymorpha*, and the quagga mussel *dreissena bugensis* (Mollusca: Bivalvia). *Parasitology Research*, 82(1), 1-7. doi: [10.1007/s004360050058](https://doi.org/10.1007/s004360050058)
- Connell, D.W., Bowman, M. & Hawker, D.W. 1988. Bioconcentration of chlorinated hydrocarbons from sediment by oligochaetes. *Ecotoxicology and Environmental Safety*, 16(3), 293-302. doi: [10.1016/0147-6513\(88\)90058-9](https://doi.org/10.1016/0147-6513(88)90058-9)
- Conrad, A.U., Comber, S.D. & Simkiss, K. 2000. New method for the assessment of contaminant uptake routes in the oligochaete *Lumbriculus variegatus*. *Bulletin of Environmental Contamination and Toxicology*, 65(1), 16-21. doi: [10.1007/s001280000088](https://doi.org/10.1007/s001280000088)
- Corbi, J.J., Roque, F.O., Trivinho-Strixino, S. & Alves, R.G. 2005. Records of oligochaetes in freshwater sponges, on bryozoans, and on colonial hydrozoans from Brazil. *Brazilian Journal of Biology – Revista Brasileira de Biologia*, 65(1), 187-188. doi: [10.1590/S1519-6984200500010002](https://doi.org/10.1590/S1519-6984200500010002)
- Corteleazzi, A., Armendáriz, L.C., van Oosterom, M.V.L., Cepeda, R. & Capítulo, A.R. 2011. Different levels of taxonomic resolution in bioassessment: A case study of Oligochaeta in lowland streams. [Diferentes níveis de resolução taxonômica em biomonitoramento: Um estudo de caso de oligoquetados em rios de planícies]. *Acta Limnologica Brasiliensis*, 23(4), 412-425. Retrieved from: www.scopus.com
- Courtney, C.C. & Christensen, B.M. 1988. Host-parasite relationships of caryophyllaeid cestodes and aquatic oligochaetes. II. Effects of host age and mixed infections. *Journal of Parasitology*, 74(4), 573-581. doi: [10.2307/3282172](https://doi.org/10.2307/3282172)
- Croce, V., De Angelis, S., Patrolecco, L., Polesello, S. & Valsecchi, S. 2005. Uptake and accumulation of sediment associated 4-nonylphenol in a benthic invertebrate (*Lumbriculus variegatus*, freshwater oligochaete). *Environmental Toxicology and Chemistry*, 24(5), 1165-1171. doi: [10.1897/04-337R.1](https://doi.org/10.1897/04-337R.1)
- Cui, Y. & Wang, H. 2009. Three new species of Tubificinae, Oligochaeta, from two plateau lakes in southwest China. *Zootaxa*, (2143), 45-54. Retrieved from: www.scopus.com
- Cui, Y.- & Wang, H.-. 2005. Potamothrix scleropenis sp. nov. (Oligochaeta: Tubificidae) from Fuxian Lake, the deepest lake in southwest China. *Zoological Science*, 22(12), 1353-1357. doi: [10.2108/zsj.22.1353](https://doi.org/10.2108/zsj.22.1353)
- Dafoe, L.T., Rygh, A.L., Yang, B., Gingras, M.K. & Pemberton, S.G. 2011. A new technique for assessing tubificid burrowing activities, and recognition of biogenic grading formed by these oligochaetes. *Palaios*, 26(1), 66-80. doi: [10.2110/palo.2010.p10-023r](https://doi.org/10.2110/palo.2010.p10-023r)
- Danijela, S., Radujković, B.M. & Krpo-Ćetković, J. 2011. Catalogue of aquatic Oligochaeta (Annelida: Clitellata) of Montenegro, exclusive of Naidinae and Pristininae. *Zootaxa*, (2985), 1-25. Retrieved from: www.scopus.com
- de Eguileor, M., Valvassori, R., Lanzavecchia, G. & Giorgi, S. 1990. Body wall muscles in the haplotaxids *Haplotaenia gordiooides* and *Pelodrilus leruthi* (Annelida, Oligochaeta). *Zoomorphology*, 110(1), 27-36. doi: [10.1007/BF01632809](https://doi.org/10.1007/BF01632809)
- De Jonge, M., Eyckmans, M., Blust, R. & Bervoets, L. 2011. Are accumulated sulfide-bound metals metabolically available in the benthic oligochaete *Tubifex tubifex*? *Environmental Science & Technology*, 45(7), 3131-3137. doi: [10.1021/es1037595](https://doi.org/10.1021/es1037595)
- de Pascar, C.G. 1987. Aquatic Oligochaeta in some tributaries of the Río de la Plata, Buenos Aires, Argentina. *Hydrobiologia*, 144(2), 125-130. doi: [10.1007/BF00014526](https://doi.org/10.1007/BF00014526)
- Deeds, J.R. & Klerks, P.L. 1999. Metallothionein-like proteins in the freshwater oligochaete *Limnodrilus udekemianus* and their role as a homeostatic mechanism against cadmium toxicity. *Environmental Pollution*, 106(3), 381-389. doi: [10.1016/S0269-7491\(99\)00100-1](https://doi.org/10.1016/S0269-7491(99)00100-1)
- Des Châtelliers, M.C., Juget, J., Lafont, M. & Martin, P. 2009. Subterranean aquatic Oligochaeta. *Freshwater Biology*, 54(4), 678-690. doi: [10.1111/j.1365-2427.2009.02173.x](https://doi.org/10.1111/j.1365-2427.2009.02173.x)
- Di Maio, J. & Stacey, D.F. 1998. New distribution records of aquatic Oligochaeta (Annelida) from running waters in Ontario. *Canadian Field-Naturalist*, 112(1), 69-74. Retrieved from: www.scopus.com
- Ding, J., Drewes, C.D. & Hsu, W.H. 2001. Behavioral effects of ivermectin in a freshwater oligochaete, *Lumbriculus variegatus*. *Environmental Toxicology and Chemistry*, 20(7), 1584-1590. doi: [10.1002/etc.5620200724](https://doi.org/10.1002/etc.5620200724)
- Dózsa-Farkas, K. 1996. An interesting reproduction type in enchytraeids (Oligochaeta). *Acta Zoologica Academiae Scientiarum Hungaricae*, 42(1), 3-10. Retrieved from: www.scopus.com
- Dózsa-Farkas, K. 2009. Review of the fridericia species (Oligochaeta: Enchytraeidae) possessing two spermathecal diverticula and description of a new species. *Journal of Natural History*, 43(17-18), 1043-1065. doi: [10.1080/00222930902767441](https://doi.org/10.1080/00222930902767441)
- Dózsa-Farkas, K. 2010. Significance of using nephridia in the taxonomy of family Enchytraeidae: (Annelida: Oligochaeta). *Zoology in the Middle East*, 51, 41-53. doi: [10.1080/09397140.2010.10638456](https://doi.org/10.1080/09397140.2010.10638456)
- Dumnicka, E. 2004. A description of *Cernosvitoviella tridentina*, a new species of Enchytraeidae (Oligochaeta) from the Italian Alps. *Annales de Limnologie*, 40(2), 133-137. doi: [10.1051/limn/2004011](https://doi.org/10.1051/limn/2004011)
- Dumnicka, E. 1998. A new species of fridericia (Oligochaeta: Enchytraeidae) found in a spring in the Sudety Mountains (Poland). *Annales de Limnologie*, 34(2), 155-158. doi: [10.1051/limn/1998014](https://doi.org/10.1051/limn/1998014)
- Dumnicka, E. 1994. Communities of oligochaetes in mountain streams of Poland. *Hydrobiologia*, 278(1-3), 107-110. doi: [10.1007/BF00142317](https://doi.org/10.1007/BF00142317)
- Dumnicka, E. 2006. Composition and abundance of oligochaetes (Annelida: Oligochaeta) in springs of Kraków-Częstochowa Upland (southern Poland): Effect of spring encasing and environmental factors. *Polish Journal of Ecology*, 54(2), 231-242. Retrieved from: www.scopus.com
- Dumnicka, E. 1986. Ecology of some waters in the forest-agricultural basin of the River Brynica near the upper Silesian industrial region. 9. Communities of oligochaetes. *Acta Hydrobiologica*, 27(4), 535-545. Retrieved from: www.scopus.com
- Dumnicka, E. 1994. Habitat preferences of invertebrates (especially Oligochaeta) in a stream. *Acta Hydrobiologica*, 36(1), 91-101. Retrieved from: www.scopus.com
- Dumnicka, E. 2009. New for Poland tubificid (Oligochaeta) species from karstic springs. *Polish Journal of Ecology*, 57(2), 395-401. Retrieved from: www.scopus.com
- Dumnicka, E. 2014. Stygobitic oligochaetes (Annelida, Clitellata) in Poland with remarks on their distribution in central Europe. *Subterranean Biology*, 14(1), 15-24. doi: [10.3897/subbtol.14.7700](https://doi.org/10.3897/subbtol.14.7700)
- Dumnicka, E. 1987. The effect of dam reservoirs on oligochaete communities in the River Dunajec (southern Poland). *Acta Hydrobiologica*, 29(1), 25-34. Retrieved from: www.scopus.com
- Dumnicka, E. 2010. Two new freshwater enchytraeid species (Oligochaeta) from the Italian Alps. *Italian Journal of Zoology*, 77(1), 38-43. doi: [10.1080/1125000902855505](https://doi.org/10.1080/1125000902855505)

- Dumnicka, E. 2002. Upper Vistula River: Response of aquatic communities to pollution and impoundment. X. oligochaete taxocens. *Polish Journal of Ecology*, 50(2), 237-247. Retrieved from: www.scopus.com
- Dumnicka, E. & Boggero, A. 2007. Freshwater Oligochaeta in two mountain ranges in europe: The tatra mountains (Poland) and the alps (Italy). *Fundamental and Applied Limnology*, 168(3), 231-242. doi: [10.1127/1863-9135/2007/0168-0231](https://doi.org/10.1127/1863-9135/2007/0168-0231)
- Dumnicka, E. & Galas, J. 2002. Factors affecting the distribution of Oligochaeta in small high mountain ponds (Tatra Mountains, Poland). *Archiv Fur Hydrobiologie*, 156(1), 121-133. doi: [10.1127/0003-9136/2002/0156-0121](https://doi.org/10.1127/0003-9136/2002/0156-0121)
- Dumnicka, E. & Galas, J. 2012. Temporal changes in oligochaete fauna of three alpine ponds in the Tatra Mountains (Poland). *Boreal Environment Research*, 17(3-4), 252-262. Retrieved from: www.scopus.com
- Dumnicka, E. & Koszałka, J. 2005. The effect of drought on Oligochaeta communities in small woodland streams. *Biologia – Section Zoology*, 60(2), 143-150. Retrieved from: www.scopus.com
- Dumnicka, E. & Krodkiewska, M. 2003. Studies on freshwater Oligochaeta in the upper Silesia region (southern Poland). *Biologia – Section Zoology*, 58(5), 897-902. Retrieved from: www.scopus.com
- Dumnicka, E. & Poznańska, M. 2006. Novel polish recordings of rare aquatic Oligochaeta species. *Oceanological and Hydrobiological Studies*, 35(2), 111-120. Retrieved from: www.scopus.com
- Dumnicka, E., Steingruber, S., Colombo, L., Zaupa, S. & Boggero, A. 2015. Oligochaete assemblages of swiss alpine lakes. *Italian Journal of Zoology*, 82(1), 112-123. doi: [10.1080/11250003.2014.965230](https://doi.org/10.1080/11250003.2014.965230)
- El-Mansy, A., Molnár, K. & Székely, C. 1998. Development of Myxobolus portugalensis saraiva and molnar, 1990 (Myxosporea: Myxobolidae) in the oligochaete *Tubifex tubifex* (Müller). *Systematic Parasitology*, 41(2), 95-103. doi: [10.1023/A:1006034527844](https://doi.org/10.1023/A:1006034527844)
- El-Matbouli, M., McDowell, T.S., Antonio, D.B., Andree, K.B. & Hedrick, R.P. 1999. Effect of water temperature on the development, release and survival of the triactinomyxon stage of *Myxobolus cerebralis* in its oligochaete host. *International Journal for Parasitology*, 29(4), 627-641. doi: [10.1016/S0020-7519\(99\)00009-0](https://doi.org/10.1016/S0020-7519(99)00009-0)
- Erickson, R.J., Ankley, G.T., Defoe, D.L., Kosian, P.A. & Makynen, E.A. 1999. Additive toxicity of binary mixtures of phototoxic polycyclic aromatic hydrocarbons to the oligochaete *Lumbriculus variegatus*. *Toxicology and Applied Pharmacology*, 154(1), 97-105. doi: [10.1006/taap.1998.8563](https://doi.org/10.1006/taap.1998.8563)
- Erséus, C. 1990. Cladistic analysis of the subfamilies within the Tubificidae (Oligochaeta). *Zoologica Scripta*, 19(1), 57-63. doi: [10.1111/j.1463-6409.1990.tb00240.x](https://doi.org/10.1111/j.1463-6409.1990.tb00240.x)
- Erséus, C. 1987. Phylogenetic analysis of the aquatic Oligochaeta under the principle of parsimony. *Hydrobiologia*, 155(1), 75-89. doi: [10.1007/BF00025633](https://doi.org/10.1007/BF00025633)
- Erséus, C. 2005. Phylogeny of oligochaetous Clitellata. *Hydrobiologia*, 535(1), 357-372. doi: [10.1007/s10750-004-4426-x](https://doi.org/10.1007/s10750-004-4426-x)
- Erséus, C., Envall, I., Marchese, M. & Gustavsson, L. 2010. The systematic position of opistocystidae (Annelida, Clitellata) revealed by DNA data. *Molecular Phylogenetics and Evolution*, 54(1), 309-313. doi: [10.1016/j.ympev.2009.09.037](https://doi.org/10.1016/j.ympev.2009.09.037)
- Erséus, C. & Grimm, R. 2002. A new species of ainudrilus (Tubificidae) from south Georgia, and other subantarctic freshwater oligochaetes. *Hydrobiologia*, 468, 77-81. doi: [10.1023/A:1015249024244](https://doi.org/10.1023/A:1015249024244)
- Erséus, C. & Källersjö, M. 2004. 18S rDNA phylogeny of Clitellata (Annelida). *Zoologica Scripta*, 33(2), 187-196. doi: [10.1111/j.1463-6409.2004.00146.x](https://doi.org/10.1111/j.1463-6409.2004.00146.x)
- Erséus, C. & Paoletti, A. 1986. An italian record of the aquatic oligochaete *Monopylephorus limosus* (Tubificidae), previously known only from Japan and China. *Bollettino di Zoologia*, 53(1), 115-118. doi: [10.1080/11250008609355492](https://doi.org/10.1080/11250008609355492)
- Erséus, C., Rota, E., Matamoros, L. & De Wit, P. 2010. Molecular phylogeny of Enchytraeidae (Annelida, Clitellata). *Molecular Phylogenetics and Evolution*, 57(2), 849-858. doi: [10.1016/j.ympev.2010.07.005](https://doi.org/10.1016/j.ympev.2010.07.005)
- Erséus, C., Rota, E., Timm, T., Grimm, R., Healy, B. & Lundberg, S. 2005. Riverine and riparian clitellates of three drainages in southern Sweden. *Annales de Limnologie*, 41(3), 183-194. doi: [10.1051/limn:20054130183](https://doi.org/10.1051/limn:20054130183)
- Erséus, C. & Sang, Q. 1985. Two aberrant Tubificidae (Oligochaeta) from Pearl River in the people's Republic of China. *Hydrobiologia*, 127(3), 193-196. doi: [10.1007/BF00000001](https://doi.org/10.1007/BF00000001)
- Eszterbauer, E., Székely, C., Molnár, K. & Baska, F. 2000. Development of *Myxobolus bramae* (Myxosporea: Myxobolidae) in an oligochaete alternate host, *Tubifex tubifex*. *Journal of Fish Diseases*, 23(1), 19-25. doi: [10.1046/j.1365-2761.2000.00202.x](https://doi.org/10.1046/j.1365-2761.2000.00202.x)
- Ewald, G., Berglund, O. & Svensson, J.M. 1997. Effect of oligochaete bioturbation on sediment accumulation of 2,2',4,4'-tetrachlorobiphenyl. *Ecotoxicology and Environmental Safety*, 36(1), 66-71. doi: [10.1006/eesa.1996.1489](https://doi.org/10.1006/eesa.1996.1489)
- Falci Theza Rodrigues, L., Jabour Vescovi Rosa, B.F., Lobo, H., Campos Divino, A. & Da Gama Alves, R. 2015. Diversity and distribution of oligochaetes in tropical forested streams, southeastern Brazil. *Journal of Limnology*, 74(3), 433-443. doi: [10.4081/jlimnol.2015.1024](https://doi.org/10.4081/jlimnol.2015.1024)
- Famme, P. & Knudsen, J. 1985. Aerotaxis by the freshwater oligochaete *Tubifex* sp. *Oecologia*, 65(4), 599-601. doi: [10.1007/BF00379679](https://doi.org/10.1007/BF00379679)
- Fend, S.V., Rodriguez, P. & Lenat, D.R. 2015. *Uktena riparia* gen. nov., sp. nov. (Annelida, Clitellata, Lumbriculidae), a new spermatophore-producing oligochaete. *Zootaxa*, 3994(3), 411-424. doi: [10.11646/zootaxa.3994.3.5](https://doi.org/10.11646/zootaxa.3994.3.5)
- Ferraguti, M. 1985. Slanted centriole and transient anchoring apparatus during spermiogenesis of an oligochaete (Annelida). *Biology of the Cell*, 52(2), 175-180. doi: [10.1111/j.1768-322X.1985.tb00334.x](https://doi.org/10.1111/j.1768-322X.1985.tb00334.x)
- Ferraguti, M., Bernardini, G., Melone, G. & Dallai, R. 1988. Structure and function of the metachronal wave in *Tubifex* *Tubifex* spermatozeugmata (Annelida, Oligochaeta). *Journal of Ultrastructure Research and Molecular Structure Research*, 99(1), 79-95. doi: [10.1016/0889-1605\(88\)90035-3](https://doi.org/10.1016/0889-1605(88)90035-3)
- Ferraguti, M. & Erséus, C. 1999. Sperm types and their use for a phylogenetic analysis of aquatic clitellates. *Hydrobiologia*, 402, 225-237. doi: [10.1023/A:1003752811830](https://doi.org/10.1023/A:1003752811830)
- Ferraguti, M., Erséus, C., Kaygorodova, I. & Martin, P. 1999. New sperm types in Naididae and Lumbriculidae (Annelida: Oligochaeta) and their possible phylogenetic implications. *Hydrobiologia*, 406, 213-222. doi: [10.1023/A:1003708906089](https://doi.org/10.1023/A:1003708906089)
- Ferraguti, M., Erséus, C. & Pinder, A. 1996. The spermatozoon of capilloventer australis and the systematic position of the capilloventridae (Annelida: Oligochaeta). *Australian Journal of Zoology*, 44(5), 469-478. doi: [10.1071/ZO9960469](https://doi.org/10.1071/ZO9960469)
- Ferraguti, M., Fascio, U. & Boi, S. 2002. Mass production of basal bodies in paraspermiogenesis of *Tubificinae* (Annelida, Oligochaeta). *Biology of the Cell*, 94(2), 109-115. doi: [10.1016/S0248-4900\(02\)01185-1](https://doi.org/10.1016/S0248-4900(02)01185-1)
- Finder, A.M., Hill, B.D. & Green, P.E. 1998. A new species of *Dero* (Allodero) (Naididae: Oligochaeta) inhabiting the wolffian ducts of the green tree frog (*Litoria caerulea*) in Queensland. *Memoirs of the Queensland Museum*, 42(2), 559-564. Retrieved from: www.scopus.com
- Finogenova, N.P. & Lobasheva, T.M. 1987. Growth of *Tubifex* *tubifex* Müller (Oligochaeta, Tubificidae) under various trophic conditions. *Internationale Revue Der Gesamten Hydrobiologie Und Hydrographie*, 72(6), 709-726. doi: [10.1002/iroh.19870720608](https://doi.org/10.1002/iroh.19870720608)
- Fisher, S.W., Chordas III, S.W. & Landrum, P.F. 1999. Lethal and sublethal body residues for PCB intoxication in the oligochaete, *Lumbriculus*

- variegatus. *Aquatic Toxicology*, 45(2-3), 115-126. doi: [10.1016/S0166-445X\(98\)00103-9](https://doi.org/10.1016/S0166-445X(98)00103-9)
- Fisk, A.T., Wiens, S.C., Barrie Webster, G.R., Bergman, Å. & Muir, D.C.G. 1998. Accumulation and depuration of sediment-sorbed C12- and C16-polychlorinated alkanes by oligochaetes (*Lumbriculus variegatus*). *Environmental Toxicology and Chemistry*, 17(10), 2019-2026. doi: [10.1897/1551-5028\(1998\)017<2019:AADSO>2.3.CO;2](https://doi.org/10.1897/1551-5028(1998)017<2019:AADSO>2.3.CO;2)
- Flores-Tena, F.J. & Martínez-Tabche, L. 2001. The effect of chromium on the hemoglobin concentration of *Limnodrilus hoffmeisteri* (Oligochaeta: Tubificidae). *Ecotoxicology and Environmental Safety*, 50(3), 196-202. doi: [10.1006/eesa.2001.2104](https://doi.org/10.1006/eesa.2001.2104)
- Florou, H., Tsytsgina, V., Polikarpov, G.G., Vosniakos, F.K. & Evangelou, N. 2009. A comparative study on genotoxic effects in Oligochaeta from areas with elevated natural and artificial radioactivity. *Journal of Environmental Protection and Ecology*, 10(3), 807-814. Retrieved from: www.scopus.com
- Gabric, A.J., Connell, D.W. & Bell, P.R.F. 1990. A kinetic model for bioconcentration of lipophilic compounds by oligochaetes. *Water Research*, 24(10), 1225-1231. doi: [10.1016/0043-1354\(90\)90045-8](https://doi.org/10.1016/0043-1354(90)90045-8)
- Gainutdinov, M.K., Kalinnikova, T.B., Gainutdinov, T.M. & Yargunov, V.G. 2002. Self-regulation of Oligochaetan worm organisms by their chemical signals effused by them. *Journal of Evolutionary Biochemistry and Physiology*, 38(3), 364-366. doi: [10.1023/A:1020701130402](https://doi.org/10.1023/A:1020701130402)
- Gelder, S.R. 1989. Histophysiology of digestion and observations on the structure of the alimentary canal in the ectosymbiont *Chaetogaster limnaei limnaei* Baer, 1827 (Annelida: Oligochaeta). *Hydrobiologia*, 180(1), 115-125. doi: [10.1007/BF00027544](https://doi.org/10.1007/BF00027544)
- Gerhardt, A. 2007. Importance of exposure route for behavioural responses in *Lumbriculus variegatus* Müller (Oligochaeta: Lumbriculida) in short-term exposures to PB. *Environmental Science & Pollution Research*, 14(6), 430-434. doi: [10.1065/espr2006.12.371](https://doi.org/10.1065/espr2006.12.371)
- Gerhardt, A. 2009. Screening the toxicity of NI, CD, CU, ivermectin, and imidacloprid in a short-term automated behavioral toxicity test with *Tubifex tubifex* (Müller, 1774) (Oligochaeta). *Human and Ecological Risk Assessment*, 15(1), 27-40. doi: [10.1080/10807030802614983](https://doi.org/10.1080/10807030802614983)
- Giani, N., Martin, P. & Juget, J. 1995. A new species of Phreodrilidae (Oligochaeta), *Astacopsisdrilus naceri* sp. nov., from Morocco (north africa), with notes on the biogeography of the family. *Canadian Journal of Zoology*, 73(12), 2375-2381. doi: [10.1139/z95-277](https://doi.org/10.1139/z95-277)
- GIANI, N. & RODRIGUEZ, P. 1994. New species of the genus trichodrilus (Oligochaeta, Lumbriculidae). *Zoologica Scripta*, 23(1), 33-41. doi: [10.1111/j.1463-6409.1994.tb00371.x](https://doi.org/10.1111/j.1463-6409.1994.tb00371.x)
- Giani, N., Sambugar, B., Martínez-Ansemil, E., Martin, P. & Schmelz, R.M. 2011. The groundwater oligochaetes (Annelida, Clitellata) of Slovenia. *Subterranean Biology*, 9(1), 85-102. doi: [10.3897/subbtbiol.9.2512](https://doi.org/10.3897/subbtbiol.9.2512)
- Gilbert, M.A. & Granath Jr., W.O. 2008. Susceptibility of *Tubifex tubifex* (Annelida: Oligochaeta: Tubificidae) from the rock creek drainage of west central Montana, USA, to *Myxobolus cerebralis* (Myxozoa: Myxosporea: Myxobolidae), the causative agent of salmonid whirling disease. *Comparative Parasitology*, 75(1), 92-97. doi: [10.1654/4310.1](https://doi.org/10.1654/4310.1)
- Gillis, P.L., Diener, L.C., Reynoldson, T.B. & Dixon, D.G. 2002. Cadmium-induced production of a metallothioneinlike protein in *Tubifex tubifex* (Oligochaeta) and *Chironomus riparius* (Diptera): Correlation with reproduction and growth. *Environmental Toxicology and Chemistry*, 21(9), 1836-1844. Retrieved from: www.scopus.com
- Gillis, P.L., Reynoldson, T.B. & Dixon, D.G. 2006. Metallothionein-like protein and tissue metal concentrations in invertebrates (Oligochaetes and Chironomids) collected from reference and metal contaminated field sediments. *Journal of Great Lakes Research*, 32(3), 565-577. doi: [10.3394/0380-1330\(2006\)32\[565:MPATMC\]2.0.CO;2](https://doi.org/10.3394/0380-1330(2006)32[565:MPATMC]2.0.CO;2)
- Gluzman, C. 1990. A new species of the freshwater genus allonais (Oligochaeta, Naididae) from among the roots of *Eichhornia crassipes* (Argentina). *Studies on Neotropical Fauna and Environment*, 25(3), 121-124. doi: [10.1080/01650529009360811](https://doi.org/10.1080/01650529009360811)
- Gluzman, C. 1997. Sperm cells in *Aeolosoma marcusii* (Annelida, Oligochaeta). *Biocell*, 21(2), 137-142. Retrieved from: www.scopus.com
- Gluzman, C. 1998. Ultrastructural study of the spermiogenesis in two species of Naididae, *paranaensis* litoralis and *Nais communis* (Annelida, Oligochaeta). *Biocell*, 22(2), 141-148. Retrieved from: www.scopus.com
- Gluzmán, C. 1999. The spermatozoon of *Slavina appendiculata* (Annelida, Oligochaeta). *Biocell*, 23(2), 97-102. Retrieved from: www.scopus.com
- Gnaiger, E., Kaufmann, R. & Staudigl, I. 1987. Physiological reactions of aquatic oligochaetes to environmental anoxia. *Hydrobiologia*, 155(1), 155. doi: [10.1007/BF00025641](https://doi.org/10.1007/BF00025641)
- Goretti, E., Marcucci, C., Di Veroli, A., Fabrizi, A. & Gaino, E. 2014. The Tubificids (Annelida, Oligochaeta) of Lake Trasimeno and Lake Piediluco in central Italy, with a study of SEM morphology of some species. *Turkish Journal of Zoology*, 38(3), 334-341. doi: [10.3906/zoo-1308-40](https://doi.org/10.3906/zoo-1308-40)
- Gorni, G., Peiro, D.F. & Sanches, N. 2015. Aquatic Oligochaeta (Annelida: Clitellata) from state of São Paulo, Brazil: Diversity and occurrence review. [Oligochaeta aquática (Annelida: Clitellata) do estado de São Paulo: revisão da diversidade e ocorrência]. *Biota Neotropica*, 15(1), 1-8. doi: [10.1590/1676-06032015006314](https://doi.org/10.1590/1676-06032015006314)
- Gorni, G.R. & Alves, R.D.G. 2007. Naididae (Annelida, Oligochaeta) associated with briophytes in Brotas, state of São Paulo, Brazil. *Revista Brasileira de Zoologia*, 24(2), 518-519. doi: [10.1590/S0101-81752007000200036](https://doi.org/10.1590/S0101-81752007000200036)
- Gorni, G.R. & Alves, R.D.G. 2006. Naididae (Annelida, Oligochaeta) associated with *Pomacea bridgesii* (Reeve) (Gastropoda, Ampullariidae). *Revista Brasileira de Zoologia*, 23(4), 1059-1061. doi: [10.1590/S0101-81752006000400011](https://doi.org/10.1590/S0101-81752006000400011)
- Gorni, G.R. & Alves, R.G. 2008. Oligochaeta (Annelida: Clitellata) in headwater streams of the Parque Estadual de Campos do Jordão (São Paulo, Brazil). [Oligochaeta (Annelida: Clitellata) em córregos de baixa ordem do Parque Estadual de Campos do Jordão (São Paulo, Brasil)]. *Biota Neotropica*, 8(4), 161-165. Retrieved from: www.scopus.com
- Gorni, G.R. & Alves, R.G. 2012. Oligochaetes (Annelida, Clitellata) in a neotropical stream: A mesohabitat approach. [Oligoquetos (Annelida, Clitellata) em um córrego neotropical: Uma abordagem do mesohabitat]. *Iheringia – Série Zoologia*, 102(1), 106-110. doi: [10.1590/S0073-47212012000100015](https://doi.org/10.1590/S0073-47212012000100015)
- Grabowski, M. & Jabłońska, A. 2009. First records of Branchiura sowerbyi Beddard, 1892 (Oligochaeta: Tubificidae) in greece. *Aquatic Invasions*, 4(2), 365-367. doi: [10.3391/ai.2009.4.2.10](https://doi.org/10.3391/ai.2009.4.2.10)
- Granath Jr., W.O. & Gilbert, M.A. 2002. The role of *Tubifex tubifex* (Annelida: Oligochaeta: Tubificidae) in the transmission of *Myxobolus cerebralis* (Myxozoa: Myxosporea: Myxobolidae). *American Fisheries Society Symposium*, 2002(29), 79-85. Retrieved from: www.scopus.com
- Gregory, T.R. & Hebert, P.D.N. 2002. Genome size estimates for some oligochaete annelids. *Canadian Journal of Zoology*, 80(8), 1485-1489. doi: [10.1139/Z02-145](https://doi.org/10.1139/Z02-145)
- Grimm, R. 1987. Contributions towards the taxonomy of the african Naididae (Oligochaeta). IV. Zoogeographical and taxonomical considerations on african Naididae. *Hydrobiologia*, 155(1), 27-37. doi: [10.1007/BF00025628](https://doi.org/10.1007/BF00025628)
- Groh, S., Claus, E., Manz, W., Möhlenkamp, C., Immermann, R.-. & Blübaum-Gronau, E. 2010. Influence of increasing sediment organic matter on reproduction and growth of the benthic oligochaete *Lumbriculus variegatus*. [Einfluss organischer substanz auf die reproduktion, biomasse und akkumulation hydrophober schadstoffe in *Lumbriculus variegatus*]. *Umweltwissenschaften Und Schadstoff-Forschung*, 22(2), 99-106. doi: [10.1007/s12302-010-0115-8](https://doi.org/10.1007/s12302-010-0115-8)

- Guérin, C. & Labroue, L. 1991. A laboratory study of the effect of *Tubifex tubifex* Müll. (Tubificidae, Oligochaeta) on the release of phosphorus through the sediment. [Etude expérimentale de l'influence de *Tubifex tubifex* Müll. (Tubificidae, Oligochaeta) sur le relargage du phosphore par le sédiment]. *Annales de Limnologie*, 27(3), 253-265. doi: [10.1051/limn/1991019](https://doi.org/10.1051/limn/1991019)
- Gupta, R.K., Kumar, S., Gorai, A.C. & Pandey, P.N. 1998. Oligochaetes in a freshwater pond in dhanbad. *Ecology, Environment and Conservation*, 4(3), 123-126. Retrieved from: www.scopus.com
- Gust, K.A. & Fleeger, J.W. 2006. Exposure to cadmium-phenanthrene mixtures elicits complex toxic responses in the freshwater tubificid oligochaete, *Ilyodrilus templetoni*. *Archives of Environmental Contamination and Toxicology*, 51(1), 54-60. doi: [10.1007/s00244-005-1075-7](https://doi.org/10.1007/s00244-005-1075-7)
- Gustavsson, L.M. 2001. Comparative study of the cuticle in some aquatic oligochaetes (Annelida: Clitellata). *Journal of Morphology*, 248(2), 185-195. doi: [10.1002/jmor.1029](https://doi.org/10.1002/jmor.1029)
- Gustavsson, L.M. & Erséus, C. 1997. Morphogenesis of the genital ducts and spermathecae in *Clitellio arenarius*, *Heterochaeta costata*, *Tubificoides benedii* (Tubificidae) and *Stylaria lacustris* (Naididae) (Annelida, Oligochaeta). *Acta Zoologica*, 78(1), 9-31. doi: [10.1111/j.1463-6395.1997.tb01122.x](https://doi.org/10.1111/j.1463-6395.1997.tb01122.x)
- Gustavsson, L.M. & Erséus, C. 1999. Morphology and phylogenetic implications of oesophageal modifications in the limnodriloidinae (Oligochaeta, Tubificidae). *Journal of Zoology*, 248(4), 467-482. doi: [10.1017/S0952836999008067](https://doi.org/10.1017/S0952836999008067)
- Hallett, S.L., Atkinson, S.D. & Bartholomew, J.L. 2005. Countering morphological ambiguities: Development of a PCR assay to assist the identification of *Tubifex tubifex* oligochaetes. *Hydrobiologia*, 543(1), 305-309. doi: [10.1007/s10750-004-6954-9](https://doi.org/10.1007/s10750-004-6954-9)
- Hartwich, G. & Kilias, I. 1989. Die Oligochaeten-Typen des zoologischen museums in berlin. *Mitteilungen Aus Dem Museum Für Naturkunde in Berlin – Zoologisches Museum Und Institut Für Spezielle Zoologie (Berlin)*, 65(2), 249-295. doi: [10.1002/mmzn.19890650210](https://doi.org/10.1002/mmzn.19890650210)
- He, X., Cui, Y. & Wang, H. 2010. *Limnodrilus simplex* sp. nov. (Oligochaeta: Naididae: Tubificinae) from Changjiang River, China. *Zoological Science*, 27(9), 768-770. doi: [10.2108/zsj.27.768](https://doi.org/10.2108/zsj.27.768)
- Healy, B. 1987. The depth distribution of Oligochaeta in an Irish quaking marsh. *Hydrobiologia*, 155(1), 235-247. doi: [10.1007/BF00025656](https://doi.org/10.1007/BF00025656)
- Healy, B. & Coates, K.A. 1999. Finding enchytraeid oligochaetes (Clitellata) in hot climates: Species occurrence on the shores of Bermuda. *Hydrobiologia*, 406, 111-117. doi: [10.1023/A:1003760922928](https://doi.org/10.1023/A:1003760922928)
- Healy, B. & Fend, S. 2002. The occurrence of *mesenchytraeus* (Enchytraeidae: Oligochaeta) in riffle habitats of north-west american rivers, with description of a new species. *Journal of Natural History*, 36(1), 15-23. doi: [10.1080/713833842](https://doi.org/10.1080/713833842)
- Hegde, P.R. & Sreepada, K.S. 2015. Freshwater oligochaetes (Annelida) from western ghats and the west Coast of Karnataka (India). *Turkish Journal of Zoology*, 39(3), 523-526. doi: [10.3906/zoo-1311-46](https://doi.org/10.3906/zoo-1311-46)
- Higgins, C.P., Mcleod, P.B., Macmanus-Spencer, L.A. & Luthy, R.G. 2007. Bioaccumulation of perfluorochemicals in sediments by the aquatic oligochaete *Lumbriculus variegatus*. *Environmental Science & Technology*, 41(13), 4600-4606. doi: [10.1021/es0627920](https://doi.org/10.1021/es0627920)
- Hipp, E., Bickel, U., Mustafa, T. & Hoffmann, K.H. 1986. Integumentary uptake of acetate and propionate (VFA) by *Tubifex* sp., a freshwater oligochaete. II. Role of VFA as nutritional resources and effects of anaerobiosis. *Journal of Experimental Zoology*, 240(3), 299-308. doi: [10.1002/jez.1402400303](https://doi.org/10.1002/jez.1402400303)
- Hipp, E., Mustafa, T., Bickel, U. & Hoffmann, K.H. 1986. Integumentary uptake of acetate and propionate (VFA) by *Tubifex* sp., a freshwater oligochaete. I. Uptake rates and transport kinetics. *Journal of Experimental Zoology*, 240(3), 289-297. doi: [10.1002/jez.1402400302](https://doi.org/10.1002/jez.1402400302)
- Hipp, E., Mustafa, T. & Hoffmann, K.H. 1985. Integumentary uptake of volatile fatty acids by the freshwater oligochaete *Tubifex*. *Naturwissenschaften*, 72(3), 148-149. doi: [10.1007/BF00490402](https://doi.org/10.1007/BF00490402)
- Hirabayashi, K., Oga, K., Yoshizawa, K., Yoshida, N. & Kazama, F. 2012. A long-term eutrophication process observed from the changes in the horizontal distribution of profundal oligochaete fauna in mesotrophic-eutrophic Lake Kawaguchi, Japan. *Turkish Journal of Zoology*, 36(1), 39-46. doi: [10.3906/zoo-0912-18](https://doi.org/10.3906/zoo-0912-18)
- Hoffmann, H., Hipp, E. & Sedlmeier, U.A. 1987. Aerobic and anaerobic metabolism of the freshwater oligochaete *Tubifex* sp. *Hydrobiologia*, 155(1), 157-158. doi: [10.1007/BF00025642](https://doi.org/10.1007/BF00025642)
- Hoffmann, K.H. & Wulf, A. 1993. Comparative studies of integumentary uptake of short-chain carboxylic acids by freshwater oligochaetes. *Comparative Biochemistry and Physiology – Part A: Physiology*, 104(1), 169-174. doi: [10.1016/0300-9629\(93\)90025-Y](https://doi.org/10.1016/0300-9629(93)90025-Y)
- Huang, T.-., Nie, X.-., Zhang, J.-., Li, W., Li, X.-., Zhang, S. & Wang, B.-. 2010. Study on migration modes and correlative factors of *Tubifex* tubifex (Oligochaeta, Tubificidae) in source water. *Water Science and Technology: Water Supply*, 10(6), 887-895. doi: [10.2166/ws.2010.503](https://doi.org/10.2166/ws.2010.503)
- Hyötyläinen, T. & Oikari, A. 2004. Bioaccumulation of PAHs from creosote-contaminated sediment in a laboratory-exposed freshwater oligochaete, *Lumbriculus variegatus*. *Chemosphere*, 57(2), 159-164. doi: [10.1016/j.chemosphere.2004.05.001](https://doi.org/10.1016/j.chemosphere.2004.05.001)
- Ibrahim, M.M. 2007. Population dynamics of *Chaetogaster limnaei* (Oligochaeta: Naididae) in the field populations of freshwater snails and its implications as a potential regulator of trematode larvae community. *Parasitology Research*, 101(1), 25-33. doi: [10.1007/s00436-006-0436-0](https://doi.org/10.1007/s00436-006-0436-0)
- Ilyashuk, B.P. 1999. Littoral oligochaete (Annelida: Oligochaeta) communities in neutral and acidic lakes in the Republic of Karelia, Russia. *Boreal Environment Research*, 4(3), 277-284. Retrieved from: www.scopus.com
- Irmler, U. 1989. Population-ecology and migration of *Dero multibranchiata* stieren, 1892 (Naididae, Oligochaeta) in the central amazon inundation forest. *Amazoniana*, 11(1), 31-52. Retrieved from: www.scopus.com
- Jabłońska, A. 2014. Oligochaete communities of highly degraded urban streams in Poland, central europe. *North-Western Journal of Zoology*, 10(1), 74-82. Retrieved from: www.scopus.com
- Jabłońska, A. & Pešić, V. 2011. Five aquatic Oligochaeta species new for the fauna of Montenegro. *Turkish Journal of Zoology*, 35(1), 119-121. doi: [10.3906/zoo-0903-12](https://doi.org/10.3906/zoo-0903-12)
- Jabłońska, A. & Pešić, V. 2014. Five species of aquatic oligochaetes new to Iran with an updated checklist. *Oceanological and Hydrobiological Studies*, 43(1), 100-105. doi: [10.2478/s13545-014-0121-3](https://doi.org/10.2478/s13545-014-0121-3)
- Jabłońska, A., Szlauer-Łukaszewska, A. & Bańkowska, A. 2015. *Aulodrilus pigueti* Kowalewski, 1914 (Annelida: Clitellata) – A new record for the polish fauna from the oder river and remarks on other oligochaetes rarely noticed in Poland. *Oceanological and Hydrobiological Studies*, 44(4), 456-465. doi: [10.1515/ohs-2015-0043](https://doi.org/10.1515/ohs-2015-0043)
- Jamieson, B.G.M. 1988. On the phylogeny and higher classification of the Oligochaeta. *Cladistics*, 4(4), 367-401. doi: [10.1111/j.1096-0031.1988.tb00520.x](https://doi.org/10.1111/j.1096-0031.1988.tb00520.x)
- Jamieson, B.G.M., Erseus, C. & Ferraguti, M. 1987. Parsimony analysis of the phylogeny of some Oligochaeta (Annelida) using spermatozoal ultrastructure. *Cladistics*, 3(2), 145-155. doi: [10.1111/j.1096-0031.1987.tb00503.x](https://doi.org/10.1111/j.1096-0031.1987.tb00503.x)
- Jenderedjian, K. 1996. Energy budget of Oligochaeta and its relationship with the primary production of Lake Sevan, Armenia. *Hydrobiologia*, 334(1-3), 133-140. doi: [10.1007/BF00017362](https://doi.org/10.1007/BF00017362)
- Jiang, T., Du, S., Sun, P. & Zhu, M. 2012. Effect of oligochaete worm body fluids on biological phosphorus removal in a bench-scale EBPR system.

- Environmental Technology (United Kingdom)*, 33(15), 1755-1762. doi: [10.1080/09593330.2011.644866](https://doi.org/10.1080/09593330.2011.644866)
- Johnson, P. & Ladle, M. 1989. The Enchytraeidae (Oligochaeta) of streams of southern England. [Les Enchytraeidae (Oligochaeta) des eaux courantes du sud de l'Angleterre]. *Annales de Limnologie*, 25(2), 121-129. doi: [10.1051/limn/1989012](https://doi.org/10.1051/limn/1989012)
- Juget, J. 1987. Contribution to the study of the rhyacodrilinae (Tubificidae, Oligochaeta), with description of two new stygobiont species from the alluvial plain of the french upper rhone, *Rhyacodrilus amphigenus*, sp. nov., and *Rhizodriloides phreaticola*, gen. nov., sp. nov. *Hydrobiologia*, 155(1), 107-118. doi: [10.1007/BF00025635](https://doi.org/10.1007/BF00025635)
- Juget, J., Goubier, V. & Barthélémy, D. 1989. Intrinsic and extrinsic variables controlling the productivity of asexual populations of *Nais* sp., (Naididae, Oligochaeta). *Hydrobiologia*, 180(1), 177-184. doi: [10.1007/BF00027550](https://doi.org/10.1007/BF00027550)
- Juget, J. & Lafont, M. 1994. Distribution of Oligochaeta in some lakes and pools of Bolivia. *Hydrobiologia*, 278(1-3), 125-127. doi: [10.1007/BF00142319](https://doi.org/10.1007/BF00142319)
- Juget, J. & Lafont, M. 1994. Theoretical habitat templets, species traits, and species richness: Aquatic oligochaetes in the upper rhône river and its floodplain. *Freshwater Biology*, 31(3), 327-340. doi: [10.1111/j.1365-2427.1994.tb01744.x](https://doi.org/10.1111/j.1365-2427.1994.tb01744.x)
- Juget, J., Riera, P. & Schmelz, R. 1994. A study of the trophic relationship between biotecton and Naididae (*Nais* sp., Oligochaeta): Importance of organic carbon. [Etude des relations trophiques entre le biotecton et les Naididae (*Nais* sp., Oligochaeta): Importance du carbone organique (C.O.T.)]. *Annales de Limnologie*, 30(4), 253-265. doi: [10.1051/limn/1994019](https://doi.org/10.1051/limn/1994019)
- Juget, J. & Yacoubi-Khebiza, M. 1997. Research on the ecology of *Astacopsisdrilus naceri giani* and *martin*, 1995 (Phreodrilidae, Oligochaeta) from the groundwaters of Morocco. [Contribution à l'écologie de l'espèce *Astacopsisdrilus naceri Giani* and *Martin*, 1995 (Phreodrilidae, Oligochaeta) en provenance des eaux souterraines du Maroc]. *Annales de Limnologie*, 33(3), 149-161. Retrieved from: www.scopus.com
- Kaeser, A.J. & Sharpe, W.E. 2006. Patterns of distribution and abundance of *Tubifex tubifex* and other aquatic oligochaetes in *Myxobolus cerebralis* enzootic areas in pennsylvania. *Journal of Aquatic Animal Health*, 18(1), 64-78. doi: [10.1577/H05-016.1](https://doi.org/10.1577/H05-016.1)
- Kahl, K. & Golanska, M. 1987. Diversity and numbers of Oligochaeta against the background of other macroinvertebrates in a concrete bed of the River Widawka (central Poland). *Acta Hydrobiologica*, 29(1), 35-51. Retrieved from: www.scopus.com
- Kaigorodova, I.A. & Shcherbakov, D.I. 2006. Molecular phylogenetic study of the systematic position of Baikalian oligochaetes in the system Clitellata. *Genetika*, 42(12), 1647-1655. Retrieved from: www.scopus.com
- Kairesalo, T. & Koskimies, I. 1987. Grazing by oligochaetes and snails on epiphytes. *Freshwater Biology*, 17(2), 317-324. doi: [10.1111/j.1365-2427.1987.tb01051.x](https://doi.org/10.1111/j.1365-2427.1987.tb01051.x)
- Kaliszewicz, A. 2015. Conspecific alarm cues induce an alternative reproductive strategy in aquatic oligochaetes. *Limnology*, 16(2), 85-90. doi: [10.1007/s10201-014-0442-3](https://doi.org/10.1007/s10201-014-0442-3)
- Kaliszewicz, A., Johst, K., Grimm, V. & Uchmański, J. 2005. Predation effects on the evolution of life-history traits in a clonal oligochaete. *American Naturalist*, 166(3), 409-417. doi: [10.1086/432037](https://doi.org/10.1086/432037)
- Kaliszewicz, A. & Uchmański, J. 2009. A cross-phyla response to daphnia chemical alarm substances by an aquatic oligochaete. *Ecological Research*, 24(2), 461-466. doi: [10.1007/s11284-008-0522-0](https://doi.org/10.1007/s11284-008-0522-0)
- Kaliszewicz, A. & Uchmański, J. 2009. Damage released prey alarm substances or predator odours? Risk assessment by an aquatic oligochaete. *Hydrobiologia*, 618(1), 57-64. doi: [10.1007/s10750-008-9547-1](https://doi.org/10.1007/s10750-008-9547-1)
- Karickhoff, S.W. & Morris, K.R. 1985. Impact of tubificid oligochaetes on pollutant transport in bottom sediments. *Environmental Science & Technology*, 19(1), 51-56. doi: [10.1021/es00131a004](https://doi.org/10.1021/es00131a004)
- Kaster, J.L. 1989. Observations of predator-prey interaction on dispersal of an oligochaete prey, *Limnodrilus hoffmeisteri*. *Hydrobiologia*, 180(1), 191-193. doi: [10.1007/BF00027552](https://doi.org/10.1007/BF00027552)
- Kathman, R.D. & Wetzel, M.J. 2003. *Allonais inaequalis* (Annelida: Oligochaeta: Tubificidae) in North America. *Proceedings of the Biological Society of Washington*, 116(3), 548-556. Retrieved from: www.scopus.com
- Kaviraj, A., Bhunia, F. & Saha, N.C. 2004. Toxicity of methanol to fish, crustacean, oligochaete worm, and aquatic ecosystem. *International Journal of Toxicology*, 23(1), 55-63. doi: [10.1080/10915810490265469](https://doi.org/10.1080/10915810490265469)
- Kaygorodova, I.A. & Sherbakov, D.Y. 2006. Molecular phylogenetic study of the systematic position of Baikalian oligochaetes in Clitellata. *Russian Journal of Genetics*, 42(12), 1390-1397. doi: [10.1134/S1022795406120052](https://doi.org/10.1134/S1022795406120052)
- Kaygorodova, I.A., Verdonschot, P.F.M. & Kravtsova, L.S. 2012. Freshwater oligochaetes (Oligochaeta, Clitellata, Annelida) of north pribaikalye (east Siberia, Russia). *Turkish Journal of Zoology*, 36(1), 47-58. doi: [10.3906/zoo-1003-140](https://doi.org/10.3906/zoo-1003-140)
- Kazancı, N. & Girgin, S. 1998. Distribution of Oligochaeta species as bioindicators of organic pollution in Ankara stream and their use in biomonitoring. *Turkish Journal of Zoology*, 22(1), 83-87. Retrieved from: www.scopus.com
- Keilty, T.J. & Landrum, P.F. 1990. Population-specific toxicity responses by the freshwater oligochaete, *Stylodrilus heringianus*, in natural Lake Michigan sediments. *Environmental Toxicology and Chemistry*, 9(9), 1147-1154. doi: [10.1002/etc.5620090905](https://doi.org/10.1002/etc.5620090905)
- Keilty, T.J., White, D.S. & Landrum, P.F. 1988. Short-term lethality and sediment avoidance assays with endrin-contaminated sediment and two oligochaetes from Lake Michigan. *Archives of Environmental Contamination and Toxicology*, 17(1), 95-101. doi: [10.1007/BF01055159](https://doi.org/10.1007/BF01055159)
- Kerovec, M. & Kerovec, M. 2014. Oligochaeta and polychaeta fauna of the croatian part of the Sava River. [Fauna Oligochaeta i Polychaeta hrvatskog dijela rijeke Save]. *Natura Croatica*, 23(2), 335-348. Retrieved from: www.scopus.com
- Ketelaars, H.A.M., Kuipers, A.M.J.P. & Van Breemen, L.W.C.A. 1992. Temporal and spatial distribution of chironomid larvae and oligochaetes in two dutch storage reservoirs. *Netherlands Journal of Aquatic Ecology*, 26(2-4), 361-369. doi: [10.1007/BF02255263](https://doi.org/10.1007/BF02255263)
- Kharin, A.V., Zagainova, I.V. & Kostiuchenko, R.P. 2006. Formation of the zone of paratomy in freshwater oligochaetes. *Ontogenet*, 37(6), 424-437. Retrieved from: www.scopus.com
- Kharin, A.V., Zagainova, I.V. & Kostyuchenko, R.P. 2006. Formation of the paratomic fission zone in freshwater oligochaetes. *Russian Journal of Developmental Biology*, 37(6), 354-365. doi: [10.1134/S1062360406060038](https://doi.org/10.1134/S1062360406060038)
- Kiliç, V., Altunsoy, F. & Kiliç, G.A. 2011. Effect of thallium on the survival and morphology of *Tubifex tubifex* (Oligochaeta, Tubificidae). *Fresenius Environmental Bulletin*, 20(9 A), 2442-2445. Retrieved from: www.scopus.com
- Kitakoshi, T. & Shimizu, T. 2010. An oligochaete homologue of the brachyury gene is expressed transiently in the third quartette of micromeres. *Gene Expression Patterns*, 10(6), 306-313. doi: [10.1016/j.gep.2010.07.001](https://doi.org/10.1016/j.gep.2010.07.001)
- Klerks, P.L. & Bartholomew, P.R. 1991. Cadmium accumulation and detoxification in a cd-resistant population of the oligochaete *Limnodrilus hoffmeisteri*. *Aquatic Toxicology*, 19(2), 97-112. doi: [10.1016/0166-445X\(91\)90030-D](https://doi.org/10.1016/0166-445X(91)90030-D)
- Clump, J.V., Krezski, J.R., Smith, M. & Kaster, J.L. 1987. Dual tracer studies of the assimilation of an organic contaminant from sediments by deposit feeding oligochaetes. *Canadian Journal of Fisheries and Aquatic Sciences*, 44(9), 1574-1583. doi: [10.1139/f87-190](https://doi.org/10.1139/f87-190)

- Koprivnikar, J. & Desser, S.S. 2002. A new form of raabeia-type actinosporean (Myxozoa) from the oligochaete *Uncinaria uncinata*. *Folia Parasitologica*, 49(2), 89-92. doi: [10.14411/fp.2002.017](https://doi.org/10.14411/fp.2002.017)
- Kowalski, D.A. & Bergersen, E.P. 2003. The toxicity of bayluscide and TFM to *Tubifex tubifex*: Implications for chemical control of the oligochaete host of *Myxobolus cerebralis*, the causative agent of whirling disease. *North American Journal of Aquaculture*, 65(3), 171-178. doi: [10.1577/C02-036](https://doi.org/10.1577/C02-036)
- Krezoski, J.R. & Robbins, J.A. 1985. Vertical distribution of feeding and particle-selective transport of ¹³⁷Cs in lake sediments by Lumbriculid oligochaetes. *Journal of Geophysical Research*, 90(C6), 11,999-“12,006”. Retrieved from: www.scopus.com
- Krodkiewska, M. 2006. Freshwater Oligochaeta in mining subsidence ponds in the upper Silesia region of southern Poland. *Journal of Freshwater Ecology*, 21(1), 177-179. doi: [10.1080/02705060.2006.9664111](https://doi.org/10.1080/02705060.2006.9664111)
- Krodkiewska, M. 2007. The distribution of *Potamoithrix bavaricus* (Oeschmann, 1913) (Oligochaeta) in anthropogenic freshwater habitats of an industrialised area (upper Silesia, Poland). *Limnologica*, 37(3), 259-263. doi: [10.1016/j.limno.2007.02.001](https://doi.org/10.1016/j.limno.2007.02.001)
- Krodkiewska, M. 2005. The Oligochaeta communities in the benthos of artificially heated Rybnik dam reservoir (Poland). *Journal of Freshwater Ecology*, 20(1), 117-122. doi: [10.1080/02705060.2005.9664944](https://doi.org/10.1080/02705060.2005.9664944)
- Krodkiewska, M. & Kostecki, M. 2015. Assessment of the restoration measures in a Man-Made Reservoir: Do oligochaete communities respond to the improvement of water quality? *Environmental Monitoring and Assessment*, 187(9) doi: [10.1007/s10661-015-4787-9](https://doi.org/10.1007/s10661-015-4787-9)
- Krodkiewska, M. & Królczyk, A. 2011. Impact of environmental conditions on bottom oligochaete communities in subsidence ponds (the Silesian upland, southern Poland). *International Review of Hydrobiology*, 96(1), 48-57. doi: [10.1002/iroh.201011284](https://doi.org/10.1002/iroh.201011284)
- Krodkiewska, M. & Michalik-Kucharz, A. 2009. The bottom Oligochaeta communities in sand pits of different trophic status in upper Silesia (southern Poland). *Aquatic Ecology*, 43(2), 437-444. doi: [10.1007/s10452-008-9199-2](https://doi.org/10.1007/s10452-008-9199-2)
- Krodkiewska, M. & Spyra, A. 2015. New data on the biology and habitat preferences of the oligochaete species *ripistes parasita* (Annelida: Clitellata: Naididae): A case study in a temporary woodland pond. *Biologia (Poland)*, 70(5), 615-624. doi: [10.1515/biolog-2015-0070](https://doi.org/10.1515/biolog-2015-0070)
- Kukkonen, J. & Landrum, P.F. 1994. Toxicokinetics and toxicity of sediment-associated pyrene to *Lumbriculus variegatus* (Oligochaeta). *Environmental Toxicology and Chemistry*, 13(9), 1457-1468. doi: [10.1002/etc.5620130909](https://doi.org/10.1002/etc.5620130909)
- Lafont, M., Durbec, A. & Ille, C. 1992. Oligochaete worms as biological descriptors of the interactions between surface and groundwaters: A first synthesis. *Regulated Rivers: Research & Management*, 7(1), 65-73. doi: [10.1002/rrr.3450070109](https://doi.org/10.1002/rrr.3450070109)
- Lafont, M., Juget, J. & Rofes, G. 1991. An environmental index based on lacustrine oligochaetes. [Un indice biologique lacustre base sur l'examen des oligochetes]. *Revue des Sciences de l'Eau*, 4(2), 253-268. Retrieved from: www.scopus.com
- Lafont, M. & Vivier, A. 2006. Oligochaete assemblages in the hyporheic zone and coarse surface sediments: Their importance for understanding of ecological functioning of watercourses. *Hydrobiologia*, 564 (1 spec. iss.), 171-181. doi: [10.1007/s10750-005-1717-9](https://doi.org/10.1007/s10750-005-1717-9)
- Lafont, M., Vivier, A., Nogueira, S., Namour, P. & Breil, P. 2006. Surface and hyporheic oligochaete assemblages in a french suburban stream. *Hydrobiologia*, 564 (1 spec. iss.), 183-193. doi: [10.1007/s10750-005-1718-8](https://doi.org/10.1007/s10750-005-1718-8)
- Lagauzère, S., Pischedda, L., Cuny, P., Gilbert, F., Stora, G. & Bonzom, J.-. 2009. Influence of *Chironomus riparius* (Diptera, Chironomidae) and *Tubifex tubifex* (Annelida, Oligochaeta) on oxygen uptake by sediments. Consequences of uranium contamination. *Environmental Pollution*, 157(4), 1234-1242. doi: [10.1016/j.envpol.2008.12.004](https://doi.org/10.1016/j.envpol.2008.12.004)
- Lang, C. 2001. Anomalous response of zoobenthos (Oligochaetes and Chironomids) to the decrease of phosphorus in Lake Neuchâtel. [Réponse anormale du zoobenthos (Oligochètes et Chironomides) à la baisse du phosphore dans le lac de Neuchâtel]. *Revue Suisse de Zoologie*, 108(4), 891-903. doi: [10.5962/bhl.part.80167](https://doi.org/10.5962/bhl.part.80167)
- Lang, C. 1999. Contrasting responses of oligochaetes (Annelida) and chironomids (Diptera) to the abatement of eutrophication in Lake Neuchâtel. *Aquatic Sciences*, 61(3), 206-214. doi: [10.1007/PL00001324](https://doi.org/10.1007/PL00001324)
- Lang, C. 2010. Ecological state of sediment in two mountain lakes indicated by Oligochaetes and Chironomids. [Etat écologique des sédiments de deux lacs de montagne indiqué par les oligochètes et les chironomides]. *Bulletin de la Société Vaudoise des Sciences Naturelles*, 92(2), 47-60. Retrieved from: www.scopus.com
- Lang, C. 1985. Eutrophication of Lake Geneva indicated by the oligochaete communities of the profundal. *Hydrobiologia*, 126(3), 237-243. doi: [10.1007/BF00007501](https://doi.org/10.1007/BF00007501)
- Lang, C. 1989. Eutrophication of Lake Neuchâtel indicated by the oligochaete communities. *Hydrobiologia*, 174(1), 57-65. doi: [10.1007/BF00006058](https://doi.org/10.1007/BF00006058)
- Lang, C. 2007. How to interpret changes of zoobenthos (Oligochaetes and Chironomids) in Lake Neuchâtel between 1984 and 2002? [Comment interpréter les modifications du zoobenthos (Oligochètes et Chironomides) dans le lac de Neuchâtel entre 1984 et 2002?]. *Bulletin de la Société Vaudoise des Sciences Naturelles*, 90(3), 133-149. Retrieved from: www.scopus.com
- Lang, C. 2009. Indices based on Oligochaetes and Chironomids indicating the ecological recovery of sediments in Lake Geneva (Léman). [Indices basés sur les oligochètes et les chironomides indiquant la restauration écologique des sédiments du Léman]. *Bulletin de la Société Vaudoise des Sciences Naturelles*, 91(3), 283-300. Retrieved from: www.scopus.com
- Lang, C. 1997. Oligochaetes, organic sedimentation, and trophic state: How to assess the biological recovery of sediments in lakes? *Aquatic Sciences*, 59(1), 26-33. doi: [10.1007/BF02522547](https://doi.org/10.1007/BF02522547)
- LANG, C. 1990. Quantitative relationships between oligochaete communities and phosphorus concentrations in lakes. *Freshwater Biology*, 24(2), 327-334. doi: [10.1111/j.1365-2427.1990.tb00713.x](https://doi.org/10.1111/j.1365-2427.1990.tb00713.x)
- Lang, C. 2000. Response of oligochaete (Tubificidae and Lumbriculidae) and diptera Chironomidae communities to the decrease of phosphorus concentrations in Lake Geneva (little lake). [Reponse des communautés d'oligochètes (Tubificides et Lumbriculides) et de diptères Chironomides à la baisse des teneurs en phosphore dans le Leman (Petit Lac)]. *Annales de Limnologie*, 36(1), 13-20. doi: [10.1051/limm/2000001](https://doi.org/10.1051/limm/2000001)
- Lang, C. 1998. Using oligochaetes to monitor the decrease of eutrophication: The 1982-1996 trend in Lake Geneva. *Archiv Fur Hydrobiologie*, 141(4), 447-458. Retrieved from: www.scopus.com
- Lang, C. & Lods-Crozet, B. 1997. Oligochaetes versus chironomids as indicators of trophic state in two swiss lakes recovering from eutrophication. *Archiv Fur Hydrobiologie*, 139(2), 187-195. Retrieved from: www.scopus.com
- Lang, C. & Reymond, O. 1995. Contrasting responses of oligochaete communities to the abatement of eutrophication in Lake Geneva. *Hydrobiologia*, 308(1), 77-82. doi: [10.1007/BF00037789](https://doi.org/10.1007/BF00037789)
- Lang, C. & Reymond, O. 1995. Empirical relationships between oligochaete communities and phosphorus in lakes before and after the abatement of eutrophication. *Archiv Fur Hydrobiologie*, 134(4), 491-497. Retrieved from: www.scopus.com
- Lang, C. & Reymond, O. 1996. Empirical relationships between oligochaetes, phosphorus and organic deposition during the recovery of Lake Geneva from eutrophication. *Archiv Fur Hydrobiologie*, 136(2), 237-245. Retrieved from: www.scopus.com

- Lang, C. & Reymond, O. 1995. Recovery from eutrophication delayed in the profundal of Lake Neuchatel: Evidence from the oligochaete communities. *Revue Suisse de Zoologie*, 102(4), 907-912. Retrieved from: www.scopus.com
- Lang, C. & Reymond, O. 1992. Reversal of eutrophication in Lake Geneva: Evidence from the oligochaete communities. *Freshwater Biology*, 28(2), 145-148. doi: [10.1111/j.1365-2427.1992.tb00571.x](https://doi.org/10.1111/j.1365-2427.1992.tb00571.x)
- Lanzavecchia, G., Valvassori, R. & de Egileor, M. 1994. Body wall muscles in oligochaetes. *Hydrobiologia*, 278(1-3), 179-188. doi: [10.1007/BF00142326](https://doi.org/10.1007/BF00142326)
- Larsson, J. 1989. Light and electron microscope studies on *Jirovecia involuta* sp. nov. (Microspora, Bacillidiidae), a new microsporidian parasite of oligochaetes in Sweden. *European Journal of Protistology*, 25(2), 172-181. doi: [10.1016/S0932-4739\(89\)80028-8](https://doi.org/10.1016/S0932-4739(89)80028-8)
- Lasier, P.J. 2009. Alternative substrates for culturing the freshwater oligochaete *Lumbriculus variegatus*. *North American Journal of Aquaculture*, 71(1), 87-92. doi: [10.1577/A07-104.1](https://doi.org/10.1577/A07-104.1)
- Lauritsen, D.D., Mozley, S.C. & White, D.S. 1985. Distribution of oligochaetes in Lake Michigan and comments on their use as indices of pollution. *Journal of Great Lakes Research*, 11(1), 67-76. doi: [10.1016/S0380-1330\(85\)71745-5](https://doi.org/10.1016/S0380-1330(85)71745-5)
- Lazim, M.N. & Learner, M.A. 1987. The influence of sediment composition and leaf litter on the distribution of Tubificid worms (Oligochaeta) – A field and laboratory study. *Oecologia*, 72(1), 131-136. doi: [10.1007/BF00385057](https://doi.org/10.1007/BF00385057)
- Lazim, M.N. & Learner, M.A. 1986. The life-cycle and productivity of *Tubifex* (Oligochaeta; Tibificidae) in the Moat-Feeder stream, Cardiff, south wales. *Ecography*, 9(3), 185-192. doi: [10.1111/j.1600-0587.1986.tb01208.x](https://doi.org/10.1111/j.1600-0587.1986.tb01208.x)
- Lazim, M.N., Learner, M.A. & Cooper, S. 1989. The importance of worm identity and life history in determining the vertical distribution of Tibificids (Oligochaeta) in a riverine mud. *Hydrobiologia*, 178(1), 81-92. doi: [10.1007/BF00006115](https://doi.org/10.1007/BF00006115)
- Leppänen, M.T. & Kukkonen, J.V.K. 1998. Factors affecting feeding rate, reproduction and growth of an oligochaete *Lumbriculus variegatus* (Müller). *Hydrobiologia*, 377(1-3), 183-194. Retrieved from: www.scopus.com
- Leppänen, M.T. & Kukkonen, J.V.K. 1998. Relative importance of ingested sediment and pore water as bioaccumulation routes for pyrene to oligochaete (*Lumbriculus variegatus*, Müller). *Environmental Science & Technology*, 32(10), 1503-1508. doi: [10.1021/es970941k](https://doi.org/10.1021/es970941k)
- Leppänen, M.T. & Kukkonen, J.V.K. 2004. Toxicokinetics of sediment-associated polybrominated diphenylethers (Flame retardants) in benthic invertebrates (*Lumbriculus variegatus*, Oligochaeta). *Environmental Toxicology and Chemistry*, 23(1), 166-172. doi: [10.1897/03-68](https://doi.org/10.1897/03-68)
- Lesiuk, N.M. & Drewes, C.D. 1999. Autotomy reflex in a freshwater oligochaete, *Lumbriculus variegatus* (Clitellata: Lumbriculidae). *Hydrobiologia*, 406, 253-261. doi: [10.1023/A:1003756722019](https://doi.org/10.1023/A:1003756722019)
- Lesiuk, N.M. & Drewes, C.D. 2001. Behavioral plasticity and central regeneration of locomotor reflexes in the freshwater oligochaete, *Lumbriculus variegatus*. II. Ablation studies. *Invertebrate Biology*, 120(3), 259-268. doi: [10.1111/j.1744-7410.2001.tb00036.x](https://doi.org/10.1111/j.1744-7410.2001.tb00036.x)
- Liebig, M., Egeler, P., Oehlmann, J. & Knacker, T. 2005. Bioaccumulation of 14C-17 α -ethinylestradiol by the aquatic oligochaete *Lumbriculus variegatus* in spiked artificial sediment. *Chemosphere*, 59(2), 271-280. doi: [10.1016/j.chemosphere.2004.10.051](https://doi.org/10.1016/j.chemosphere.2004.10.051)
- Lietz, D.M. 1987. Potential for aquatic oligochaetes as live food in commercial aquaculture. *Hydrobiologia*, 155(1), 309-310. doi: [10.1007/BF00025665](https://doi.org/10.1007/BF00025665)
- Lin, K.-. & Yo, S.-. 2008. The effect of organic pollution on the abundance and distribution of aquatic oligochaetes in an urban water basin, Taiwan. *Hydrobiologia*, 596(1), 213-223. doi: [10.1007/s10750-007-9098-x](https://doi.org/10.1007/s10750-007-9098-x)
- Lindegaard, C., Hamburger, K. & Dall, P.C. 1994. Population dynamics and energy budget of *Marionina southerni* (Cernosvitov) (Enchytraeidae, Oligochaeta) in the shallow littoral of Lake Esrom, Denmark. *Hydrobiologia*, 278(1-3), 291-301. doi: [10.1007/BF00142338](https://doi.org/10.1007/BF00142338)
- Linhares Frizzera, G. & da Gama Alves, R. 2012. The influence of taxonomic resolution of Oligochaeta on the evaluation of water quality in an urban stream in Minas Gerais, Brazil. [A influência da resolução taxonômica de Oligochaeta na avaliação da qualidade da água em um córrego urbano em Minas Gerais, Brasil]. *Acta Limnologica Brasiliensis*, 24(4), 408-416. doi: [10.1590/S2179-975X201300500005](https://doi.org/10.1590/S2179-975X201300500005)
- Lobo, H. & Alves, R.G. 2011. Influence of body weight and substrate granulometry on the reproduction of *Limnodrilus hoffmeisteri* (Oligochaeta: Naididae: Tubificinae). *Zoologia*, 28(5), 558-564. doi: [10.1590/S1984-46702011000500002](https://doi.org/10.1590/S1984-46702011000500002)
- Lobo, H. & da Gama Alves, R. 2011. Reproductive cycle of *Branchiura sowerbyi* (Oligochaeta: Naididae: Tubificinae) cultivated under laboratory conditions. *Zoologia*, 28(4), 427-431. doi: [10.1590/S1984-46702011000400003](https://doi.org/10.1590/S1984-46702011000400003)
- Lods-Crozet, B. & Reymond, O. 2005. Ten years trends in the oligochaete and chironomid fauna of Lake Neuchâtel (Switzerland). *Revue Suisse de Zoologie*, 112(2), 543-558. doi: [10.5962/bhl.part.80313](https://doi.org/10.5962/bhl.part.80313)
- Löhlein, B. 1999. Assessment of secondary production of Naididae (Oligochaeta): An example from a north german lake. *Hydrobiologia*, 406, 191-198. doi: [10.1023/A:1003736217476](https://doi.org/10.1023/A:1003736217476)
- Löhlein, B. & Böttger, K. 1996. The Oligochaeta (Annelida) of a near-natural lowland stream in northern Germany. [Die Oligochaeta (Annelida) eines naturnahen baches im Norddeutschen tiefland]. *Limnologica*, 26(4), 393-402. Retrieved from: www.scopus.com
- López Van Oosterom, M.V., Ocon, C., Armendariz, L.C. & Rodrigues Capítulo, A. 2015. Structural and functional responses of the oligochaete and aeolosomatid assemblage in lowland streams: A one-way-pollution-modelled ecosystem. *Journal of Limnology*, 74(3) doi: [10.4081/jlimnol.2015.1142](https://doi.org/10.4081/jlimnol.2015.1142)
- Lotufo, G.R. & Fleeger, J.W. 1996. Toxicity of sediment-associated pyrene and phenanthrene to *Limnodrilus hoffmeisteri* (Oligochaeta: Tubificidae). *Environmental Toxicology and Chemistry*, 15(9), 1508-1516. Retrieved from: www.scopus.com
- Lowers, J.M. & Bartholomew, J.L. 2003. Detection of myxozoan parasites in oligochaetes imported as food for ornamental fish. *Journal of Parasitology*, 89(1), 84-91. doi: [10.1645/0022-3395\(2003\)089\[0084:DOMPIO\]2.0.CO;2](https://doi.org/10.1645/0022-3395(2003)089[0084:DOMPIO]2.0.CO;2)
- Lu, X., Reible, D.D., Fleeger, J.W. & Chai, Y. 2003. Bioavailability of desorption-resistant phenanthrene to the oligochaete *Ilyodrilus templetoni*. *Environmental Toxicology and Chemistry*, 22(1), 153-160. Retrieved from: www.scopus.com
- Lucan-Bouché, M.-., Arsac, F., Biagianti-Risbourg, S., Habets, F. & Vernet, G. 1997. Experimental study of the lethal effects induced by copper and lead on the Oligochaeta *Tubifex tubifex*. [Étude expérimentale des effets létaux induits par le cuivre et/ou le plomb chez l'Oligochète *Tubifex tubifex*]. *Bulletin de la Société Zoologique de France*, 122(4), 389-392. Retrieved from: www.scopus.com
- Lucan-Bouché, M.-., Biagianti-Risbourg, S., Arsac, F. & Vernet, G. 1999. An original decontamination process developed by the aquatic oligochaete *Tubifex tubifex* exposed to copper and lead. *Aquatic Toxicology*, 45(1), 9-17. doi: [10.1016/S0166-445X\(98\)00091-5](https://doi.org/10.1016/S0166-445X(98)00091-5)
- Lucan-Bouché, M.-., Biagianti-Risbourg, S., Arsac, F. & Vernet, G. 1999. Autotomy as a mechanism of decontamination used by the oligochaete *Tubifex tubifex*. [L'autotomie comme moyen de décontamination utilisé par l'oligochète *Tubifex tubifex*]. *Bulletin de la Société Zoologique de France*, 124(4), 383-387. Retrieved from: www.scopus.com

- Lyytikäinen, M., Rantalainen, A.-., Mikkelsen, P., Hämäläinen, H., Paasivirta, J. & Kukkonen, J.V.K. 2003. Similarities in bioaccumulation patterns of polychlorinated dibenzo-p-dioxins and furans and polychlorinated diphenyl ethers in laboratory-exposed oligochaetes and semipermeable membrane devices and in field-collected chironomids. *Environmental Toxicology and Chemistry*, 22(10), 2405-2415. doi: [10.1897/02-433](https://doi.org/10.1897/02-433)
- Mackie, J.A., Levinton, J.S., Przeslawski, R., Delambert, D. & Wallace, W. 2010. Loss of evolutionary resistance by the oligochaete Limnodrilus hoffmeisteri to a toxic substance – cost or gene flow? *Evolution*, 64(1), 152-165. doi: [10.1111/j.1558-5646.2009.00806.x](https://doi.org/10.1111/j.1558-5646.2009.00806.x)
- MacMichael, G.J., Brown, L.R. & Ladner, C.M. 1988. An oligochaete as a potential food source for fish in aquaculture. *Progressive Fish-Culturist*, 50(1), 31-35. doi: [10.1577/1548-8640\(1988\)050\[0031:AOAAPF\]2.3.CO;2](https://doi.org/10.1577/1548-8640(1988)050[0031:AOAAPF]2.3.CO;2)
- Mäenpää, K. & Kukkonen, J.V.K. 2006. Bioaccumulation and toxicity of 4-nonylphenol (4-NP) and 4-(2-dodecyl)-benzene sulfonate (LAS) in *Lumbriculus variegatus* (Oligochaeta) and *Chironomus riparius* (Insecta). *Aquatic Toxicology*, 77(3), 329-338. doi: [10.1016/j.aquatox.2006.01.002](https://doi.org/10.1016/j.aquatox.2006.01.002)
- Mäenpää, K., Leppänen, M.T. & Kukkonen, J.V.K. 2009. Sublethal toxicity and biotransformation of pyrene in *Lumbriculus variegatus* (Oligochaeta). *Science of the Total Environment*, 407(8), 2666-2672. doi: [10.1016/j.scitotenv.2009.01.019](https://doi.org/10.1016/j.scitotenv.2009.01.019)
- Mäenpää, K., Sorsa, K., Lyytikäinen, M., Leppänen, M.T. & Kukkonen, J.V.K. 2008. Bioaccumulation, sublethal toxicity, and biotransformation of sediment-associated pentachlorophenol in *Lumbriculus variegatus* (Oligochaeta). *Ecotoxicology and Environmental Safety*, 69(1), 121-129. doi: [10.1016/j.ecoenv.2006.12.019](https://doi.org/10.1016/j.ecoenv.2006.12.019)
- Mäenpää, K.A., Sormunen, A.J. & Kukkonen, J.V.K. 2003. Bioaccumulation and toxicity of sediment associated herbicides (ioxynil, pendimethalin, and bentazone) in *Lumbriculus variegatus* (Oligochaeta) and *Chironomus riparius* (Insecta). *Ecotoxicology and Environmental Safety*, 56(3), 398-410. doi: [10.1016/S0147-6513\(03\)00010-1](https://doi.org/10.1016/S0147-6513(03)00010-1)
- Maestre, Z., Martinez-Madrid, M., Rodriguez, P. & Reynoldson, T. 2007. Ecotoxicity assessment of river sediments and a critical evaluation of some of the procedures used in the aquatic oligochaete *Tubifex tubifex* chronic bioassay. *Archives of Environmental Contamination and Toxicology*, 53(4), 559-570. doi: [10.1007/s00244-006-0264-3](https://doi.org/10.1007/s00244-006-0264-3)
- Manum, S.B., Bose, M.N. & Sawyer, R.T. 1991. Clitellate cocoons in freshwater deposits since the triassic. *Zoologica Scripta*, 20(4), 347-366. doi: [10.1111/j.1463-6409.1991.tb00300.x](https://doi.org/10.1111/j.1463-6409.1991.tb00300.x)
- Marchese, M. 1994. Population dynamics of *Narapa bonettoi righi* and *varela*, 1983 (Oligochaeta: Narapidae) from the main channel of the middle Paraná River, Argentina. *Hydrobiologia*, 278(1-3), 303-308. doi: [10.1007/BF00142339](https://doi.org/10.1007/BF00142339)
- Marchese, M.R. 1987. The ecology of some benthic Oligochaeta from the Paraná River, Argentina. *Hydrobiologia*, 155(1), 209-214. doi: [10.1007/BF00025653](https://doi.org/10.1007/BF00025653)
- Marotta, R., Ferraguti, M., Erséus, C. & Gustavsson, L.M. 2008. Combined-data phylogenetics and character evolution of Clitellata (Annelida) using 18S rDNA and morphology. *Zoological Journal of the Linnean Society*, 154(1), 1-26. doi: [10.1111/j.1096-3642.2008.00408.x](https://doi.org/10.1111/j.1096-3642.2008.00408.x)
- Martin, P. & Brinkhurst, R.O. 1998. A new genus and two new species of Tubificidae (Oligochaeta) from the abyssal zone of Lake Baikal with redescriptions of *Lymphachaeta pinnigera*, *Rhyacodriloides abyssalis* and *Tubifex bazikalovae*. *Zoologica Scripta*, 27(3), 197-207. doi: [10.1111/j.1463-6409.1998.tb00437.x](https://doi.org/10.1111/j.1463-6409.1998.tb00437.x)
- Martin, P. & Brinkhurst, R.O. 1994. A new species of *Insulodrilus* (Oligochaeta, Phreodrilidae) from Lake Tanganyika (East Africa) with notes on the oligochaete fauna of the lake. *Archiv Fur Hydrobiologie*, 130(2), 249-256. Retrieved from: www.scopus.com
- Martin, P., Ferraguti, M. & Kaygorodova, I. 1998. Description of two new species of *Rhynchelmis* (Oligochaeta: Lumbriculidae) from Lake Baikal (Russia), using classical morphology and ultrastructure of spermatozoa. *Annales de Limnologie*, 34(3), 283-293. doi: [10.1051/limn/1998025](https://doi.org/10.1051/limn/1998025)
- Martin, P. & Giani, N. 1995. *Insulodrilus martensi*, a new species of Phreodrilidae (Oligochaeta) from Lake Tanganyika (East Africa). [*Insulodrilus martensi*, une nouvelle espèce de Phreodrilidae (Oligochaeta) du lac Tanganyika (Afrique de l'Est)]. *Annales de Limnologie*, 31(1), 3-8. doi: [10.1051/limn/1995003](https://doi.org/10.1051/limn/1995003)
- Martin, P. & Giani, N. 1995. Two new species of *epirodrilus* (Oligochaeta, Tubificidae) from Lake Nyasa and Tanganyika (East Africa), with redescriptions of *E. slovenicus* and *E. michaelensi*. *Zoologica Scripta*, 24(1), 13-19. doi: [10.1111/j.1463-6409.1995.tb00471.x](https://doi.org/10.1111/j.1463-6409.1995.tb00471.x)
- Martin, P., Goddeeris, B. & Martens, K. 1994. Depth distribution of oligochaetes in Lake Baikal (Siberia, Russia). *Hydrobiologia*, 278(1-3), 151-156. doi: [10.1007/BF00142323](https://doi.org/10.1007/BF00142323)
- Martin, P., Kaygorodova, I., Sherbakov, D.Y. & Verheyen, E. 2000. Rapidly evolving lineages impede the resolution of phylogenetic relationships among Clitellata (Annelida). *Molecular Phylogenetics and Evolution*, 15(3), 355-368. doi: [10.1006/mpev.1999.0764](https://doi.org/10.1006/mpev.1999.0764)
- Martin, P., Martens, K. & Goddeeris, B. 1999. Oligochaeta from the abyssal zone of Lake Baikal (Siberia, Russia). *Hydrobiologia*, 406, 165-174. doi: [10.1023/A:1003719430680](https://doi.org/10.1023/A:1003719430680)
- Martin, P., Martinez-Ansemil, E., Pinder, A., Timm, T. & Wetzel, M.J. 2008. Global diversity of Oligochaetous clitellates ("Oligochaeta"; Clitellata) in freshwater. *Hydrobiologia*, 595(1), 117-127. doi: [10.1007/s10750-007-9009-1](https://doi.org/10.1007/s10750-007-9009-1)
- Martin, P., Schmelz, R.M. & Dole-Olivier, M.-. 2015. Groundwater oligochaetes (Annelida, Clitellata) from the Mercantour National Park (France), with the descriptions of one new genus and two new stygobiont species. *Zoosystema*, 37(4), 551-569. doi: [10.5252/z2015n4a2](https://doi.org/10.5252/z2015n4a2)
- Martínez, D.E. & Levinton, J. 1996. Adaptation to heavy metals in the aquatic oligochaete *Limnodrilus hoffmeisteri*: Evidence for control by one gene. *Evolution*, 50(3), 1339-1343. Retrieved from: www.scopus.com
- Martinez-Ansemil, E. 1990. Biological and ecological study of the aquatic oligochaetes of the River Tambre and its environment (Galicia, Spain). [Etude biologique et écologique des Oligochètes aquatiques de la rivière Tambre et de ses milieux associés (Galice, Espagne)]. *Annales de Limnologie*, 26(2), 131-151. doi: [10.1051/limn/1990013](https://doi.org/10.1051/limn/1990013)
- Martínez-Ansemil, E. & Collado, R. 1996. Two new species of freshwater Oligochaeta from the north-west Iberian Peninsula: *Krenedrilus realis* sp. nov. (Tubificidae) and *Cernosvitoviella bulbducta* sp. nov. (Enchytraeidae). *Journal of Zoology*, 240(2), 363-370. doi: [10.1111/j.1469-7998.1996.tb05291.x](https://doi.org/10.1111/j.1469-7998.1996.tb05291.x)
- Martínez-Ansemil, E., Creuzé Des Châtelliers, M., Martin, P. & Sambugar, B. 2012. The Parvidrilidae – a diversified groundwater family: Description of six new species from southern Europe, and clues for its phylogenetic position within Clitellata (Annelida). *Zoological Journal of the Linnean Society*, 166(3), 530-558. doi: [10.1111/j.1096-3642.2012.00857.x](https://doi.org/10.1111/j.1096-3642.2012.00857.x)
- Martínez-Ansemil, E. & Giani, N. 1987. The distribution of aquatic oligochaetes in the south and eastern Mediterranean area. *Hydrobiologia*, 155(1), 293-303. doi: [10.1007/BF00025662](https://doi.org/10.1007/BF00025662)
- Martínez-Ansemil, E., Giani, N. & Sambugar, B. 2002. Oligochaetes from underground waters of Oman with descriptions of two new species of Phreodrilidae (Oligochaeta): *Antarctodrilus arabicus* sp. nov., and *Phreodrilus stocki* sp. nov. *Contributions to Zoology*, 71(4), 147-158. Retrieved from: www.scopus.com
- Martínez-Ansemil, E., Sambugar, B. & Giani, N. 2002. First record of Parvidrilidae (Annelida, Oligochaeta) in Europe with a description of a

- new species (*Parvidrilus spelaeus* sp. nov.) and comments on the family and its phyletic relationships. *Journal of Zoology*, 256(4), 495-503. Retrieved from: www.scopus.com
- Martínez-Ansemil, E., Sambugar, B. & Giani, N. 1997. Groundwater oligochaetes from southern-europe. I. A new genus and three new species of rhyacodrilinae (Tubificidae) with a redescription of *Tubifex pescei* (Dumnicka) comb. nov. *Annales de Limnologie*, 33(1), 33-44. doi: [10.1051/limn/1997003](https://doi.org/10.1051/limn/1997003)
- Martinez-Tabche, L., Ortega, M.D.L.A.G., Ramirez Mora, B., German Faz, C., Lopez Lopez, E. & Galar Martinez, M. 2001. Hemoglobin concentration and acetylcholinesterase activity of oligochaetes in relation to lead concentration in spiked sediments from Ignacio Ramirez Reservoir. *Ecotoxicology and Environmental Safety*, 49(1), 76-83. doi: [10.1006/eesa.2001.2030](https://doi.org/10.1006/eesa.2001.2030)
- Martinovic-Vitanovic, V., Djikanovic, V., Obradovic, S. & Kalafatic, V. 2007. Composition and structure of the Oligochaeta (Annelida) in benthic assemblages of the Danube River in the Belgrade region during may and october of 2004. *Ekologia Bratislava*, 26(2), 174-189. Retrieved from: www.scopus.com
- Martins, R.T. & Alves, R.G. 2010. Occurrence of *Chaetogaster limnaei* K. von Baer, 1927 (Oligochaeta, Naididae) associated with Gastropoda mollusks in horticultural channels in southeastern Brazil. [Ocorrência de *Chaetogaster limnaei* K. von Baer, 1927 (Oligochaeta, Naididae) associado com moluscos Gastropoda em canais de horticultura no sudeste do Brasil]. *Brazilian Journal of Biology*, 70(4), 1055-1057. doi: [10.1590/S1519-69842010000500020](https://doi.org/10.1590/S1519-69842010000500020)
- Martins, R.T. & Da Gama Alves, R. 2008. Occurrence of Naididae (Annelida: Oligochaeta) from three gastropod species in irrigation fields in southeastern Brazil. *Biota Neotropica*, 8(3), 255-257. doi: [10.1590/S1676-06032008000300023](https://doi.org/10.1590/S1676-06032008000300023)
- Martins, R.T., Silveira, L.S. & Alves, R.G. 2011. Colonization by oligochaetes (Annelida: Clitellata) in decomposing leaves of *Eichhornia azurea* (SW.) Kunth (Pontederiaceae) in a neotropical lentic system. *Annales de Limnologie*, 47(4), 339-346. doi: [10.1051/limn/2011053](https://doi.org/10.1051/limn/2011053)
- Mastrantuono, L. 1988. A note on the feeding of *Amphichaeta leydigii* (Oligochaeta, Naididae) in lacustrine sandy shores. *Hydrobiological Bulletin*, 22(2), 195-198. doi: [10.1007/BF02256823](https://doi.org/10.1007/BF02256823)
- Mathooko, J.M., Mpawenayo, B., Kipkemboi, J.K. & M'rimba, C.M. 2005. Distributional patterns of diatoms and *Limnodrilus* oligochaetes in a kenyan dry streambed following the 1999-2000 drought conditions. *International Review of Hydrobiology*, 90(2), 185-200. doi: [10.1002/irh.200310708](https://doi.org/10.1002/irh.200310708)
- Matisoff, G., Wang, X. & McCall, P.L. 1999. Biological redistribution of lake sediments by tubificid oligochaetes: Branchiura sowerbyi and *Limnodrilus hoffmeisteri*/Tubifex tubifex. *Journal of Great Lakes Research*, 25(1), 205-219. doi: [10.1016/S0380-1330\(99\)70729-X](https://doi.org/10.1016/S0380-1330(99)70729-X)
- Matsuo, K. & Shimizu, T. 2006. Embryonic expression of a decapentaplegic gene in the oligochaete annelid *Tubifex tubifex*. *Gene Expression Patterns*, 6(8), 800-806. doi: [10.1016/j.modgep.2006.01.009](https://doi.org/10.1016/j.modgep.2006.01.009)
- Matsuo, K., Yoshida, H. & Shimizu, T. 2005. Differential expression of caudal and dorsal genes in the teloblast lineages of the oligochaete annelid *Tubifex tubifex*. *Development Genes and Evolution*, 215(5), 238-247. doi: [10.1007/s00427-005-0473-1](https://doi.org/10.1007/s00427-005-0473-1)
- Measures, L.N. 1988. The development of Eustrengylides tubifex (Nematoda: Dioctophymatoidea) in oligochaetes. *Journal of Parasitology*, 74(2), 294-304. doi: [10.2307/3282459](https://doi.org/10.2307/3282459)
- Méndez-Fernández, L., De Jonge, M. & Bervoets, L. 2014. Influences of sediment geochemistry on metal accumulation rates and toxicity in the aquatic oligochaete *Tubifex tubifex*. *Aquatic Toxicology*, 157, 109-119. doi: [10.1016/j.aquatox.2014.10.009](https://doi.org/10.1016/j.aquatox.2014.10.009)
- Méndez-Fernández, L., Martínez-Madrid, M. & Rodriguez, P. 2013. Toxicity and critical body residues of CD, CU and CR in the aquatic oligochaete *Tubifex tubifex* (Müller) based on lethal and sublethal effects. *Ecotoxicology*, 22(10), 1445-1460. doi: [10.1007/s10646-013-1131-4](https://doi.org/10.1007/s10646-013-1131-4)
- Merle, C., Danielopol, D.L. & Watanamahart, P. 1997. Impact of environmental conditions on the habitat selection of interstitial-dwelling Tubificids (Oligochaeta). An experimental study. *Geobios*, 30 (suppl. 2), 91-99. doi: [10.1016/S0016-6995\(97\)80074-5](https://doi.org/10.1016/S0016-6995(97)80074-5)
- Mermilliod-Blondin, F., Creuze des Chatelliers, M., Gerino, M. & Gaudet, J.-. 2000. Testing the effect of *Limnodrilus* sp. (Oligochaeta, Tubificidae) on organic matter and nutrient processing in the hyporheic zone: A microcosm method. *Archiv Fur Hydrobiologie*, 149(3), 467-487. Retrieved from: www.scopus.com
- Mermilliod-Blondin, F., Gérino, M., Degrange, V., Lensi, R., Chassé, J.-, Rard, M. & Creuzé des Châtelliers, M. 2001. Testing the functional redundancy of *Limnodrilus* and *Tubifex* (Oligochaeta, Tubificidae) in hyporheic sediments: An experimental study in microcosms. *Canadian Journal of Fisheries and Aquatic Sciences*, 58(9), 1747-1759. doi: [10.1139/f01-119](https://doi.org/10.1139/f01-119)
- Mermilliod-Blondin, F., Poggiale, J.-, Tolla, C., Auger, P., Thuiller, W. & Des Châtelliers, M.C. 2008. Using a mathematical model to simulate the influence of Tubificid worms (Oligochaeta) on oxygen concentrations in hyporheic sediments. *Fundamental and Applied Limnology*, 172(2), 135-145. doi: [10.1127/1863-9135/2008/0172-0135](https://doi.org/10.1127/1863-9135/2008/0172-0135)
- Meyer, J.S., Boese, C.J. & Collyard, S.A. 2002. Whole-body accumulation of copper predicts acute toxicity to an aquatic oligochaete (*Lumbriculus variegatus*) as pH and calcium are varied. *Comparative Biochemistry and Physiology – Part C: Toxicology and Pharmacology*, 133(1-2), 99-109. doi: [10.1016/S1532-0456\(02\)00103-5](https://doi.org/10.1016/S1532-0456(02)00103-5)
- Mihaljević, Z., Kerovec, M., Ternjek, I. & Mrakovčić, M. 2001. Composition and depth distribution of oligochaete fauna in a mediterranean karstic lake. *Biologia*, 56(5), 461-467. Retrieved from: www.scopus.com
- Milbrink, G. 1993. Evidence for mutualistic interactions in freshwater oligochaete communities. *Oikos*, 68(2), 317-322. doi: [10.2307/3544845](https://doi.org/10.2307/3544845)
- Milbrink, G. 1994. Oligochaetes and water pollution in two deep norwegian lakes. *Hydrobiologia*, 278(1-3), 213-222. doi: [10.1007/BF00142329](https://doi.org/10.1007/BF00142329)
- Milbrink, G., Timm, T. & Lundberg, S. 2002. Indicative profundal oligochaete assemblages in selected small swedish lakes. *Hydrobiologia*, 468, 53-61. doi: [10.1023/A:1015274323026](https://doi.org/10.1023/A:1015274323026)
- Millward, R.N., Fleeger, J.W., Reible, D.D., Keteles, K.A., Cunningham, B.P. & Zhang, L. 2001. Pyrene bioaccumulation, effects of pyrene exposure on particle-size selection, and fecal pyrene content in the oligochaete *Limnodrilus hoffmeisteri* (Tubificidae, Oligochaeta). *Environmental Toxicology and Chemistry*, 20(6), 1359-1366. doi: [10.1002/etc.5620200627](https://doi.org/10.1002/etc.5620200627)
- Mischke, C.C. & Griffin, M.J. 2011. Laboratory mass culture of the freshwater oligochaete *Dero digitata*. *North American Journal of Aquaculture*, 73(1), 13-16. doi: [10.1080/15222055.2011.544936](https://doi.org/10.1080/15222055.2011.544936)
- Mischke, C.C., Terhune, J.S. & Wise, D.J. 2001. Acute toxicity of several chemicals to the oligochaete *Dero digitata*. *Journal of the World Aquaculture Society*, 32(2), 184-188. doi: [10.1111/j.1749-7345.2001.tb01093.x](https://doi.org/10.1111/j.1749-7345.2001.tb01093.x)
- Molnár, K., El-Mansy, A., Székely, C. & Baska, F. 1999. Development of *Myxobolus dispar* (Myxosporea: Myxobolidae) in an oligochaete alternate host, *Tubifex tubifex*. *Folia Parasitologica*, 46(1), 15-21. Retrieved from: www.scopus.com
- Molnár, L. & Fischer, E. 1989. The effect of saline load on the elemental composition of chloragosomes in *Limnodrilus hoffmeisteri* (Oligochaeta, Tubificidae). *Comparative Biochemistry and Physiology – Part A: Physiology*, 92(4), 551-553. doi: [10.1016/0300-9629\(89\)90364-2](https://doi.org/10.1016/0300-9629(89)90364-2)
- Montalto, L. & Marchese, M. 2005. Cyst formation in Tubificidae (Naidinae) and opistocystidae (Annelida, Oligochaeta) as an adaptive strategy for

- drought tolerance in fluvial wetlands of the Paraná River, Argentina. *Wetlands*, 25(2), 488-494. doi: [10.1672/23](https://doi.org/10.1672/23)
- Montanholi-Martins, M.C. & Takeda, A.M. 1999. Communities of benthic oligochaetes in relation to sediment structure in the upper Parana River, Brazil. *Studies on Neotropical Fauna and Environment*, 34(1), 52-58. Retrieved from: www.scopus.com
- Montanholi-Martins, M.C. & Takeda, A.M. 2001. Spatial and temporal variations of oligochaetes of the Ivinhema River and Patos Lake in the upper Paraná River basin, Brazil. *Hydrobiologia*, 463, 197-205. doi: [10.1023/A:1013163927814](https://doi.org/10.1023/A:1013163927814)
- Moretto, Y., Simões, N.R., Benedito, E. & Higuti, J. 2013. Effect of trophic status and sediment particle size on diversity and abundance of aquatic Oligochaeta (Annelida) in neotropical reservoirs. *Annales de Limnologie*, 49(1), 65-78. doi: [10.1051/limn/2013040](https://doi.org/10.1051/limn/2013040)
- Morev, Y.B. & Smilova, T.S. 1987. Food assimilation rate of Tubificid worms *Tubifex tubifex* (Oligochaeta). *Hydrobiological Journal*, 23(2), 69-72. Retrieved from: www.scopus.com
- Mosleh, Y.Y., Paris-Palacios, S., Arnoult, F., Couderchet, M., Biagiante-Risbourg, S. & Vernet, G. 2004. Metallothionein induction in aquatic oligochaete *Tubifex tubifex* exposed to herbicide isoproturon. *Environmental Toxicology*, 19(1), 88-93. doi: [10.1002/tox.10153](https://doi.org/10.1002/tox.10153)
- Mosleh, Y.Y., Paris-Palacios, S., Arnoult, F., Couderchet, M. & Vernet, G. 2003. Metallothionein induction in the aquatic oligochaete *Tubifex tubifex* exposed to the herbicide isoproturon. *Communications in Agricultural and Applied Biological Sciences*, 68(4 Pt B), 759-770. Retrieved from: www.scopus.com
- Mosleh, Y.Y., Paris-Palacios, S. & Biagiante-Risbourg, S. 2006. Metallothioneins induction and antioxidative response in aquatic worms *Tubifex tubifex* (Oligochaeta, Tubificidae) exposed to copper. *Chemosphere*, 64(1), 121-128. doi: [10.1016/j.chemosphere.2005.10.045](https://doi.org/10.1016/j.chemosphere.2005.10.045)
- Mosleh, Y.Y., Paris-Palacios, S., Couderchet, M., Biagiante-Risbourg, S. & Vernet, G. 2005. Metallothionein induction, antioxidative responses, glycogen and growth changes in *Tubifex tubifex* (Oligochaete) exposed to the fungicide, fenhexamid. *Environmental Pollution*, 135(1), 73-82. doi: [10.1016/j.envpol.2004.10.012](https://doi.org/10.1016/j.envpol.2004.10.012)
- Mount, D.R., Highland, T.L., Mattson, V.R., Dawson, T.D., Lott, K.G. & Ingersoll, C.G. 2006. Use of the oligochaete, *Lumbriculus variegatus*, as a prey organism for toxicant exposure of fish through the diet. *Environmental Toxicology and Chemistry*, 25(10), 2760-2767. doi: [10.1897/06-138.1](https://doi.org/10.1897/06-138.1)
- Müller, M.C.M. 2004. Nerve development, growth and differentiation during regeneration in *Enchytraeus fragmentosus* and *Stylaria lacustris* (Oligochaeta). *Development Growth & Differentiation*, 46(5), 471-478. doi: [10.1111/j.1440-169x.2004.00763.x](https://doi.org/10.1111/j.1440-169x.2004.00763.x)
- Naitoh, T. & Matsuno, A. 1985. Demonstration of the structural connections of the longitudinal muscle cells and circular muscle cells, and interconnections between the two, in the alimentary canal of an oligochaete, *Branchiura sowerbyi* Beddard. *Experientia*, 41(3), 370-372. doi: [10.1007/BF02004510](https://doi.org/10.1007/BF02004510)
- Nascimento, H.L.S. & Alves, R.D.G. 2008. Cocoon production and hatching rates of *Branchiura sowerbyi* Beddard (Oligochaeta: Tubificidae). *Revista Brasileira de Zoologia*, 25(1), 16-19. doi: [10.1590/S0101-81752008000100003](https://doi.org/10.1590/S0101-81752008000100003)
- Nascimento, H.L.S. & Alves, R.G. 2009. The effect of temperature on the reproduction of *Limnodrilus hoffmeisteri* (Oligochaeta: Tubificidae). *Zoologia*, 26(1), 191-193. doi: [10.1590/S1984-46702009000100026](https://doi.org/10.1590/S1984-46702009000100026)
- Nasi, M., Piol, M.N., Di Risio, C. & Verrengia Guerrero, N.R. 2011. Investigating arsenic bioavailability and bioaccumulation by the freshwater oligochaete *Lumbriculus variegatus*. *Archives of Environmental Contamination and Toxicology*, 61(3), 426-434. doi: [10.1007/s00244-010-9639-6](https://doi.org/10.1007/s00244-010-9639-6)
- Naveed, M.I. 2012. Preliminary studies on aquatic Oligochaeta in and around Chennai, Tamil Nadu, India. *Turkish Journal of Zoology*, 36(1), 25-37. doi: [10.3906/zoo-1002-33](https://doi.org/10.3906/zoo-1002-33)
- Nazaraghghi, F., Timm, T., Nadoushan, R.M., Shabanipour, N., Fatemi, M.R. & Moradi, A.M. 2014. Oligochaetes (Annelida, Clitellata) in the Anzali International Wetland, north-western Iran. *Estonian Journal of Ecology*, 63(3), 130-144. doi: [10.3176/eco.2014.3.02](https://doi.org/10.3176/eco.2014.3.02)
- Negredo, C. & Mulcahy, M.F. 2001. Actinosporean infections in oligochaetes in a river system in southwest Ireland with descriptions of three new forms. *Diseases of Aquatic Organisms*, 46(1), 67-77. doi: [10.3354/dao046067](https://doi.org/10.3354/dao046067)
- Newrka, P. & Mutayoba, S. 1987. Why and where do oligochaetes hide their cocoons? *Hydrobiologia*, 155(1), 171-178. doi: [10.1007/BF00025646](https://doi.org/10.1007/BF00025646)
- Newrka, P. & Wijegoonawardana, N. 1987. Vertical distribution and abundance of benthic invertebrates in profundal sediments of mondsee, with special reference to oligochaetes. *Hydrobiologia*, 155(1), 227-234. doi: [10.1007/BF00025655](https://doi.org/10.1007/BF00025655)
- Nie, X.-, Huang, T.-, Zhang, J.-, Li, W., Li, X.-, Zhang, S. & Wang, B.-. 2010. Study on inactivation effect and toxic mechanism of chlorine for the aquatic worm *Tubifex tubifex* (Oligochaeta, Tubificidae). *Sustainable Environment Research*, 20(5), 341-346. Retrieved from: www.scopus.com
- Nijboer, R.C., Wetzel, M.J. & Verdonschot, P.F.M. 2004. Diversity and distribution of Tubificidae, Naididae, and Lumbriculidae (Annelida: Oligochaeta) in the netherlands: An evaluation of twenty years of monitoring data. *Hydrobiologia*, 520(1-3), 127-141. doi: [10.1023/B:HYDR.0000027732.88238.61](https://doi.org/10.1023/B:HYDR.0000027732.88238.61)
- Nikkilä, A., Halme, A. & Kukkonen, J.V.K. 2003. Toxicokinetics, toxicity and lethal body residues of two chlorophenols in the oligochaete worm, *Lumbriculus variegatus*, in different sediments. *Chemosphere*, 51(1), 35-46. doi: [10.1016/S0045-6535\(02\)00791-9](https://doi.org/10.1016/S0045-6535(02)00791-9)
- Nikkilä, A., Kukkonen, J.V.K. & Oikari, A. 2001. Bioavailability of sediment-associated retene to an oligochaete worm *Lumbriculus variegatus*: Effects of sediment organic carbon and retene concentration. *Journal of Soils and Sediments*, 1(3), 137-145. doi: [10.1007/BF02986476](https://doi.org/10.1007/BF02986476)
- Nishino, M., Ohtaka, A. & Narita, T. 1999. Studies on the aquatic oligochaete fauna in Lake Biwa, central Japan. III. Distribution of respective taxa within the lake. *Japanese Journal of Limnology (Rikusuigaku Zasshi)*, 60(4), 539-558. doi: [10.3739/rikusui.60.539](https://doi.org/10.3739/rikusui.60.539)
- Nuutinen, S. & Kukkonen, J.V.K. 1998. The effect of selenium and organic material in lake sediments on the bioaccumulation of methylmercury by *Lumbriculus variegatus* (Oligochaeta). *Biogeochemistry*, 40(2-3), 267-278. Retrieved from: www.scopus.com
- Nylander, J.A.A., Erséus, C. & Källersjö, M. 1999. A test of monophyly of the gutless Phalodrilinae (Oligochaeta, Tubificidae) and the use of a 573-bp region of the mitochondrial cytochrome oxidase I gene in analysis of annelid phylogeny. *Zoologica Scripta*, 28(3-4), 305-313. doi: [10.1046/j.1463-6409.1999.00001.x](https://doi.org/10.1046/j.1463-6409.1999.00001.x)
- Oda, F.H., Petsch, D.K., Ragonha, F.H., Batista, V.G., Takeda, A.M. & Takemoto, R.M. 2015. Dero (Allodero) Lutzi michaelsen, 1926 (Oligochaeta: Naididae) associated with *Scinax fuscovarius* (Lutz, 1925) (Anura: Hylidae) from semi-deciduous atlantic rain forest, southern Brazil. [Dero (Allodero) Lutzi Michaelsen, 1926 (Oligochaeta: Naididae) associado com *Scinax fuscovarius* (Lutz, 1925) (Anura: Hylidae) na Floresta Estacional Semidecidual, Sul do Brasil]. *Brazilian Journal of Biology*, 75(1), 86-90. doi: [10.1590/1519-6984.07613](https://doi.org/10.1590/1519-6984.07613)
- Odabaşı, S., Odabaşı, D.A. & Arslan, N. 2015. The first record of Chaetogaster limnaei limnaei Baer 1827 (Annelida: Clitellata) on *Pseudobithynia yildirimi* (Gastropoda: Prosobranchia) from northwest of Turkey. *Turkish Journal of Fisheries and Aquatic Sciences*, 15(2), 367-369. doi: [10.4194/1303-2712-v15_2_19](https://doi.org/10.4194/1303-2712-v15_2_19)

- O'Gara, B.A., Bohannon, V.K., Teague, M.W. & Smeaton, M.B. 2004. Copper-induced changes in locomotor behaviors and neuronal physiology of the freshwater oligochaete, *Lumbriculus variegatus*. *Aquatic Toxicology*, 69(1), 51-66. doi: [10.1016/j.aquatox.2004.04.006](https://doi.org/10.1016/j.aquatox.2004.04.006)
- Ohtaka, A. 2004. A new species of the subterranean genus *Krenedrilus Dunnicki* (Oligochaeta, Tubificidae) from the deep bottom of an oligotrophic Caldera Lake in Japan. *Zoological Science*, 21(4), 465-471. doi: [10.2108/zsj.21.465](https://doi.org/10.2108/zsj.21.465)
- Ohtaka, A. & Ichiyanagi, H. 2015. Water environment and species composition of profundal oligochaetes in dam lakes of Japan. *Ecology and Civil Engineering*, 18(2), 87-98. doi: [10.3825/ece.18.87](https://doi.org/10.3825/ece.18.87)
- Ohtaka, A. & Iwakuma, T. 1993. Redescription of *ophidonais serpentina* (Muller, 1773) (Naididae, Oligochaeta) from Lake Yunoko, central Japan, with record of the oligochaete composition in the lake. *Japanese Journal of Limnology (Rikusuirigaku Zasshi)*, 54(4), 251-259. doi: [10.3739/rikusui.54.251](https://doi.org/10.3739/rikusui.54.251)
- Ohtaka, A. & Martin, P. 2011. Redescriptions of *Spiroperma apapillatus* and *Embocephalus nikolskyi* (Annelida, Clitellata, Tubificinae) from Japan, with reference to distribution of Papillate tubificines in Japanese freshwaters. *Zoological Science*, 28(5), 368-379. doi: [10.2108/zsj.28.368](https://doi.org/10.2108/zsj.28.368)
- Ohtaka, A. & Morino, H. 1986. Seasonal changes in the epiphytic animals on the *Potamogeton malaianus* in Lake Kita-Ura, with special reference to oligochaetes. *Japanese Journal of Limnology (Rikusuirigaku Zasshi)*, 47(1), 63-75. doi: [10.3739/rikusui.47.63](https://doi.org/10.3739/rikusui.47.63)
- Ohtaka, A. & Nishino, M. 2012. Diversity and distribution of aquatic oligochaetes (Annelida, Clitellata) in Lake Biwa. *Lake Biwa: Interactions between nature and people*. (pp. 79-83) doi: [10.1007/978-94-007-1783-1_2](https://doi.org/10.1007/978-94-007-1783-1_2). Retrieved from: www.scopus.com
- Ohtaka, A. & Nishino, M. 1995. Studies on the aquatic oligochaete fauna in Lake Biwa, central Japan. I. Checklist with taxonomic remarks. *Japanese Journal of Limnology (Rikusuirigaku Zasshi)*, 56(3), 167-182. doi: [10.3739/rikusui.56.167](https://doi.org/10.3739/rikusui.56.167)
- Ohtaka, A. & Nishino, M. 1999. Studies on the aquatic oligochaete fauna in Lake Biwa, central Japan. II. Records and taxonomic remarks of nine species. *Hydrobiologia*, 406, 33-47. doi: [10.1023/A:100379531154](https://doi.org/10.1023/A:100379531154)
- Ohtaka, A. & Nishino, M. 2006. Studies on the aquatic oligochaete fauna in Lake Biwa, Central Japan. IV. Faunal characteristics in the attached lakes (Naiko). *Limnology*, 7(2), 129-142. doi: [10.1007/s10201-006-0173-1](https://doi.org/10.1007/s10201-006-0173-1)
- Oliver, B.G. 1987. Biouptake of chlorinated hydrocarbons from laboratory-spiked and field sediments by oligochaete worms. *Environmental Science & Technology*, 21(8), 785-790. doi: [10.1021/es00162a009](https://doi.org/10.1021/es00162a009)
- Omodeo, P. 1987. Some new species of Haplotaenidae (Oligochaeta) from guinea and remarks on the history of the family. *Hydrobiologia*, 155(1), 1-13. doi: [10.1007/BF00025626](https://doi.org/10.1007/BF00025626)
- Oyama, A. & Shimizu, T. 2007. Transient occurrence of vasa-expressing cells in nongenital segments during embryonic development in the oligochaete annelid *Tubifex tubifex*. *Development Genes and Evolution*, 217(10), 675-690. doi: [10.1007/s00427-007-0180-1](https://doi.org/10.1007/s00427-007-0180-1)
- Oyama, A., Yoshida, H. & Shimizu, T. 2008. Embryonic expression of p68, a DEAD-box RNA helicase, in the oligochaete annelid *Tubifex tubifex*. *Gene Expression Patterns*, 8(6), 464-470. doi: [10.1016/j.gep.2008.02.003](https://doi.org/10.1016/j.gep.2008.02.003)
- Özer, A. & Wootten, R. 2001. Ultrastructural observations on the development of some actinosporean types within their oligochaete hosts. *Turkish Journal of Zoology*, 25(3), 199-216. Retrieved from: www.scopus.com
- Pasteris, A., Vecchi, M. & Bonomi, G. 1999. A comparison among different population models for *Limnodrilus hoffmeisteri claparede* (Oligochaeta, Tubificidae). *Hydrobiologia*, 406, 183-189. doi: [10.1023/A:1003748520202](https://doi.org/10.1023/A:1003748520202)
- Pasteris, A., Vecchi, M., Reynoldson, T.B. & Bonomi, G. 2003. Toxicity of copper-spiked sediments to *Tubifex tubifex* (Oligochaeta, Tubificidae): A comparison of the 28-day reproductive bioassay with a 6-month cohort experiment. *Aquatic Toxicology*, 65(3), 253-265. doi: [10.1016/S0166-445X\(03\)00136-X](https://doi.org/10.1016/S0166-445X(03)00136-X)
- Pathiratne, A. & Weerasundara, A. 2004. Bioassessment of selected inland water bodies in sri lanka using benthic oligochaetes with consideration of temporal variations. *International Review of Hydrobiology*, 89(3), 305-316. doi: [10.1002/iroh.200310676](https://doi.org/10.1002/iroh.200310676)
- Paumen, M.L., Stol, P., Laak, T.L.T., Kraak, M.H.S., Van Gestel, C.A.M. & Admiraal, W. 2008. Chronic exposure of the oligochaete *Lumbriculus variegatus* to polycyclic aromatic compounds (PACs): Bioavailability and effects on reproduction. *Environmental Science & Technology*, 42(9), 3434-3440. doi: [10.1021/es702500t](https://doi.org/10.1021/es702500t)
- Paunovic, M., Kalafatic, V., Jakovcev, D. & Martinovic-Vitanovic, V. 2003. Oligochaetes (Annelida, Oligochaeta) of the Vlasina River (south-east Serbia): Diversity and distribution. *Biologia – Section Zoology*, 58(5), 903-911. Retrieved from: www.scopus.com
- Pelegrí, S.P. & Blackburn, T.H. 1995. Effects of *Tubifex tubifex* (Oligochaeta: Tubificidae) on N-mineralization in freshwater sediments, measured with ¹⁵N isotopes. *Aquatic Microbial Ecology*, 9(3), 289-294. doi: [10.3354/ame009289](https://doi.org/10.3354/ame009289)
- Penttilinen, O.-., Kilpi-Koski, J., Jokela, M., Toivainen, K. & Väistönen, A. 2008. Importance of dose metrics for lethal and sublethal sediment metal toxicity in the oligochaete worm *Lumbriculus variegatus*. *Journal of Soils and Sediments*, 8(1), 59-66. doi: [10.1065/jss2007.12.267](https://doi.org/10.1065/jss2007.12.267)
- Peralta, L., Escobar, E., Alcocer, J. & Lugo, A. 2002. Oligochaetes from six tropical crater lakes in Central Mexico: Species composition, density and biomass. *Hydrobiologia*, 467, 109-116. doi: [10.1023/A:1014913503577](https://doi.org/10.1023/A:1014913503577)
- Petersen, S., Arlt, G., Faubel, A. & Carman, K.R. 1998. On the nutritive significance of dissolved free amino acid uptake for the cosmopolitan oligochaete *Nais elongata* Müller (Naididae). *Estuarine, Coastal and Shelf Science*, 46(1), 85-91. doi: [10.1006/ecss.1997.0253](https://doi.org/10.1006/ecss.1997.0253)
- Petsch, D.K., Ragonha, F.H., Gimenez, B.C.G., Barboza, L.G.A. & Takeda, A.M. 2015. Partitioning beta diversity of aquatic Oligochaeta in different environments of a neotropical floodplain. [Partição da diversidade beta de Oligochaeta aquático em diferentes ambientes de uma planície de inundação neotropical]. *Acta Scientiarum – Biological Sciences*, 37(1), 41-49. doi: [10.4025/actascibiolsci.v37i1.21738](https://doi.org/10.4025/actascibiolsci.v37i1.21738)
- Phipps, G.L., Ankle, G.T., Benoit, D.A. & Mattson, V.R. 1993. Use of the aquatic oligochaete *Lumbriculus variegatus* for assessing the toxicity and bioaccumulation of sediment-associated contaminants. *Environmental Toxicology and Chemistry*, 12(2), 269-279. doi: [10.1002/etc.5620120210](https://doi.org/10.1002/etc.5620120210)
- Pinder, A.M. 2003. First australian records of three species and two genera of aquatic oligochaetes (Clitellata: Annelida). *Proceedings of the Linnean Society of New South Wales*, 124, 109-114. Retrieved from: www.scopus.com
- Pinder, A.M. & Brinkhurst, R.O. 1997. First records of *slavina* (Oligochaeta: Naididae) in Australia and description of a new species. *Proceedings of the Royal Society of Victoria*, 109(2), 149-155. Retrieved from: www.scopus.com
- Pinder, A.M. & Brinkhurst, R.O. 1997. Review of the Phreodrilidae (Annelida: Oligochaeta: Tubificida) of Australia. *Invertebrate Taxonomy*, 11(3), 443-523. Retrieved from: www.scopus.com
- Pinder, A.M. & Brinkhurst, R.O. 1997. The family capilloventridae (Annelida, Clitellata) in Australia, with descriptions of two new species of capilloventer. *Zoologica Scripta*, 26(3), 255-265. doi: [10.1111/j.1463-6409.1997.tb00415.x](https://doi.org/10.1111/j.1463-6409.1997.tb00415.x)
- Polatdemir Arslan, N. & Şahin, Y. 2003. Nine new Naididae (Oligochaeta) species for Sakarya River, Turkey. *Turkish Journal of Zoology*, 27(1), 27-38. Retrieved from: www.scopus.com
- Prenda, J. & Gallardo, A. 1992. The influence of environmental factors and microhabitat availability on the distribution of an aquatic oligochaete

- assemblage in a Mediterranean River basin. *Internationale Revue Der Gesamten Hydrobiologie Und Hydrographie*, 77(3), 421-434. doi: [10.1002/iroh.19920770306](https://doi.org/10.1002/iroh.19920770306)
- Probst, L. 1987. Sublitoral and profundal Oligochaeta fauna of the Lake Constance (Bodensee-Obersee). *Hydrobiologia*, 155(1), 277-282. doi: [10.1007/BF00025660](https://doi.org/10.1007/BF00025660)
- Pujals, M.A. 1988. Especies del género Dero Oken, 1815 (Oligochaeta, Naididae) en la provincia de Buenos Aires, Argentina. *Studies on Neotropical Fauna and Environment*, 23(4), 203-212. doi: [10.1080/01650528809360764](https://doi.org/10.1080/01650528809360764)
- Purschke, G. 1999. Terrestrial polychaetes – models for the evolution of the Clitellata (Annelida)? *Hydrobiologia*, 406, 87-99. doi: [10.1023/A:1003780032497](https://doi.org/10.1023/A:1003780032497)
- Putzer, V., Gnaiger, E. & Lackner, R. 1985. Flexibility of anaerobic metabolism in aquatic oligochaetes (*Tubifex* sp.). Biochemical and calorimetric changes induced by a deproteinized hydrolysate of bovine blood. *Comparative Biochemistry and Physiology – Part A: Physiology*, 82(4), 965-970. doi: [10.1016/0300-9629\(85\)90513-4](https://doi.org/10.1016/0300-9629(85)90513-4)
- Putzer, V.M., de Zwaan, A. & Wieser, W. 1990. Anaerobic energy metabolism in the oligochaete *Lumbriculus variegatus* Müller. *Journal of Comparative Physiology B*, 159(6), 707-715. doi: [10.1007/BF00691716](https://doi.org/10.1007/BF00691716)
- Qi, S. 1987. Some ecological aspects of aquatic oligochaetes in the lower Pearl River (people's Republic of China). *Hydrobiologia*, 155(1), 199-208. doi: [10.1007/BF00025652](https://doi.org/10.1007/BF00025652)
- Rácz, O.Z., Eszterbauer, E. & Molnár, K. 2005. Hungactinomyxon, a new actinosporean type and collective group (Myxozoa) from Branchiura sowerbyi Beddard (Oligochaeta). *Systematic Parasitology*, 61(2), 107-113. doi: [10.1007/s11230-005-3136-8](https://doi.org/10.1007/s11230-005-3136-8)
- Ragonha, F.H., Chiaramonte, J.B., Junior, H.M.F., da Cunha, E.R., Benedito, E. & Takeda, A.M. 2013. Spatial distribution of aquatic Oligochaeta in Ilha Grande National Park, Brazil. [Distribuição espacial de Oligochaeta aquático no Parque Nacional de Ilha Grande, Brasil]. *Acta Scientiarum – Biological Sciences*, 35(1), 63-70. doi: [10.4025/actascibiolsci.v35i1.10768](https://doi.org/10.4025/actascibiolsci.v35i1.10768)
- Ragonha, F.H., Petsch, D.K., Alves, G.H.Z., Santana, H.S., Michelan, T.S. & Takeda, A.M. 2014. Tributaries as richness source for Oligochaeta assemblage (Annelida) of Neotropical dammed river. [Tributários como fontes de riqueza para assembléia de Oligochaeta (Annelida) de um rio Neotropical represado]. *Brazilian Journal of Biology*, 74(4), 861-869. doi: [10.1590/1519-6984.05613](https://doi.org/10.1590/1519-6984.05613)
- Ragonha, F.H. & Takeda, A.M. 2014. Does richness of Oligochaeta (Annelida) follows a linear distribution with habitat structural heterogeneity in aquatic sediments? *Journal of Limnology*, 73(1), 146-156. doi: [10.4081/jlimnol.2014.791](https://doi.org/10.4081/jlimnol.2014.791)
- Ramskov, T., Thit, A., Croteau, M.-. & Selck, H. 2015. Biodynamics of copper oxide nanoparticles and copper ions in an oligochaete – Part I: Relative importance of water and sediment as exposure routes. *Aquatic Toxicology*, 164, 81-91. doi: [10.1016/j.aquatox.2015.04.022](https://doi.org/10.1016/j.aquatox.2015.04.022)
- Raposeiro, P.M., Ramos, J.C. & Costa, A.C. 2009. First record of Branchiura sowerbyi Beddard, 1892 (Oligochaeta: Tubificidae) in azores. *Aquatic Invasions*, 4(3), 487-490. doi: [10.3391/ai.2009.4.3.8](https://doi.org/10.3391/ai.2009.4.3.8)
- Ratsak, C.H., Koouman, S.A.L.M. & Kooi, B.W. 1993. Modelling the growth of an oligochaete on activated sludge. *Water Research*, 27(5), 739-747. doi: [10.1016/0043-1354\(93\)90136-6](https://doi.org/10.1016/0043-1354(93)90136-6)
- Ratsak, C.H. & Verkuijlen, J. 2006. Sludge reduction by predatory activity of aquatic oligochaetes in wastewater treatment plants: Science or fiction? A review. *Hydrobiologia*, 564 (1 spec. iss.), 197-211. doi: [10.1007/s10750-005-1719-7](https://doi.org/10.1007/s10750-005-1719-7)
- Real, M. & Prat, N. 1992. Factors influencing the distribution of chironomids and oligochaetes in profundal areas of spanish reservoirs. *Netherlands Journal of Aquatic Ecology*, 26(2-4), 405-410. doi: [10.1007/BF02255269](https://doi.org/10.1007/BF02255269)
- Real, M., Sabater, F. & Morguí, J.A. 1992. Significant physiographic disturbances in the ebro basin reservoirs (N.E. Spain) reflected by Oligochaeta size spectra. *Hydrobiologia*, 235-236(1), 363-374. doi: [10.1007/BF00026226](https://doi.org/10.1007/BF00026226)
- Rech, K.C., Guereschi, R.M., de Oliveira Nuñez, A.P. & Sticca, S.C. 2013. Density and organic matter influence on egg-laying and hatching of Branchiura sowerbyi (Oligochaeta). *Journal of the World Aquaculture Society*, 44(2), 267-272. doi: [10.1111/jwas.12026](https://doi.org/10.1111/jwas.12026)
- Redeker, E.S., Bervoets, L. & Blust, R. 2004. Dynamic model for the accumulation of cadmium and zinc from water and sediment by the aquatic oligochaete, *Tubifex tubifex*. *Environmental Science & Technology*, 38(23), 6193-6200. doi: [10.1021/es0496470](https://doi.org/10.1021/es0496470)
- Redeker, E.S. & Blust, R. 2004. Accumulation and toxicity of cadmium in the aquatic oligochaete *Tubifex tubifex*: A kinetic modeling approach. *Environmental Science & Technology*, 38(2), 537-543. doi: [10.1021/es0343858](https://doi.org/10.1021/es0343858)
- Reible, D.D., Popov, V., Valsaraj, K.T., Thibodeaux, L.J., Lin, F., Dikshit, M., ... Fleeger, J.W. 1996. Contaminant fluxes from sediment due to tubificid oligochaete bioturbation. *Water Research*, 30(3), 704-714. doi: [10.1016/0043-1354\(95\)00187-5](https://doi.org/10.1016/0043-1354(95)00187-5)
- Reynoldson, T.B. 1994. A field test of a sediment bioassay with the oligochaete worm *Tubifex tubifex* (Müller, 1774). *Hydrobiologia*, 278(1-3), 223-230. doi: [10.1007/BF00142330](https://doi.org/10.1007/BF00142330)
- Reynoldson, T.B. 1987. The role of environmental factors in the ecology of tubificid oligochaetes - an experimental study. *Ecography*, 10(4), 241-248. doi: [10.1111/j.1600-0587.1987.tb00765.x](https://doi.org/10.1111/j.1600-0587.1987.tb00765.x)
- Reynoldson, T.B., Thompson, S.P. & Bamsey, J.L. 1991. A sediment bioassay using the tubificid oligochaete worm *Tubifex tubifex*. *Environmental Toxicology and Chemistry*, 10(8), 1061-1072. doi: [10.1002/etc.5620100811](https://doi.org/10.1002/etc.5620100811)
- Rieradevall, M. & Real, M. 1994. On the distribution patterns and population dynamics of sublitoral and profundal Oligochaeta fauna from Lake Banyoles (Catalonia, NE Spain). *Hydrobiologia*, 278(1-3), 139-149. doi: [10.1007/BF00142322](https://doi.org/10.1007/BF00142322)
- Rodrigues, L.F.T., Leite, F.S. & Alves, R.G. 2013. Inventory and distribution of Oligochaeta (Annelida, Clitellata) in first-order streams in preserved areas of the state of Minas Gerais, Brazil. [Inventário e distribuição de Oligochaeta (Annelida, Clitellata) em córregos de primeira ordem localizados em áreas preservadas do estado de Minas Gerais, Brasil]. *Biota Neotropica*, 13(1), 245-254. doi: [10.1590/S1676-06032013000100024](https://doi.org/10.1590/S1676-06032013000100024)
- Rodrigues, L.F.T., Santana, L.D. & Alves, R.G. 2013. Aquatic oligochaetes associated with bryophytes in an atlantic forest stream. [Oligoquetas aquáticos associados a briófitas em um córrego de Mata Atlântica]. *Biota Neotropica*, 13(4), 371-375. doi: [10.1590/S1676-06032013000400031](https://doi.org/10.1590/S1676-06032013000400031)
- Rodriguez, P. 1988. On some species of Lumbriculidae (Annelida: Oligochaeta) from the north of the Iberian Peninsula. [Sur certaines espèces de Lumbriculidae (Annelida: Oligochaeta) du nord de la péninsule ibérique]. *Annales de Limnologie*, 24(3), 203-211. doi: [10.1051/limn/1988018](https://doi.org/10.1051/limn/1988018)
- Rodriguez, P. 1996. *Stylodrilus californianus* sp. nov., a new Lumbriculid (Annelida: Oligochaeta) from North America. *Hydrobiologia*, 333(3), 161-164. doi: [10.1007/BF00013430](https://doi.org/10.1007/BF00013430)
- Rodriguez, P. 1987. The variability of setae of *Pristina longiseta* Ehrenberg (Oligochaeta, Naididae). *Hydrobiologia*, 155(1), 39-44. doi: [10.1007/BF00025629](https://doi.org/10.1007/BF00025629)
- Rodriguez, P. & Achurra, A. 2010. New species of aquatic oligochaetes (Annelida: Clitellata) from groundwaters in karstic areas of northern Spain, with taxonomic remarks on *Lophochaeta ignota* Štolc, 1886. *Zootaxa*, (2332), 21-39. Retrieved from: www.scopus.com
- Rodriguez, P., Arrate, J., Martinez-Madrid, M., Reynoldson, T.B., Schumacher, V. & Viguri, J. 2006. Toxicity of santander bay sediments to the euryhaline

- freshwater oligochaete Limnodrilus hoffmeisteri. *Hydrobiologia*, 564 (1 spec. iss.), 157-169. doi: [10.1109/MAP.2006.1650857](https://doi.org/10.1109/MAP.2006.1650857)
- Rodriguez, P. & Coates, K.A. 1996. A new american Stylodrilus species (Lumbriculidae, Oligochaeta). *Canadian Journal of Zoology*, 74(1), 92-96. doi: [10.1139/z96-012](https://doi.org/10.1139/z96-012)
- Rodriguez, P. & Fend, S.V. 2013. New species of Rhyacodrilus (Annelida: Clitellata: Rhyacodrilinae) of North America, with re-description of *R. sodalis* (Eisen, 1879). *Zootaxa*, 3664(1), 1-44. doi: [10.11646/zootaxa.3664.1.1](https://doi.org/10.11646/zootaxa.3664.1.1)
- Rodriguez, P., Fend, S.V. & Lenat, D.R. 2014. Sylphella pucoo gen. nov., sp. nov., and two additional new species of aquatic oligochaetes (Lumbriculidae, Clitellata) from poorly-known lotic habitats in north carolina (USA). *ZooKeys*, (451), 1-32. doi: [10.3897/zookeys.451.7304](https://doi.org/10.3897/zookeys.451.7304)
- Rodriguez, P. & Giani, N. 1994. A preliminary review of the taxonomic characters used for the systematics of the genus Trichodrilus Claparède (Oligochaeta, Lumbriculidae). *Hydrobiologia*, 278(1-3), 35-51. doi: [10.1007/BF00142310](https://doi.org/10.1007/BF00142310)
- Rodriguez, P. & Giani, N. 1989. New species of Phalodrilus (Oligochaeta, Tubificidae) from caves of northern Spain and southwestern France. *Hydrobiologia*, 180(1), 57-63. doi: [10.1007/BF00027537](https://doi.org/10.1007/BF00027537)
- Rodriguez, P. & Rico, E. 2008. A new freshwater oligochaete species (Clitellata: Enchytraeidae) from Livingston Island, Antarctica. *Polar Biology*, 31(10), 1267-1279. doi: [10.1007/s00300-008-0465-5](https://doi.org/10.1007/s00300-008-0465-5)
- Rosa, B.F.V., Martins, R.T. & Alves, R.G. 2015. Distribution of oligochaetes in a stream in the atlantic forest in southeastern Brazil. [Distribuição de oligoquetas em um riacho da mata atlântica, sudeste do Brasil]. *Brazilian Journal of Biology*, 75(1), 1-7. doi: [10.1590/1519-6984.02313](https://doi.org/10.1590/1519-6984.02313)
- Rosa, B.F.V., Rodrigues, L.F.T., de Oliveira, G.S. & da Gama Alves, R. 2014. Chironomidae and Oligochaeta for water quality evaluation in an urban river in southeastern Brazil. *Environmental Monitoring and Assessment*, 186(11), 7771-7779. doi: [10.1007/s10661-014-3965-5](https://doi.org/10.1007/s10661-014-3965-5)
- Rosso, A., Lafont, M. & Exinger, A. 1993. Effects of heavy metals on oligochaete communities of the river ill and its tributaries (Haut-Rhin, France). [Effets des métaux lourds sur les peuplements d'oligochètes de l'Ill et de ses affluents (Haut-Rhin, France)]. *Annales de Limnologie*, 29(3), 295-305. doi: [10.1051/limn/1993024](https://doi.org/10.1051/limn/1993024)
- Rota, E. & Manconi, R. 2004. Taxonomy and ecology of sponge-associate Marionina sp. (Clitellata: Enchytraeidae) from the horomatangi geothermal system of Lake Taupo, New Zealand. *International Review of Hydrobiologia*, 89(1), 58-67. doi: [10.1002/iroh.200310695](https://doi.org/10.1002/iroh.200310695)
- Růžičková, S., Schenková, J., Weissová, V. & Helešic, J. 2014. Environmental impact of heated mining waters on clitellate (Annelida: Clitellata) assemblages. *Biologia (Poland)*, 69(9), 1179-1189. doi: [10.2478/s11756-014-0424-2](https://doi.org/10.2478/s11756-014-0424-2)
- Sager, M. & Pucsko, R. 1991. Trace element concentrations of oligochaetes and relations to sediment characteristics in the reservoir at Altenwörth, Austria. *Hydrobiologia*, 226(1), 39-49. doi: [10.1007/BF00007778](https://doi.org/10.1007/BF00007778)
- Şahin, S.K. & Yıldız, S. 2011. Species distribution of oligochaetes related to environmental parameters in Lake Sapanca (Marmara Region, Turkey). [Oligoket türlerinin Sapanca gölünde (Marmara Bölgesi, Türkiye) çevresel parametrelerle ilişkili dağılımının mikrolojisi]. *Turkish Journal of Fisheries and Aquatic Sciences*, 11(3), 359-366. doi: [10.4194/1303-2712-v11_3_04](https://doi.org/10.4194/1303-2712-v11_3_04)
- Sang, Q. & Erséus, C. 1985. Ecological survey of the aquatic oligochaetes in the lower pearl river (people's Republic of China). *Hydrobiologia*, 128(1), 39-44. doi: [10.1007/BF00008938](https://doi.org/10.1007/BF00008938)
- Sardo, A.M. & Soares, A.M.V.M. 2010. Can behavioural responses of *Lumbriculus variegatus* (Oligochaeta) assess sediment toxicity? A case study with sediments exposed to acid mine drainage. *Environmental Pollution*, 158(2), 636-640. doi: [10.1016/j.envpol.2009.08.015](https://doi.org/10.1016/j.envpol.2009.08.015)
- Sardo, A.M. & Soares, A.M.V.M. 2011. Short- and long-term exposure of *Lumbriculus variegatus* (Oligochaete) to metal lead: Ecotoxicological and behavioral effects. *Human and Ecological Risk Assessment*, 17(5), 1108-1123. doi: [10.1080/10807039.2011.605721](https://doi.org/10.1080/10807039.2011.605721)
- Sardo, A.M., Soares, A.M.V.M. & Gerhardt, A. 2007. Behavior, growth, and reproduction of *Lumbriculus variegatus* (Oligochaetae) in different sediment types. *Human and Ecological Risk Assessment*, 13(3), 519-526. doi: [10.1080/10807030701341043](https://doi.org/10.1080/10807030701341043)
- Särkkä, J. 1994. Lacustrine, profundal meiobenthic oligochaetes as indicators of trophy and organic loading. *Hydrobiologia*, 278(1-3), 231-241. doi: [10.1007/BF00142331](https://doi.org/10.1007/BF00142331)
- Särkkä, J. 1989. Meiobenthic naidid and aeolosomatid oligochaetes from the profundal zone, and relations of species to eutrophication. *Hydrobiologia*, 180(1), 185-190. doi: [10.1007/BF00027551](https://doi.org/10.1007/BF00027551)
- Särkkä, J. 1987. The occurrence of oligochaetes in lake chains receiving pulp mill waste and their relation to eutrophication on the trophic scale. *Hydrobiologia*, 155(1), 259-266. doi: [10.1007/BF00025658](https://doi.org/10.1007/BF00025658)
- Schenková, J., Helešic, J. & Jarkovský, J. 2006. Seasonal dynamics of *Bythonomus lemani* and *Bothrioneurum vejvodskyanum* (Oligochaeta, Annelida) in relation to environmental variables. *Biologia*, 61(5), 517-523. Retrieved from: www.scopus.com
- Schenková, J., Komárek, O. & Zahrádková, S. 2001. Oligochaeta of the morava and odra river basins (Czech Republic): Species distribution and community composition. *Hydrobiologia*, 463, 235-240. doi: [10.1023/A:1013128413701](https://doi.org/10.1023/A:1013128413701)
- Schenková, J., Paøil, P., Petøivalská, K. & Bojková, J. 2010. Aquatic oligochaetes (Annelida: Clitellata) of the Czech Republic: Check-list, new records, and ecological remarks. *Zootaxa*, (2676), 29-44. Retrieved from: www.scopus.com
- Schierwater, B. & Hauenschild, C. 1990. A photoperiod determined life-cycle in an oligochaete worm. *Biological Bulletin*, 178(2), 111-117. doi: [10.2307/1541969](https://doi.org/10.2307/1541969)
- Schlegel, M., Steinbrück, G., Kramer, M. & Brockmeyer, V. 1991. Restriction fragment patterns as molecular markers for species identification and phylogenetic analysis in the genus *Enchytraeus* (Oligochaeta). *Journal of Zoological Systematics and Evolutionary Research*, 29(5-6), 362-372. doi: [10.1111/j.1439-0469.1991.tb00458.x](https://doi.org/10.1111/j.1439-0469.1991.tb00458.x)
- Schloesser, D.W., Reynoldson, T.B. & Manny, B.A. 1995. Oligochaete fauna of western Lake Erie 1961 and 1982: Signs of sediment quality recovery. *Journal of Great Lakes Research*, 21(3), 294-306. doi: [10.1016/S0380-1330\(95\)71040-1](https://doi.org/10.1016/S0380-1330(95)71040-1)
- Schmelz, R.M. 1994. Oligochaetes of the River Rhine. Preliminary records. *Hydrobiologia*, 278(1-3), 85-86. doi: [10.1007/BF00142314](https://doi.org/10.1007/BF00142314)
- Schmelz, R.M., Jocque, M. & Collado, R. 2015. Microdrile Oligochaeta in bromeliad pools of a honduran cloud forest. *Zootaxa*, 3947(4), 508-526. doi: [10.1164/zootaxa.3947.4.3](https://doi.org/10.1164/zootaxa.3947.4.3)
- Schmelz, R.M. & Westheide, W. 2000. Ultrastructure of oesophageal appendages ("peptonephridia") in enchytraeids (Annelida: Clitellata). *Invertebrate Biology*, 119(1), 94-103. Retrieved from: www.scopus.com
- Schuytema, G.S., Krawczyk, D.F., Grifflis, W.L., Nebeker, A.V., Robideaux, M.L., Brownawell, B.J. & Westall, J.C. 1988. Comparative uptake of hexachlorobenzene by fathead minnows, amphipods and oligochaete worms from water and sediment. *Environmental Toxicology and Chemistry*, 7(12), 1035-1045. doi: [10.1002/etc.5620071211](https://doi.org/10.1002/etc.5620071211)
- Sedlmeier, U.A. & Hoffmann, K.H. 1989. Integumentary uptake of short-chain carboxylic acids by two freshwater oligochaetes, *Tubifex tubifex* and *Lumbriculus variegatus*: Specificity of uptake and characterization of transport carrier. *Journal of Experimental Zoology*, 250(2), 128-134. doi: [10.1002/jez.1402500203](https://doi.org/10.1002/jez.1402500203)

- Shaw, R. 1992. Host tracking and photosensitivity in *Chaetogaster limnaei* (Oligochaeta). *Canadian Journal of Zoology*, 70(3), 578-581. doi: [10.1139/z92-086](https://doi.org/10.1139/z92-086)
- Sheedy, B.R., Mattson, V.R., Cox, J.S., Kosian, P.A., Phipps, G.L. & Ankley, G.T. 1998. Bioconcentration of polycyclic aromatic hydrocarbons by the freshwater oligochaete *Lumbriculus variegatus*. *Chemosphere*, 36(15), 3061-3070. doi: [10.1016/S0045-6535\(98\)00007-1](https://doi.org/10.1016/S0045-6535(98)00007-1)
- Shimizu, T., Nakamoto, A. & Nakamoto, A. 2014. Developmental significance of D quadrant micromeres 2d and 4d in the oligochaete annelid *Tubifex tubifex*. *International Journal of Developmental Biology*, 58(6-8), 445-456. doi: [10.1387/ijdb.140102tsh](https://doi.org/10.1387/ijdb.140102tsh)
- Shimizu, T. & Savage, R.M. 2002. Expression of hunchback protein in a subset of ectodermal teloblasts of the oligochaete annelid *Tubifex*. *Development Genes and Evolution*, 212(11), 520-525. doi: [10.1007/s00427-002-0278-4](https://doi.org/10.1007/s00427-002-0278-4)
- Shirakashi, S. & El-Matbouli, M. 2009. *Myxobolus cerebralis* (Myxozoa), the causative agent of whirling disease, reduces fecundity and feeding activity of *Tubifex tubifex* (Oligochaeta). *Parasitology*, 136(6), 603-613. doi: [10.1017/S0031182009005721](https://doi.org/10.1017/S0031182009005721)
- Simpson, I.C., Roger, P.A., Oficial, R. & Grant, I.F. 1993. Density and composition of aquatic oligochaete populations in different farmers' ricefields. *Biology and Fertility of Soils*, 16(1), 34-40. doi: [10.1007/BF00336512](https://doi.org/10.1007/BF00336512)
- Simpson, I.C., Roger, P.A., Oficial, R. & Grant, I.F. 1993. Impacts of agricultural practices on aquatic oligochaete populations in ricefields. *Biology and Fertility of Soils*, 16(1), 27-33. doi: [10.1007/BF00336511](https://doi.org/10.1007/BF00336511)
- Smit, H., van Heel, E.D. & Wiersma, S. 1993. Biovolume as a tool in biomass determination of Oligochaeta and Chironomidae. *Freshwater Biology*, 29(1), 37-46. doi: [10.1111/j.1365-2427.1993.tb00742.x](https://doi.org/10.1111/j.1365-2427.1993.tb00742.x)
- Smith, D.P., Kennedy, J.H. & Dickson, K.L. 1991. An evaluation of a naidid oligochaete as a toxicity test organism. *Environmental Toxicology and Chemistry*, 10(11), 1459-1465. doi: [10.1002/etc.5620101111](https://doi.org/10.1002/etc.5620101111)
- Smith, M.E. 1986. Ecology of Naididae (Oligochaeta) from an alkaline bog stream: Life history patterns and community structure. *Hydrobiologia*, 133(1), 79-90. doi: [10.1007/BF00010805](https://doi.org/10.1007/BF00010805)
- Smith, M.E. 1985. Population and reproductive dynamics of *Nais communis* (Oligochaeta: Naididae) from a Wisconsin limnophore. *American Midland Naturalist*, 114(1), 152-158. doi: [10.2307/2425250](https://doi.org/10.2307/2425250)
- Smith, M.E. & Kaster, J.L. 1986. Feeding habits and dietary overlap of Naididae (Oligochaeta) from a bog stream. *Hydrobiologia*, 137(3), 193-201. doi: [10.1007/BF00004233](https://doi.org/10.1007/BF00004233)
- Smythe, A.B., Forgrave, K., Patti, A., Hochberg, R. & Litvaitis, M.K. 2015. Untangling the ecology, taxonomy, and evolution of *Chaetogaster limnaei* (Oligochaeta: Naididae) species complex. *Journal of Parasitology*, 101(3), 320-326. doi: [10.1645/13-465.1](https://doi.org/10.1645/13-465.1)
- Snimshchikova, L.N. & Akinshina, T.W. 1994. Oligochaete fauna of Lake Baikal. *Hydrobiologia*, 278(1-3), 27-34. doi: [10.1007/BF00142309](https://doi.org/10.1007/BF00142309)
- Snimshchikova, L.N. 1998. Revision of the genus *lycodrilus* grube and description of the evolution of *Limnodrilus clavarede* (Oligochaeta, Tubificidae) species in Lake Baikal. 2. Revision of the species *lycodrilus grubei* with discussion of the current taxonomic position of *Limnodrilus* species from Lake Baikal and their evolution. *Zoologicheskii Zhurnal*, 77(6), 646-XII. Retrieved from: www.scopus.com
- Snimshchikova, L.N. & Akinshina, T.V. 1995. *Rhyacodrilus stephensi* (Oligochaeta, Tubificidae) from Lake Baikal. *Hydrobiological Journal*, 31(4), 70-76. Retrieved from: www.scopus.com
- Snitgen, J.L. & Sibley, P.K. 1998. New record for the oligochaete amphichaeta americana (Oligochaeta: Naididae) in Minnesota. *Journal of Freshwater Ecology*, 13(3), 369-370. doi: [10.1080/02705060.1998.9663631](https://doi.org/10.1080/02705060.1998.9663631)
- Snow, P. & Buss, L.W. 1994. HOM/Hox-type homeoboxes from *Stylaria lacustris* (Annelida: Oligochaeta). *Molecular Phylogenetics and Evolution*, 3(4), 360-364. doi: [10.1006/mpev.1994.1042](https://doi.org/10.1006/mpev.1994.1042)
- Sormunen, A.J., Leppänen, M.T. & Kukkonen, J.V.K. 2008. Influence of sediment ingestion and exposure concentration on the bioavailable fraction of sediment-associated tetrachlorobiphenyl in oligochaetes. *Environmental Toxicology and Chemistry*, 27(4), 854-863. doi: [10.1897/07-334.1](https://doi.org/10.1897/07-334.1)
- Soster, F.M., Harvey, D.T., Troksa, M.R. & Grooms, T. 1992. The effect of tubificid oligochaetes on the uptake of zinc by Lake Erie sediments. *Hydrobiologia*, 248(3), 249-258. doi: [10.1007/BF00006151](https://doi.org/10.1007/BF00006151)
- Soster, F.M. & McCall, P.L. 1989. Predation by coelotanypus (Diptera: Chironomidae) on laboratory populations of Tubificid oligochaetes. *Journal of Great Lakes Research*, 15(3), 408-417. doi: [10.1016/S0380-1330\(89\)71496-9](https://doi.org/10.1016/S0380-1330(89)71496-9)
- Spencer, D.R. & Denton, R.L. 2003. Aquatic Oligochaeta (Annelida: Lumbriculidae, Haplodtaxidae, Naididae, Tubificidae) of Utah. *Western North American Naturalist*, 63(3), 343-352. Retrieved from: www.scopus.com
- Spencer, D.R. & Hudson, P.L. 2003. The Oligochaeta (Annelida, Clitellata) of the St. Lawrence great lakes region: An update. *Journal of Great Lakes Research*, 29(1), 89-104. doi: [10.1016/S0380-1330\(03\)70418-3](https://doi.org/10.1016/S0380-1330(03)70418-3)
- Sporka, F. 1996. First record of *Aulodrilus pigueti* (Oligochaeta, Tubificidae) from the River Morava in Slovakia. *Biologia*, 51(5), 494. Retrieved from: www.scopus.com
- Sporka, F. 1998. The typology of floodplain water bodies of the middle Danube (Slovakia) on the basis of the superficial polychaete and oligochaete fauna. *Hydrobiologia*, 386(1-3), 55-62. Retrieved from: www.scopus.com
- Standley, L.J. 1997. Effect of sedimentary organic matter composition on the partitioning and bioavailability of dieldrin to the oligochaete *Lumbriculus variegatus*. *Environmental Science & Technology*, 31(9), 2577-2583. doi: [10.1021/es961005s](https://doi.org/10.1021/es961005s)
- Steinbach Elwell, L.C., Kerans, B.L., Rasmussen, C. & Winton, J.R. 2006. Interactions among two strains of *Tubifex tubifex* (Oligochaeta: Tubificidae) and *Myxobolus cerebralis* (Myxozoa). *Diseases of Aquatic Organisms*, 68(2), 131-139. doi: [10.3354/dao068131](https://doi.org/10.3354/dao068131)
- Steinhagen, D. & Korting, W. 1990. The role of tubificid oligochaetes in the transmission of *Goussia carpelli*. *Journal of Parasitology*, 76(1), 104-107. doi: [10.2307/3282636](https://doi.org/10.2307/3282636)
- Steinlechner, R. 1987. Identification of immature Tubificids (Oligochaeta) of Lake Constance and its influence on the evaluation of species distribution. *Hydrobiologia*, 155(1), 57-63. doi: [10.1007/BF00025631](https://doi.org/10.1007/BF00025631)
- Stern, M.S., Vinogradov, S.N., Sharma, P.K., Ereifej, K. & Walz, D.A. 1990. Amino acid sequence of the monomer subunit of the giant extracellular hemoglobin of the aquatic oligochaete, *Tubifex tubifex*. *European Journal of Biochemistry*, 194(1), 67-73. doi: [10.1111/j.1432-1033.1990.tb19428.x](https://doi.org/10.1111/j.1432-1033.1990.tb19428.x)
- Stoll, S., Früh, D., Westerwald, B., Hormel, N. & Haase, P. 2013. Density-dependent relationship between *Chaetogaster limnaei* (Oligochaeta) and the freshwater snail *Physa acuta* (Pulmonata). *Freshwater Science*, 32(2), 642-649. doi: [10.1899/12-072.1](https://doi.org/10.1899/12-072.1)
- Strayer, D. & Bannon-O'donnell, E. 1988. Aquatic microannelids (Oligochaeta and Aphanoneura) of underground waters of southeastern New York. *American Midland Naturalist*, 119(2), 327-335. doi: [10.2307/2425815](https://doi.org/10.2307/2425815)
- Strayer, D.L. 2001. Ecology and distribution of hyporheic microannelids (Oligochaeta, Aphanoneura, and Polychaeta) from the eastern United States. *Archiv Fur Hydrobiologie*, 151(3), 493-510. Retrieved from: www.scopus.com
- Struck, T., Hessling, R. & Purschke, G. 2002. The phylogenetic position of the aeolosomatidae and parergodrilidae, two enigmatic oligochaete-like taxa of the 'polychaeta', based on molecular data from 18S rDNA sequences. *Journal of Zoological Systematics and Evolutionary Research*, 40(3), 155-163. doi: [10.1046/j.1439-0469.2002.00200.x](https://doi.org/10.1046/j.1439-0469.2002.00200.x)

- Sturmbauer, C., Opadiya, G.B., Niederstätter, H., Riedmann, A. & Dallinger, R. 1999. Mitochondrial DNA reveals cryptic oligochaete species differing in cadmium resistance. *Molecular Biology and Evolution*, 16(7), 967-974. doi: [10.1093/oxfordjournals.molbev.a026186](https://doi.org/10.1093/oxfordjournals.molbev.a026186)
- Šundic, D., Radujkovic, B.M. & Krpo-Cetkovic, J. 2011. Catalogue of Naidinae and Pristininae (Annelida: Oligochaeta: Naididae) with twenty species new for Montenegro. *Zootaxa*, (2737), 1-18. Retrieved from: www.scopus.com
- Surianni-Affonso, A.L., França, R.S., Marchese, M. & Rocha, O. 2011. Environmental factors and benthic Oligochaeta (Annelida, Clitellata) assemblages in a stretch of the upper São Francisco River (Minas Gerais state, Brazil). [Fatores ambientais e associações de Oligochaeta (Annelida, Clitellata) em um trecho do alto Rio São Francisco (Minas Gerais, Brasil)]. *Brazilian Journal of Biology*, 71(2), 437-446. Retrieved from: www.scopus.com
- Svein-Erik, S.-. 1994. Oligochaete response to changes in water flow in the Dokka Delta, Lake Randsfjorden (Norway), caused by hydroelectric power development. *Hydrobiologia*, 278(1-3), 243-249. doi: [10.1007/BF00142332](https://doi.org/10.1007/BF00142332)
- Svensson, J.M., Enrich-Prast, A. & Leonardson, L. 2001. Nitrification and denitrification in a eutrophic lake sediment bioturbated by oligochaetes. *Aquatic Microbial Ecology*, 23(2), 177-186. doi: [10.3354/ame023177](https://doi.org/10.3354/ame023177)
- Swayne, H., Day, M. & Wetzel, M.J. 2004. Limnodrilus hoffmeisteri (Annelida: Oligochaeta: Tubificidae) in pop's cave, wisconsin, USA. *Journal of Cave and Karst Studies*, 66(1), 28-31. Retrieved from: www.scopus.com
- Syrovátková, V., Schenková, J. & Brabec, K. 2009. The distribution of chironomid larvae and oligochaetes within a stony-bottomed river stretch: The role of substrate and hydraulic characteristics. *Fundamental and Applied Limnology*, 174(1), 43-62. doi: [10.1127/1863-9135/2009/0174-0043](https://doi.org/10.1127/1863-9135/2009/0174-0043)
- Szarowska, M. 1987. Annual life cycle of Limnodrilus udekemianus (Oligochaeta, Tubificidae) in the River Biala Przemsza (southern Poland). *Acta Hydrobiologica*, 29(1), 53-69. Retrieved from: www.scopus.com
- Székely, C., Molnár, K., Eszterbauer, E. & Baska, F. 1999. Experimental detection of the actinospores of Myxobolus pseudodispar (Myxosporea: Myxobolidae) in oligochaete alternate hosts. *Diseases of Aquatic Organisms*, 38(3), 219-224. doi: [10.3354/dao038219](https://doi.org/10.3354/dao038219)
- Székely, C., Rácz, O., Molnár, K. & Eszterbauer, E. 2002. Development of Myxobolus macrocapsularis (Myxosporea: Myxobolidae) in an oligochaete alternate host, *Tubifex tubifex*. *Diseases of Aquatic Organisms*, 48(2), 117-123. Retrieved from: www.scopus.com
- Székely, C., Sitjà-Bobadilla, A. & Alvarez-Pellitero, P. 2000. First report on the occurrence of an actinosporean stage (Myxozoa) in oligochaetes from spanish freshwaters. *Acta Veterinaria Hungarica*, 48(4), 433-441. doi: [10.1556/004.48.2000.4.6](https://doi.org/10.1556/004.48.2000.4.6)
- Takada, K., Kato, K. & Okino, T. 1992. Environmental parameters and estivation of *Rhyacodrilus* (Tubificidae, Oligochaeta) in Lake Suwa, Japan. *Ecography*, 15(3), 328-333. doi: [10.1111/j.1600-0587.1992.tb00043.x](https://doi.org/10.1111/j.1600-0587.1992.tb00043.x)
- Takahashi, H., Shimizu, T. & Aruga, J. 2008. Expression pattern of annelid zic in embryonic development of the oligochaete *Tubifex tubifex*. *Development Genes and Evolution*, 218(10), 553-560. doi: [10.1007/s00427-008-0252-x](https://doi.org/10.1007/s00427-008-0252-x)
- Takeda, A.M. 1999. Oligochaete community of alluvial upper Paraná River, Brazil: Spatial and temporal distribution (1987-1988). *Hydrobiologia*, 412, 35-42. doi: [10.1023/A:1003844131148](https://doi.org/10.1023/A:1003844131148)
- Taş, M., Kirgiz, T. & Arslan, N. 2011. Dynamics of Oligochaeta fauna in sazlidere stream (Edirne, Turkey) with relation to environmental factors. *Acta Zoologica Bulgarica*, 63(2), 179-185. Retrieved from: www.scopus.com
- Timm, H. 1994. Production of some oligochaete species in large estonian lakes. *Hydrobiologia*, 278(1-3), 309-313. doi: [10.1007/BF00142340](https://doi.org/10.1007/BF00142340)
- Timm, T. 2006. A nearctic tubificid *Varichaetadrilus harmani* (Loden) comb. nov., in a dutch wetland, with remarks on *Tubifex tubifex* (Müller) (Annelida: Oligochaeta). *Zootaxa*, (1281), 21-39. Retrieved from: www.scopus.com
- Timm, T. 1999. Distribution of freshwater oligochaetes in the west and east coastal regions of the north pacific ocean. *Hydrobiologia*, 406, 67-81. doi: [10.1023/A:1003749519662](https://doi.org/10.1023/A:1003749519662)
- Timm, T. 1997. Freshwater Oligochaeta of some urban watercourses in the russian far east. *Internationale Revue Der Gesamten Hydrobiologie Und Hydrographie*, 82(4), 437-467. doi: [10.1002/iroh.19970820402](https://doi.org/10.1002/iroh.19970820402)
- Timm, T. 1998. Lumbriculidae (Oligochaeta) of Lake Taimyr. *Journal of Natural History*, 32(9), 1291-1301. doi: [10.1080/00222939800770641](https://doi.org/10.1080/00222939800770641)
- Timm, T. 2012. Profundal oligochaete assemblages in palaeartic lakes. *Turkish Journal of Zoology*, 36(1), 121-131. doi: [10.3906/zoo-1002-51](https://doi.org/10.3906/zoo-1002-51)
- Timm, T. 1994. Propappidae and aquatic Enchytraeidae (Oligochaeta) from the farthest southeast of Russia. *Hydrobiologia*, 278(1-3), 67-78. doi: [10.1007/BF00142312](https://doi.org/10.1007/BF00142312)
- Timm, T. 2013. The genus *Potamothisrix* (Annelida, Oligochaeta, Tubificidae): A literature review. *Estonian Journal of Ecology*, 62(2), 121-136. doi: [10.3176/eco.2013.2.04](https://doi.org/10.3176/eco.2013.2.04)
- Timm, T. 1996. *Tubifex tubifex* (MÜLLER, 1774) (Oligochaeta, Tubificidae) in the profundal of estonian lakes. *Internationale Revue Der Gesamten Hydrobiologie Und Hydrographie*, 81(4), 589-596. doi: [10.1002/iroh.19960810412](https://doi.org/10.1002/iroh.19960810412)
- Timm, T., Arslan, N., Rüzgar, M., Martinsson, S. & Erséus, C. 2013. Oligochaeta (Annelida) of the profundal of Lake Hazar (Turkey), with description of *Potamothisrix alatus* Hazaricus ssp. nov. *Zootaxa*, 3716(2), 144-156. doi: [10.11646/zootaxa.3716.2.2](https://doi.org/10.11646/zootaxa.3716.2.2)
- Timm, T. & Grimm, R. 2005. What is *Homochaeta naidina* Bretscher, 1896 (Annelida, Oligochaeta, Naididae)? *Zoosystema*, 27(3), 469-482. Retrieved from: www.scopus.com
- Timm, T. & Martin, P.J. 2014. Clitellata: Oligochaeta. *Thorp and covich's freshwater invertebrates: Ecology and general biology: Fourth edition.* (pp. 529-549) doi: [10.1016/B978-0-12-385026-3.00021-8](https://doi.org/10.1016/B978-0-12-385026-3.00021-8). Retrieved from: www.scopus.com
- Timm, T. & Rodriguez, P. 1994. Description of a new Lumbriculus species (Oligochaeta, Lumbriculidae) from the Russian Far-East. [Description d'une nouvelle espèce de *Lumbriculus* (Oligochaeta, Lumbriculidae) de l'Extrême-Orient russe]. *Annales de Limnologie*, 30(2), 95-100. doi: [10.1051/limn/1994012](https://doi.org/10.1051/limn/1994012)
- Trodd, W.R., Kelly-Quinn, M., Sweeney, P. & Quirke, B. 2005. A review of the status and distribution of the free-living freshwater Oligochaeta of Ireland. *Biology and Environment*, 105(1), 59-64. doi: [10.3318/BIOE.2005.105.1.59](https://doi.org/10.3318/BIOE.2005.105.1.59)
- Tsytsugina, V.G., Gudkov, D.I., Nazarov, A.B. & Gorbenko, V.P. 2005. Damage to chromosomes in cells of oligochaetes *Stylaria lacustris* in water bodies of the Chernobyl NPP exclusion zone. *Hydrobiological Journal*, 41(5), 97-101. doi: [10.1615/HydrobJ.v41.i5.100](https://doi.org/10.1615/HydrobJ.v41.i5.100)
- Tsytsugina, V.G. & Polikarpov, G.G. 2000. Cytogenetic and population effects in Oligochaeta from the Chernobyl zone. *Radiatsionnaya Biologiya. Radioekologiya*, 40(2), 226-230. Retrieved from: www.scopus.com
- Tsytsugina, V.G. & Polikarpov, G.G. 2003. Radiological effects on populations of Oligochaeta in the Chernobyl contaminated zone. *Journal of Environmental Radioactivity*, 66(1-2), 141-154. doi: [10.1016/S0265-931X\(02\)00120-0](https://doi.org/10.1016/S0265-931X(02)00120-0)
- Uzunov, J., Košel, V. & Sládeček, V. 1988. Indicator value of freshwater Oligochaeta. *Acta Hydrochimica et Hydrobiologica*, 16(2), 173-186. doi: [10.1002/aheh.19880160207](https://doi.org/10.1002/aheh.19880160207)
- Van Duinen, G.A., Timm, T., Smolders, A.J.P., Brock, A.M.T., Verberk, W.C.E.P. & Esselink, H. 2006. Differential response of aquatic oligochaete species to

- increased nutrient availability – A comparative study between estonian and dutch raised bogs. *Hydrobiologia*, 564 (1 spec. iss.), 143-145. doi: [10.1007/s10750-005-1715-y](https://doi.org/10.1007/s10750-005-1715-y)
- Vandegehuchte, M.B., Roman, Y.E., Nguyen, L.T.H., Janssen, C.R. & De Schampheleire, K.A.C. 2007. Toxicological availability of nickel to the benthic oligochaete *Lumbriculus variegatus*. *Environment International*, 33(6), 736-742. doi: [10.1016/j.envint.2007.02.006](https://doi.org/10.1016/j.envint.2007.02.006)
- Vecchi, M., Reynoldson, T.B., Pasteris, A. & Bonomi, G. 1999. Toxicity of copper-spiked sediments to *Tubifex tubifex* (Oligochaeta, Tubificidae): Comparison of the 28-day reproductive bioassay with an early-life-stage bioassay. *Environmental Toxicology and Chemistry*, 18(6), 1173-1179. doi: [10.1897/1551-5028\(1999\)018<1173:TOCSST>2.3.CO;2](https://doi.org/10.1897/1551-5028(1999)018<1173:TOCSST>2.3.CO;2)
- Vedovini, A., Légier, P. & Vespini, F. 1985. Oligochaetes of a river in the South Alps, the Buech. [Les Oligochètes d'une rivière des Alpes du Sud, le Buech]. *Annales de Limnologie*, 21(2), 117-125. doi: [10.1051/limn/1985011](https://doi.org/10.1051/limn/1985011)
- Veltz, I., Arsac, F., Biagiotti-Risbourg, S., Habets, F., Lechenault, H. & Vernet, G. 1996. Effects of platinum (Pt4+) on *Lumbriculus variegatus* Müller (Annelida, Oligochaetae): Acute toxicity and bioaccumulation. *Archives of Environmental Contamination and Toxicology*, 31(1), 63-67. Retrieved from: www.scopus.com
- Veltz-Balatre, I., Biagiotti-Risbourg, S. & Vernet, G. 2000. De la régénération céphalique de l'oligochaète *Lumbriculus variegatus* Müller à l'étude de la toxicité des métaux lourds cephalic regeneration of freshwater Oligochaeta *Lumbriculus variegatus* Müller used for studying heavy metal toxicity. *Annee Biologique*, 39(1), 39-52. Retrieved from: www.scopus.com
- Verdonschot, P.F.M. 1987. Aquatic oligochaetes in ditches. *Hydrobiologia*, 155(1), 283-292. doi: [10.1007/BF00025661](https://doi.org/10.1007/BF00025661)
- Verdonschot, P.F.M. 2006. Beyond masses and blooms: The indicative value of oligochaetes. *Hydrobiologia*, 564 (1 spec. iss.), 127-142. doi: [10.1007/s10750-005-1714-z](https://doi.org/10.1007/s10750-005-1714-z)
- Verdonschot, P.F.M. 1999. Micro-distribution of oligochaetes in a soft-bottomed lowland stream (Elsbeek; the Netherlands). *Hydrobiologia*, 406, 149-163. doi: [10.1023/A:1003796403364](https://doi.org/10.1023/A:1003796403364)
- Verdonschot, P.F.M. 1989. The role of oligochaetes in the management of waters. *Hydrobiologia*, 180(1), 213-227. doi: [10.1007/BF00027554](https://doi.org/10.1007/BF00027554)
- Verdonschot, P.F.M. & Ter Braak, C.J.F. 1994. An experimental manipulation of oligochaete communities in mesocosms treated with chlorpyrifos or nutrient additions: Multivariate analyses with Monte Carlo permutation tests. *Hydrobiologia*, 278(1-3), 251-266. doi: [10.1007/BF00142333](https://doi.org/10.1007/BF00142333)
- Vidal, D.E. & Horne, A.J. 2003. Inheritance of mercury tolerance in the aquatic oligochaete *Tubifex tubifex*. *Environmental Toxicology and Chemistry*, 22(9), 2130-2135. doi: [10.1897/02-407](https://doi.org/10.1897/02-407)
- Vidal, D.E. & Horne, A.J. 2003. Mercury toxicity in the aquatic oligochaete *sparganophilus pearsei*: I. Variation in resistance among populations. *Archives of Environmental Contamination and Toxicology*, 45(2), 184-189. doi: [10.1007/s00244-003-0119-0](https://doi.org/10.1007/s00244-003-0119-0)
- Vivien, R. 2013. Note on the diversity of aquatic oligochaetes in the Geneva area (Switzerland). [Note sur la diversité des oligochètes aquatiques dans la région genevoise (Suisse)]. *Revue Suisse de Zoologie*, 120(3), 415-420. Retrieved from: www.scopus.com
- Vivien, R., La Font, M. & Perfetta, J. 2011. Proposal of heavy metal toxicity thresholds of sediments indicated by oligochaete worms in some watercourses. [Proposition d'un seuil de toxicité des métaux lourds des sédiments mis en évidence par les vers oligochètes dans quelques cours d'eau]. *Bulletin de la Société Vaudoise des Sciences Naturelles*, 92(3-4), 153-164. Retrieved from: www.scopus.com
- Vivien, R. & Lafont, M. 2013. Diversity of aquatic oligochaetes in the Geneva area (Switzerland). [Diversité des oligochètes aquatiques dans la région Genevoise (Suisse)]. *Revue Suisse de Zoologie*, 120(1), 161-173. Retrieved from: www.scopus.com
- Vivien, R. & Lafont, M. 2015. Faunistic note on the aquatic oligochaetes of the Geneva area and of Switzerland. [Note faunistique sur les oligochètes aquatiques de la région Genevoise et de Suisse]. *Revue Suisse de Zoologie*, 122(2), 207-212. doi: [10.5281/zenodo.29997](https://doi.org/10.5281/zenodo.29997)
- Vivien, R., Tixier, G. & Lafont, M. 2014. Use of oligochaete communities for assessing the quality of sediments in watercourses of the Geneva area (Switzerland) and artois-picardie basin (France): Proposition of heavy metal toxicity thresholds. *Ecohydrology and Hydrobiology*, 14(2), 142-151. doi: [10.1016/j.ecohyd.2014.03.003](https://doi.org/10.1016/j.ecohyd.2014.03.003)
- Vorob'ev, D.S., Frank, Y.A., Lushnikov, S.V., Zaloznyi, N.A. & Noskov, Y.A. 2010. Oil decontamination of bottom sediments using *Limnodrilus hoffmeisteri* (Oligochaeta: Tubificidae). *Contemporary Problems of Ecology*, 3(1), 15-18. doi: [10.1134/S1995425510010042](https://doi.org/10.1134/S1995425510010042)
- Wang, H. & Liang, Y. 2001. A preliminary study of oligochaetes in Poyang Lake, the largest freshwater lake of China, and its vicinity, with description of a new species of *Limnodrilus*. *Hydrobiologia*, 463, 29-38. doi: [10.1023/A:1013126918728](https://doi.org/10.1023/A:1013126918728)
- Wang, H., Xie, Z. & Liang, Y. 1999. Records of Enchytraeidae (Clitellata) from the people's Republic of China. *Hydrobiologia*, 406, 57-66. doi: [10.1023/A:1003732116567](https://doi.org/10.1023/A:1003732116567)
- Wang, X. & Matisoff, G. 1997. Solute transport in sediments by a large freshwater oligochaete, *Branchiura sowerbyi*. *Environmental Science & Technology*, 31(7), 1926-1933. doi: [10.1021/es960716n](https://doi.org/10.1021/es960716n)
- Weerasundara, A., Pathiratne, A. & Costa, H. 2000. Species composition and abundance of littoral oligochaete fauna in Lunuwila Reservoir, Sri Lanka. *International Review of Hydrobiology*, 85(2-3), 223-230. doi: [10.1002/\(SICI\)1522-2632\(200004\)85:2/3<223::AID-IROH223>3.0.CO;2-1](https://doi.org/10.1002/(SICI)1522-2632(200004)85:2/3<223::AID-IROH223>3.0.CO;2-1)
- Wegener, J.-M., van den Berg, G.A., Stroomberg, G.J. & van Velzen, M.J.M. 2002. The role of sediment-feeding oligochaete *Tubifex* on the availability of trace metals in sediment pore waters as determined by diffusive gradients in thin films (DGT). *Journal of Soils and Sediments*, 2(2), 71-76. doi: [10.1007/BF02987874](https://doi.org/10.1007/BF02987874)
- Wei, Y. & Liu, J. 2005. The discharged excess sludge treated by oligochaete. Retrieved from: www.scopus.com
- Wei, Y., Van Houten, R.T., Borger, A.R., Eikelboom, D.H. & Fan, Y. 2003. Comparison performances of membrane bioreactor and conventional activated sludge processes on sludge reduction induced by oligochaete. *Environmental Science & Technology*, 37(14), 3171-3180. doi: [10.1021/es026176d](https://doi.org/10.1021/es026176d)
- Wei, Y., Wang, Y., Guo, X. & Liu, J. 2009. Sludge reduction potential of the activated sludge process by integrating an oligochaete reactor. *Journal of Hazardous Materials*, 163(1), 87-91. doi: [10.1016/j.jhazmat.2008.06.065](https://doi.org/10.1016/j.jhazmat.2008.06.065)
- Wei, Y., Zhu, H., Wang, Y., Li, J., Zhang, P., Hu, J. & Liu, J. 2009. Nutrients release and phosphorus distribution during oligochaetes predation on activated sludge. *Biochemical Engineering Journal*, 43(3), 239-245. doi: [10.1016/j.bej.2008.10.004](https://doi.org/10.1016/j.bej.2008.10.004)
- Weider, L.J. 1992. Allozymic variation in tubificid oligochaetes from the Laurentian great lakes. *Hydrobiologia*, 234(2), 79-85. doi: [10.1007/BF00010863](https://doi.org/10.1007/BF00010863)
- West, C.W. & Ankley, G.T. 1998. A laboratory assay to assess avoidance of contaminated sediments by the freshwater oligochaete *Lumbriculus variegatus*. *Archives of Environmental Contamination and Toxicology*, 35(1), 20-24. doi: [10.1007/s002449900343](https://doi.org/10.1007/s002449900343)
- Westheide, W., Purschke, G. & Middendorf, K. 1991. Spermatozoal ultrastructure of the taxon *Enchytraeus* (Annelida, Oligochaeta) and its significance for species discrimination and identification. *Journal of Zoological Systematics and Evolutionary Research*, 29(5-6), 323-342. doi: [10.1111/j.1439-0469.1991.tb00456.x](https://doi.org/10.1111/j.1439-0469.1991.tb00456.x)

- Wetzel, M.J., Oberlin, G.E. & Blinn, D.W. 1999. The aquatic Oligochaeta (Annelida: Clitellata) of Montezuma Well, Arizona: A near thermally constant limnocrene. *Southwestern Naturalist*, 44(4), 514-518. doi: [10.2307/3672352](https://doi.org/10.2307/3672352)
- Wetzel, M.J. & Taylor, S.J. 2001. First records of freshwater oligochaetes (Annelida, Clitellata) from caves in Illinois and Missouri, USA. *Journal of Cave and Karst Studies*, 63(3), 99-104. Retrieved from: www.scopus.com
- Whittaker, J.W., Pote, L.M.W. & Hanson, L.A. 2005. Assay to detect the actinospore and myxospore stages of proliferative gill disease in oligochaetes and pond water. *North American Journal of Aquaculture*, 67(2), 133-137. doi: [10.1577/A03-059.1](https://doi.org/10.1577/A03-059.1)
- White, D.S. & Keily, T.J. 1988. Burrowing avoidance assays of contaminated Detroit River sediments, using the freshwater oligochaete *Stylodrilus heringianus* (Lumbriculidae). *Archives of Environmental Contamination and Toxicology*, 17(5), 673-681. doi: [10.1007/BF01055837](https://doi.org/10.1007/BF01055837)
- White, D.S., Klahr, P.C. & Robbins, J.A. 1987. Effects of temperature and density on sediment reworking by *Stylodrilus heringianus* (Oligochaeta: Lumbriculidae). *Journal of Great Lakes Research*, 13(2), 147-156. doi: [10.1016/S0380-1330\(87\)71638-4](https://doi.org/10.1016/S0380-1330(87)71638-4)
- Wieltschnig, C., Kirschner, A.K.T., Fischer, U.R. & Velimirov, B. 2003. Top-down control of benthic heterotrophic nanoflagellates by oligochaetes and microcrustaceans in a littoral freshwater habitat. *Freshwater Biology*, 48(10), 1840-1849. doi: [10.1046/j.1365-2427.2003.01133.x](https://doi.org/10.1046/j.1365-2427.2003.01133.x)
- Wolf, K., Markiw, M.E. & Hiltunen, J.K. 1986. Salmonid whirling disease: *Tubifex tubifex* (Müller) identified as the essential oligochaete in the protozoan life cycle. *Journal of Fish Diseases*, 9(1), 83-85. doi: [10.1111/j.1365-2761.1986.tb00984.x](https://doi.org/10.1111/j.1365-2761.1986.tb00984.x)
- Xiao, C. & Desser, S.S. 1998. Actinosporean stages of myxozoan parasites of oligochaetes from Lake Sasajewun, Algonquin Park, Ontario: New forms of *echinactinomyxon*, *neoactinomyxum*, *aurantiactinomyxon*, *guyenotia*, *synactinomyxon* and *antonactinomyxon*. *Journal of Parasitology*, 84(5), 1010-1019. doi: [10.2307/3284635](https://doi.org/10.2307/3284635)
- Xiao, C. & Desser, S.S. 1998. Actinosporean stages of myxozoan parasites of oligochaetes from Lake Sasajewun, Algonquin Park, Ontario: New forms of *triactinomyxon* and *rabeiae*. *Journal of Parasitology*, 84(5), 998-1009. doi: [10.2307/3284634](https://doi.org/10.2307/3284634)
- Xiao, C. & Desser, S.S. 1998. The oligochaetes and their actinosporean parasites in Lake Sasajewun, Algonquin Park, Ontario. *Journal of Parasitology*, 84(5), 1020-1026. doi: [10.2307/3284636](https://doi.org/10.2307/3284636)
- Xie, L., Lambert, D., Martin, C., Cain, D.J., Luoma, S.N. & Buchwalter, D. 2008. Cadmium biodynamics in the oligochaete *Lumbriculus variegatus* and its implications for trophic transfer. *Aquatic Toxicology*, 86(2), 265-271. doi: [10.1016/j.aquatox.2007.11.008](https://doi.org/10.1016/j.aquatox.2007.11.008)
- Xie, Z. & Chen, J. 2008. A redescription of *Piguetiella denticulata liang and xie* (Oligochaeta, Tubificidae, Naidinae) from China, with a key to the genus *Piguetiella*. *Proceedings of the Biological Society of Washington*, 121(3), 321-325. doi: [10.2988/08-08.1](https://doi.org/10.2988/08-08.1)
- Xie, Z., Shu, S., Zhang, J., Chen, J. & Cai, Q. 2008. Oligochaete assemblages associated with macrophytes in the Liangzi Lake District, China. *Journal of Freshwater Ecology*, 23(2), 237-244. doi: [10.1080/02705060.2008.9664196](https://doi.org/10.1080/02705060.2008.9664196)
- Yanling, L. 1987. Preliminary study of the aquatic Oligochaeta of the Changjiang (Yangtze) River. *Hydrobiologia*, 155(1), 195-198. doi: [10.1007/BF00025651](https://doi.org/10.1007/BF00025651)
- Yasuda, K. & Okino, T. 1987. Distribution and seasonal changes of aquatic Oligochaeta in Lake Suwa. *Japanese Journal of Limnology (Rikusuigaku Zasshi)*, 48(1), 1-8. doi: [10.3739/rikusu.48.1](https://doi.org/10.3739/rikusu.48.1)
- Yasuda, K. & Okino, T. 1988. Method for distinguishing *Limnodrilus hoffmeisteri* and *Limnodrilus claparedaeianus* (Oligochaeta, Tubificidae) and its applicability in Lake Suwa. *Hydrobiologia*, 169(3), 307-311. doi: [10.1007/BF00007553](https://doi.org/10.1007/BF00007553)
- Yıldız, S. & Ahiska, S. 2010. *Nais stolci* hrabě, 1981: A new oligochaete (Annelida: Naididae) species for Turkey. [Nais stolci Hrabě, 1981: Türkiye için yeni bir Oligoket (Annelida, Oligochaeta) Türü]. *Turkish Journal of Zoology*, 34(4), 547-549. doi: [10.3906/zoo-0906-3](https://doi.org/10.3906/zoo-0906-3)
- Yıldız, S. & Balık, S. 2006. The Oligochaeta (Annelida) fauna of topçam Dam Lake (Aydın, Turkey). *Turkish Journal of Zoology*, 30(1), 83-89. Retrieved from: www.scopus.com
- Yıldız, S. & Balık, S. 2010. *Nais christinae* Kasprzak, 1973: An Oligochaeta species new for Turkey (Annelida: Naididae). *Zoology in the Middle East*, 50(1), 151-152. doi: [10.1080/09397140.2010.10638432](https://doi.org/10.1080/09397140.2010.10638432)
- Yıldız, S., Ustaoglu, M.R. & Balık, S. 2007. Contributions to the knowledge of the Oligochaeta (Annelida) fauna of some lakes in the taurus mountain range (Turkey). *Turkish Journal of Zoology*, 31(3), 249-254. Retrieved from: www.scopus.com
- Yokoyama, H., Ogawa, K. & Wakabayashi, H. 1993. Some biological characteristics of actinosporeans from the oligochaete *Branchiura sowerbyi*. *Diseases of Aquatic Organisms*, 17(3), 223-228. doi: [10.3354/dao017223](https://doi.org/10.3354/dao017223)
- Zendt, J.S. & Bergersen, E.P. 2000. Distribution and abundance of the aquatic oligochaete host *Tubifex tubifex* for the salmonid whirling disease parasite *Myxobolus cerebralis* in the upper Colorado River basin. *North American Journal of Fisheries Management*, 20(2), 502-512. doi: [10.1577/1548-8675\(2000\)020<0502:DAAOTA>2.3.CO;2](https://doi.org/10.1577/1548-8675(2000)020<0502:DAAOTA>2.3.CO;2)
- Zeybek, M., Kalyoncu, H. & Ertan, O.O. 2013. The effects of environmental variables on distribution of aquatic Oligochaeta and Chironomidae at kovada channel and the Linked Lakes (Isparta, Turkey). *Fresenius Environmental Bulletin*, 22(11), 3160-3169. Retrieved from: www.scopus.com
- Zhang, B., Li, H., Wei, Y. & You, J. 2013. Bioaccumulation kinetics of polybrominated diphenyl ethers and decabromodiphenyl ethane from field-collected sediment in the oligochaete, *Lumbriculus variegatus*. *Environmental Toxicology and Chemistry*, 32(12), 2711-2718. doi: [10.1002/etc.2384](https://doi.org/10.1002/etc.2384)
- Zhang, J., Chen, J. & Xie, Z. 2007. Redescription of *Nais inflata liang* (Annelida: Oligochaeta) from China. *Proceedings of the Biological Society of Washington*, 120(3), 264-270. doi: [10.2988/0006-324X\(2007\)120\[264:RONILA\]2.0.CO;2](https://doi.org/10.2988/0006-324X(2007)120[264:RONILA]2.0.CO;2)
- Zhu, H., Wei, Y.-., Wang, Y.-. & Liu, J.-. 2008. Advances in sludge reduction induced by oligochaetes and theirs growth kinetics. *Guocheng Gongcheng Xuebao – The Chinese Journal of Process Engineering*, 8(5), 1030-1040. Retrieved from: www.scopus.com
- Zvereva, Y.M., Mizandrontsev, I.B., Zaytseva, Y.P. & Timoshkin, O.A. 2015. Respiration of aquatic oligochaetes (Annelida, Oligochaeta) and peculiarities of oxygen uptake by endemic Enchytraeidae of Lake Baikal. *Hydrobiological Journal*, 51(5), 14-24. doi: [10.1615/HydrobJ.v51.i5.20](https://doi.org/10.1615/HydrobJ.v51.i5.20)