

Two newly introduced Heteroptera (Insecta: Hemiptera) species in Colombia: *Brachyplatys subaenus* (Plataspidae) and *Thaumastocoris peregrinus* (Thaumastocoridae)

Dimitri Forero¹

¹ Pontificia Universidad Javeriana, Facultad de Ciencias, Departamento de Biología, Laboratorio de Entomología, Unidad de Ecología y Sistemática (UNESIS). Bogotá, D.C., Colombia.
ORCID: <http://orcid.org/0000-0002-6358-757X>. E-mail: forero-i@javeriana.edu.co

Abstract. The introduction of alien species can carry negative consequences to the areas in which they appear. Early detection of introduced species is key if management practices are to be effectively implemented. Here, based on specimens from collections and citizen science observations, we document the recent introduction of two alien Heteroptera (Insecta: Hemiptera) species in Colombia: *Brachyplatys subaenus* (Westwood, 1837) (Plataspidae) and *Thaumastocoris peregrinus* Carpintero & Dellapé, 2006 (Thaumastocoridae). *Brachyplatys subaenus* was found in two localities (Bolívar and Valle del Cauca), whereas *T. peregrinus* was found in a single locality (Bogotá) so far. Future research activities should establish the geographic range of these species, as well as document their host plant associations, in Colombia.

Keywords. Exotic species; Neotropical region; Black bean bug; Bronze bug; South America.

INTRODUCTION

Exotic invasive species are costly to society (Simberloff *et al.*, 2013), and among these, insect species are considered the most expensive (Diagne *et al.*, 2021). Among all living organisms insects constitute about 40% of all known species (Adler & Foottit, 2017; Stork, 2018), thus it is not surprising that the most common group of introduced animals found worldwide are insects (Liebhold *et al.*, 2018). Because of the ample negative societal impacts of alien species, ranging from ecosystem function to human well-being (Pyšek *et al.*, 2020), early detection is key to implement appropriate management practices (Pyšek & Richardson, 2010; Simberloff *et al.*, 2013).

Hemiptera is the fifth most diverse order of insects, with Heteroptera containing the largest proportion of its diversity (Henry, 2017). Several species of Heteroptera are considered of economic importance, mostly in agricultural settings (Schaefer & Panizzi, 2000), and many are now exotic elements of the fauna of several regions of the world (*e.g.*, Scudder & Foottit, 2006; Rabitsch, 2008, 2010; Henry *et al.*, 2013), in many cases causing economic losses (Lee, 2015; McPherson, 2018). Economically important alien invasive Heteroptera include phytophagous species such

as *Bagrada hilaris* (Burmeister, 1835) (Palumbo *et al.*, 2016; Carvajal *et al.*, 2019), *Halyomorpha halys* (Stål, 1855) (Nielsen & Hamilton, 2009; Lee *et al.*, 2013), *Leptoglossus occidentalis* Heidemann, 1910 (Lesieur *et al.*, 2019), and *Nezara viridula* (Linnaeus, 1758) (McPherson & McPherson, 2000; Esquivel *et al.*, 2018). Other introduced Heteroptera may include predatory species, such as *Zelus renardii* Kolenati, 1857, which is now widely distributed in the world (Weirauch *et al.*, 2012; Zhang *et al.*, 2016).

Several Neotropical Heteroptera species, specially within Pentatomoidea, have been identified as potential invasives for other regions (Panizzi, 2015), but little research has been carried out to identify introduced Heteroptera within the Neotropics. In South America, just a few species have been identified as alien, for instance, *B. hilaris* and *H. halys* have been recorded from Argentina and Chile (Faúndez *et al.*, 2016; Faúndez & Rider, 2017; Carpintero *et al.*, 2021b); *L. occidentalis* from Chile, Argentina, and Brazil (Kun & Masciocchi, 2019; van der Heyden & Faúndez, 2020); *Boisea trivittata* (Say, 1825) from Chile (Faúndez *et al.*, 2020); *Holcocranum saturejae* (Kolenati, 1845) (Carpintero *et al.*, 2021a) from Argentina; *Z. renardii* from Chile and Argentina (Curkovic *et al.*, 2004; D'Hervé *et al.*, 2018); and *Amphibolus venator* (Klug, 1830) from

Pap. Avulsos Zool., 2021; v.61: e20216196

<http://doi.org/10.11606/1807-0205/2021.61.96>

<http://www.revistas.usp.br/paz>

<http://www.scielo.br/paz>

Edited by: Cristiano Feldens Schwertner

Received: 06/07/2021

Accepted: 27/09/2021

Published: 27/10/2021

ISSN On-Line: 1807-0205

ISSN Printed: 0031-1049

ISNI: 0000-0004-0384-1825



Venezuela (Forero *et al.*, 2004). In Colombia, the checklist of introduced and invasive species (Baptiste *et al.*, 2020), only list seven Hemiptera species, of which just one is a Heteroptera (Miridae, *Tropidosteptes chapingoensis* Carvalho & Rosas, 1965). Nonetheless, there are other recently introduced species, such as *Spilostethus pandurus* (Scopoli, 1763) (Rengifo-Correa & González, 2011) now common in Colombia (see iNaturalist records: <https://bit.ly/3qagAbe>), which are not listed yet, therefore, it is likely that additional species might be discovered with the study of biological collections, more fieldwork, and the involvement of research taxonomists.

Here, I document two introduced species in Colombia that possess the potential to become invasive in agricultural settings, highlighting the importance of an early detection to better implement management strategies.

MATERIAL AND METHODS

Examined specimens are deposited at the Colección Entomológica, Museo Javeriano de Historia Natural, Pontificia Universidad Javeriana, Bogotá, Colombia (MPUJ_ENT). In addition, some observations uploaded to the citizen science initiative iNaturalist (<https://www.inaturalist.org>), were used to complement the distribution documented from specimens. In this case, the observations were all identified or corroborated by me. Images of specimens were carried out using a dissecting scope Nikon SMZ1270 equipped with a Nikon D5300 digital camera.

RESULTS

Brachyplatys subaenus (Westwood, 1837) (Fig. 1)

Remarks: The highest diversity of Plataspidae is found in tropical and subtropical regions of the Eastern Hemisphere, with a few species widely distributed in the temperate Palearctic (Schuh & Weirauch, 2020). The first species of Plataspidae recorded from the Western Hemisphere was *Megacocta cribraria* (Fabricius, 1798), discovered in the USA in 2009 (Eger *et al.*, 2010). This species is found from China (temperate and tropical) to tropical Asia and Australia, and it is associated mainly with kudzu *Pueraria montana* var. *lobata* (Willd.) Maesen & Almeida, and which has been considered recently a pest of soybeans *Glycine max* (L.) (Ruberson *et al.*, 2013). A second plataspid found in the Western Hemisphere is *B. subaenus*, which was first discovered in Panama in 2012, being also the first record from the Neotropical region (Aiello *et al.*, 2016). This species was initially misidentified in Panama as *B. vahlii* (Fabricius, 1787) but later correctly identified as *B. subaenus* by Rédei (2016). Rédei (2016) provided diagnostic characters that help identify this species and differentiate it from *B. vahlii*.

Plataspidae species feed primarily on legumes (Schaefer, 1988), being *B. subaenus* no exception, which

is known to feed on a variety of legumes species in its native range in Indomalaya, although other plant species in various families have also been recorded as host plants (Rédei, 2016). In the Neotropical countries in which *B. subaenus* has been found, it has been recorded on the following Fabaceae host plants: the cultivated *Cajanus cajan* (L.) Mill sp., *Glycine max*, and *Mucuna pruriens* (L.) DC (Aiello *et al.*, 2016; Añino *et al.*, 2018, 2020; Carmona-Ríos, 2019; Perez-Gelabert *et al.*, 2019); the endemic and cultivated *Phaseolus vulgaris* L. (Perez-Gelabert *et al.*, 2019), *Gliricidia sepium* (Jacq.) Kunth ex Walp. (Añino *et al.*, 2018; Carmona-Ríos, 2019); and the endemic *Leptolobium panamense* (Benth.) Sch.Rodr. & A.M.G.Azevedo (Aiello *et al.*, 2016). It has also been found on the following non-Fabaceae host species: *Bactris gasipaes* Kunth, *Schefflera actinophylla* (Endl.) Harms (Aiello *et al.*, 2016), and *Zea mays* L. (Añino *et al.*, 2020). In Colombia, *B. subaenus* has no recorded host plant species.

Distribution: Known in the Neotropical region from Costa Rica (Carmona-Ríos, 2019), Panama (Añino *et al.*, 2018), Dominican Republic (Perez-Gelabert *et al.*, 2019), Guadeloupe (Anonymous, 2020), and Ecuador (Añino *et al.*, 2020). Additional localities can be found on iNaturalist (<http://www.inaturalist.org>) for some of these countries. Newly recorded from Colombia from two localities, in Cali (Valle del Cauca) and Isla Grande (Bolívar).

Examined material: COLOMBIA. Valle del Cauca, 1 female, Cali, calle 5 con carrera 39, 03.4265°N, 76.5454°W, 974 m, 11 Abr 2019, D. Forero, dentro de edificio, en vuelo [within building, flying], MPUJ_ENT0063186 (MPUJ_ENT).

iNaturalist observations: COLOMBIA. Bolívar, 1 adult, sex not determined, Isla Grande, 10.1807°N, 75.7249°W, 16 Mar 2019, C.M. Rangel, <https://www.inaturalist.org/observations/21342531>. **Valle del Cauca,** 1 adult, sex not determined, Cali, Villa Fatima, 03.3519°N, 76.5097°W, 02 Jan 2021, L. Rubio, <https://www.inaturalist.org/observations/68191884>; nymphs, Cali, Mariposario Andoke, km 6, Parcelación La Reforma, vía a Cristo Rey, 03.4261°N, 76.5719°W, 01 May 2021, <https://www.inaturalist.org/observations/76232278>.

Thaumastocoris peregrinus Carpintero & Dellapé, 2006 (Fig. 2)

Remarks: *Thaumastocoris peregrinus* was described based on specimens collected in Argentina but acknowledging its Australian origin (Carpintero & Dellapé, 2006). Noack & Coviella (2006) simultaneously reported a *Thaumastocoris* species from Buenos Aires (Argentina) but misidentified it as *T. australicus* Kirkaldy, which was later identified as *T. peregrinus* (Noack *et al.*, 2011). Carpintero & Dellapé (2006) also assigned to *T. peregrinus* specimens found in South Africa but also misidentified as *T. australicus*. The taxonomy of this Australian endemic genus *Thaumastocoris* was clarified by Noack *et al.* (2011),

in which they provide a revised diagnosis and comparative notes for *T. peregrinus*.

Most of the host plant associations of *Thaumastocoris* are with species of Myrtaceae, in which nearly all species are associated with species of *Eucalyptus* L'Hér. (Noack *et al.*, 2011). *Thaumastocoris peregrinus* has been documented to feed on one species of *Corymbia* K.D. Hill & L.A.S. Johnson and at least 13 species of *Eucalyptus* (Jacobs & Nesar, 2005; Noack & Coviella, 2006; Noack *et al.*, 2011; Nascimento-Machado *et al.*, 2019). The Colombian examined specimens were collected in an unidentified species of *Eucalyptus*. In all places in which *T. peregrinus* has been accidentally introduced, it has caused damages to ornamental or cultivated *Eucalyptus* species (Machado

et al., 2020), and it is likely that biological control might be the best management option (Nadel & Noack, 2012; Souza *et al.*, 2012). Pinzón-Florián (2020) reviewed the pest species of *Eucalyptus* in Colombia, for which the only Heteroptera listed as pest was the mirid *Monalonion velezangeli* Carvalho & Costa. *Thaumastocoris peregrinus* is the only other Heteroptera pest species associated with *Eucalyptus* in Colombia.

Distribution: Originally from Australia, but accidentally introduced in South America (Brazil, Argentina, Uruguay, Paraguay, Chile), North America (Mexico, USA), Africa (South Africa, Kenya, Zimbabwe), Reunion Islands, Israel, New Zealand, and more recently in Europe (Italy,




Figure 1. *Brachyplatys subaenus*, female specimen from Cali (Valle del Cauca, Colombia) [MPUJ_ENT0063186]. (A) Dorsal view; (B) ventral view; (C) lateral view; (D) detail of scent gland apparatus in ventral view.




Figure 2. *Thaumastocoris peregrinus*, female specimen from Bogotá (Bogotá, D.C., Colombia). (A) Dorsal view; (B) ventral view [MPUJ_ENT0065778]; (C) detail of the male pygophore in ventral view [MPUJ_ENT0065773].

Portugal, Spain, Albania, Greece) (Carpintero & Dellapé, 2006; Noack *et al.*, 2011; Nascimento-Machado *et al.*, 2019). Newly recorded from Colombia from a single locality in the high Andes (Bogotá).

Examined material: 5 males, 3 females, 1 nymph, 1 adult of undetermined sex (damaged abdomen), **COLOMBIA, Bogotá D.C.**, parque Santa Helena de Baviera, 04.7313°N, 74.0569°W, 2,559 m, 26 Mar 2019, D. Forero; ex. *Eucalyptus* sp., MPUJ_ENT0065771 – MPUJ_ENT0065780 (MPUJ_ENT).

DISCUSSION

In Colombia, *B. subaenus* is only known from two localities, one in the Cauca valley (Cali) and the other in the Caribbean coast (Isla Grande). In all the other Neotropical countries from Central and South America where it has

been recorded, the localities are close to the seashore (Aiello *et al.*, 2016; Añino *et al.*, 2018, 2020; Carmona-Ríos, 2019). In Colombia, the Caribbean locality is very close to Cartagena, a coastal touristic city receiving people from all over the world, mostly from cruises. This locality is also very close (less than 100 km) to Barranquilla, a city with the main port and with the most important industrial facilities on the Colombian Caribbean coast. Therefore, it is likely that *B. subaenus* might have arrived from another place in the Caribbean via either commercial or touristic routes. This also means, that given the proximity of the documented locality to potential areas of easy dispersion by human means, it is likely that *B. subaenus* would be soon found in other localities in Colombia. On the other hand, although Cali is in an inter-Andean valley far away from the seashore, it is the main city connecting the Buenaventura port in the Pacific coast to the rest of the cities in Colombia. Therefore, it seems plausible that individuals of *B. subaenus* found in Cali, might have ar-

rived there from Buenaventura through sea commerce, and dispersed via terrestrial transportation to the interior of the country.

Thaumastocoris peregrinus, although widely distributed now in the Western Hemisphere, has not been recorded before from northern South America. It is intriguing that the locality in which it was found was in the middle of this large city in the high Andes. This could mean either that *T. peregrinus* arrived through commerce goods directly to Bogotá, or that it arrived at some lowland locality, and it later dispersed into the city. The first scenario would imply a contained distribution, whereas the second one would imply a much larger distribution already for *T. peregrinus*.

For both *B. subaenus* and *T. peregrinus* it is thus very important to document their present distribution and host plant associations in Colombia. An analysis of their geographic distribution will help to better understand the biology and development of each species regarding abiotic variables such as temperature or rainfall. Host plant associations will allow to, first, evaluate if these alien species are causing economic losses to cultivated crops, and second, to assess the extent of plant species serving as secondary hosts. This baseline information would be important to later implement adequate management practices for both species. Finding *B. subaenus* and *T. peregrinus* in Colombia for the first time highlights the importance of accurate taxonomic information derived from natural history collections and citizen science initiatives.

ACKNOWLEDGMENTS

I thank Wolfgang Rabitsch (University of Vienna) and two anonymous reviewers for their criticism that improved the manuscript. This paper is a contribution to the project “Actividades docentes y de investigación como apoyo al conocimiento de la biodiversidad colombiana” ID PPTA 00006416 to DF of the Pontificia Universidad Javeriana.

CONFLICT OF INTEREST

The author declares that he has no conflict of interest and confirms that there are no disputes over the ownership of the data presented, and all contributions have been attributed appropriately.

REFERENCES

- Adler, P.H. & Foottit, R.G. 2017. Introduction. In: Foottit, R.G. & Adler, P.H. (Eds.). *Insect biodiversity*. 2 ed. Hoboken, John Wiley. v. 1, p. 1-7.
- Aiello, A.; Saltonstall, K. & Young, V. 2016. *Brachyplatys vahlii* (Fabricius, 1787), an introduced bug from Asia: first report in the Western Hemisphere (Hemiptera: Plataspidae: Brachyplatidinae). *BiolInvasions Records*, 5: 7-12.
- Añino, Y.J.; Santos Murgas, A.; Henríquez Chirú, G.N.; Carranza, R. & Villareal, C. 2018. Distribución espacial del chinche invasor *Brachyplatys subaenus* (Westwood, 1837) (Hemiptera: Heteroptera: Plataspidae) en Panamá. *Insecta Mundi*, 630: 1-6.
- Añino, Y.J.; Sumba-Zhongor, M.B.; Naranjo-Morán, J.A.; Rodríguez, R.; Santos-Murgas, A. & Zachrisson, B. 2020. First report by *Brachyplatys subaenus* (Westwood) (Hemiptera: Plataspidae) for Ecuador and the synoptic list of its host plants. *IDESIA*, 38: 113-118.
- Anonymous. 2020. *Premier signalement d'une punaise invasive en Guadeloupe*. 3p.
- Baptiste, M.P.; García, L.M.; Acevedo-Charry, O.; Acosta, A.; Alarcón, J.; Arévalo, E.; Avella, G.C.; Blanco, A.; Botero, J.; Caicedo-Portilla, J.R.; Camelo Martínez, C.; Camelo-Calvo, M.P.; Certuche-Cubillos, K.; Chasqui, L.; Cifuentes, Y.; Contreras, P.J.; Córdoba, S.; Correa, J.; Díaz, M.F.; DoNascimento, C.; Duque, R.A.; Flechas, S.V.; Forero, D.; Gómez Hoyos, A.J.; González Durán, G.; Guayara, S.; Guetiva, J.C.; Jiménez, G.; Larrahondo, M.; Maldonado Ocampo, J.; Medina-Rangel, G.F.; Merino, M.C.; Mesa, L.M.; Millán, M.V.; Mojica, H.; Neita Moreno, J.C.; Parrado, M.P.; Pérez, S.C.; Ramírez, W.; Rojas, V.; Rojas, Z.; Urbina-Cardona, N.; Velásquez, P.L.; Wong, J.L. & Pagad, S. 2020. *Global Register of Introduced and Invasive Species – Colombia. Version 1.5*. Invasive Species Specialist Group ISSG. Available: <http://doi.org/10.15468/yznr8v>. Access: 06/07/2021.
- Carmona-Ríos, M.G. 2019. Primer registro de *Brachyplatys subaenus* (Westwood) (Hemiptera: Plataspidae) en Costa Rica. *Revista Chilena de Entomología*, 45: 583-586.
- Carpintero, D.L. & Dellapé, P.M. 2006. A new species of *Thaumastocoris* Kirkaldy from Argentina (Heteroptera: Thaumastocoridae: Thaumastocorinae). *Zootaxa*, 1228: 61-68. [DOI](#)
- Carpintero, D.L.; de Magistris, A.A. & Faúndez, E.I. 2021a. Primer registro para el Neotrópico de la familia Artheneidae Stål, 1872 (Heteroptera: Lygaeoidea), con la especie *Holcocranum saturejae* (Kolenati, 1845) introducida en Argentina. *Revista Chilena de Entomología*, 47(2): 311-318.
- Carpintero, D.L.; Quiroga, V.N.; Celentano, E. & Holgado, M.G. 2021b. Primer registro de *Bagrada hilaris* (Burmeister, 1835) (Hemiptera: Pentatomidae) para la república Argentina. *Historia Natural*, 11: 179-184.
- Carvajal, M.A.; Alaniz, A.J.; Núñez-Hidalgo, I. & González-Céspedes, C. 2019. Spatial global assessment of the pest *Bagrada hilaris* (Burmeister) (Heteroptera: Pentatomidae): current and future scenarios. *Pest Management Science*, 75(3): 809-820. [DOI](#)
- Curkovic, T.; Araya, J.; Baena, M. & Guerrero, M.A. 2004. Presencia de *Zelus renardii* Kolenati (Heteroptera: Reduviidae) en Chile. *Boletín de la Sociedad Entomológica Aragonesa*, 34: 163-165.
- D'Hervé, F.E.; Olave, A. & Dapoto, G.L. 2018. *Zelus renardii* (Hemiptera: Reduviidae: Harpactorinae: Harpactorini): primer registro en la Argentina. *Revista de la Sociedad Entomológica Argentina*, 77: 32-35. [DOI](#)
- Diagne, C.; Leroy, B.; Vaissière, A.-C.; Gozlan, R.E.; Roiz, D.; Jarić, I.; Salles, J.M.; Bradshaw, C.J.A. & Courchamp, F. 2021. High and rising economic costs of biological invasions worldwide. *Nature*, 592(7855): 1-6. [DOI](#)
- Eger, J.E.; Ames, L.M.; Suiter, D.R.; Jenkins, T.M.; Rider, D.A. & Halbert, S.E. 2010. Occurrence of the Old World bug *Megacopta cribraria* (Fabricius) (Heteroptera: Plataspidae) in Georgia: a serious home invader and potential legume pest. *Insecta Mundi*, 0121: 1-11.
- Esquivel, J.F.; Musolin, D.L.; Jones, W.A.; Rabitsch, W.; Greene, J.K.; Toews, M.D.; Schwertner, C.F.; Grazia, J.; McPherson, R.M. & McPherson, J.E. 2018. *Nezara viridula* (L.). In: *Invasive stink bugs and related species (Pentatomidae)*. Boca Raton, CRC Press.
- Faúndez, E.I. & Rider, D.A. 2017. The brown marmorated stink bug *Halyomorpha halys* (Stål, 1855) (Heteroptera: Pentatomidae) in Chile. *Archivos Entomológicos, Revista Galega de Entomoloxía*, 17: 305-307. https://www.aegaweb.com/archivos_entomoloxicos/vol_17_2017.htm.

- Faúndez, E.I.; Carvajal, M.A. & Sarmiento, C. 2020. Detection of the boxelder bug *Boisea trivittata* (Say, 1825) (Heteroptera: Rhopalidae) in Chile. *Heteroptera Poloniae – Acta Faunistica*, 14: 125-126. [DOI](#)
- Faúndez, E.I.; Lüer, A.; Cuevas, Á.G.; Rider, D.A. & Valdebenito, P. 2016. First record of the painted bug *Bagrada hilaris* (Burmeister, 1835) (Heteroptera: Pentatomidae) in South America. *Arquivos Entomológicos, Revista Galega de Entomología*, 16: 175-179. https://www.aegaweb.com/arquivos_entomologicos/vol_16_2016.htm.
- Forero, D.; Weirauch, C. & Baena, M. 2004. Synonymy of the reduviid (Hemiptera: Heteroptera) genus *Torrealbaia* (Triatominae) with *Amphibolus* (Harpactorinae), with notes on *Amphibolus venator* (Klug, 1830). *Zootaxa*, 670: 1-12. [DOI](#)
- Henry, T.J. 2017. Biodiversity of Heteroptera. In: Footitt, R.G. & Adler, P.H. (Eds.). *Insect Biodiversity*. 2 ed. Hoboken, John Wiley. v. 1, p. 279-335.
- Henry, T.J.; Perez-Gelabert, D.E.; Jr, W.E.S. & Heiss, E. 2013. *Brachyrhynchus membranaceus* (Fabricius), an Old World flat bug (Hemiptera: Heteroptera: Aradidae: Mezirinae) newly discovered in the Western Hemisphere. *Proceedings of the Entomological Society of Washington*, 115: 342-348. [DOI](#)
- Jacobs, D.H. & Nesar, S. 2005. *Thaumastocoris australicus* Kirkaldy (Heteroptera: Thaumastocoridae): a new insect arrival in South Africa, damaging to *Eucalyptus* tree. *South African Journal of Science*, 101: 233-236.
- Kun, M.E. & Masciocchi, M. 2019. First detection of the cosmopolitan invader *Leptoglossus occidentalis* Heidemann (Heteroptera: Coreidae) in Argentina. *Anais da Academia Brasileira de Ciências*, 91: e20180493. [DOI](#)
- Lee, D.-H. 2015. Current status of research progress on the biology and management of *Halyomorpha halys* (Hemiptera: Pentatomidae) as an invasive species. *Applied Entomology and Zoology*, 50(3): 277-290. [DOI](#)
- Lee, D.-H.; Short, B.D.; Joseph, S.V.; Bergh, J.C. & Leskey, T.C. 2013. Review of the biology, ecology, and management of *Halyomorpha halys* (Hemiptera: Pentatomidae) in China, Japan, and the Republic of Korea. *Environmental Entomology*, 42(4): 627-641. [DOI](#)
- Lesieur, V.; Lombaert, E.; Guillemaud, T.; Courtial, B.; Strong, W.; Roques, A. & Auger-Rozenberg, M.A. 2019. The rapid spread of *Leptoglossus occidentalis* in Europe: a bridgehead invasion. *Journal of Pest Science*, 92(1): 189-200. [DOI](#)
- Liebholt, A.M.; Yamanaka, T.; Roques, A.; Augustin, S.; Chown, S.L.; Brockerhoff, E.G. & Pyšek, P. 2018. Plant diversity drives global patterns of insect invasions. *Scientific Reports*, 8(12095): 1-5. [DOI](#)
- Machado, D. do N.; Costa, E.C.; Guedes, J.V.C.; Barbosa, L.R.; Martínez, G.; Mayorga, S.I.; Ramos, S.O.; Branco, M.; Garcia, A.; Vanegas-Rico, J.M.; Jiménez-Quiroz, E.; Laudonia, S.; Novoselsky, T.; Hodel, D.R.; Arakelian, G.; Silva, H.; Perini, C.R.; Valmorbidia, I.; Ugalde, G.A. & Arnemann, J.A. 2020. One maternal lineage leads the expansion of *Thaumastocoris peregrinus* (Hemiptera: Thaumastocoridae) in the New and Old Worlds. *Scientific Reports*, 10: 1-12. [DOI](#)
- McPherson, J.E. 2018. *Invasive stink bugs and related species (Pentatomoidea): Biology, higher systematics, semiochemistry, and management*. In: McPherson, J.E. (Ed.). *Contemporary topics in entomology*. Boca Raton, FL, CRC Press. 819p.
- McPherson, J.E. & McPherson, R. 2000. *Stink bugs of economic importance in America North of Mexico*. Boca Raton, FL, CRC Press. 253p.
- Nadel, R.L. & Noack, A.E. 2012. Current understanding of the biology of *Thaumastocoris peregrinus* in the quest for a management strategy. *International Journal of Pest Management*, 58(3): 257-266. [DOI](#)
- Nascimento-Machado, D. do; Corrêa-Costa, E.; Perini, C.R.; Andrade-Ugalde, G.; Alves-Saldanha, M.; Leitão, J.V.; Lovato-Colpo, T.; Arnemann, J.A. & Cordero-Rivera, A. 2019. The ongoing dispersion of the *Eucalyptus* bronze bug (*Thaumastocoris peregrinus*) in Spain. *Forest Systems*, 28: eS03. [DOI](#)
- Nielsen, A.L. & Hamilton, G.C. 2009. Life history of the invasive species *Halyomorpha halys* (Hemiptera: Pentatomidae) in Northeastern United States. *Annals of the Entomological Society of America*, 102: 608-616. [DOI](#)
- Noack, A.E. & Coviella, C.E. 2006. *Thaumastocoris australicus* Kirkaldy (Hemiptera: Thaumastocoridae): first record of this invasive pest of *Eucalyptus* in the Americas. *General and Applied Entomology*, 35: 13-14.
- Noack, A.E.; Cassis, G. & Rose, H.A. 2011. Systematic revision of *Thaumastocoris* Kirkaldy (Hemiptera: Heteroptera: Thaumastocoridae). *Zootaxa*, 3121: 1-60. [DOI](#)
- Palumbo, J.C.; Perring, T.M.; Millar, J.G. & Reed, D.A. 2016. Biology, ecology, and management of an invasive stink bug, *Bagrada hilaris*, in North America. *Annual Review of Entomology*, 61: 453-473. [DOI](#)
- Panizzi, A.R. 2015. Growing problems with stink bugs (Hemiptera: Heteroptera: Pentatomidae): species invasive to the U.S. and potential Neotropical invaders. *American Entomologist*, 61: 223-233. [DOI](#)
- Perez-Gelabert, D.E.; Abud-Antún, A. & Nolasco, C. 2019. First record of the introduced Asian Plataspid *Brachyplatys subaeneus* (Westwood) (Hemiptera: Heteroptera: Plataspidae) in the Dominican Republic. *Proceedings of the Entomological Society of Washington*, 121: 135-138. [DOI](#)
- Pinzón-Florián, O.P. 2020. Pests management in Colombian forest plantations. In: Estay, S.A. (Ed.). *Forest pest and disease management in Latin America: Modern perspectives in natural forests and exotic plantations*. Cham, Springer International Publishing. p. 145-169. [DOI](#)
- Pyšek, P. & Richardson, D.M. 2010. Invasive Species, Environmental Change and Management, and Health. *Annual Review of Environment and Resources*, 35: 25-55. [DOI](#)
- Pyšek, P.; Hulme, P.E.; Simberloff, D.; Bacher, S.; Blackburn, T.M.; Carlton, J.T.; Dawson, W.; Essl, F.; Foxcroft, L.C.; Genovesi, P.; Jeschke, J.M.; Kühn, I.; Liebhold, A.M.; Mandrak, N.E.; Meyerson, L.A.; Pauchard, A.; Pergl, J.; Roy, H.E.; Seebens, H.; Kleunen, M.v.; Vilà, M.; Wingfield, M.J. & Richardson, D.M. 2020. Scientists' warning on invasive alien species. *Biological Reviews*, 95(6): 1511-1534. [DOI](#)
- Rabitsch, W. 2008. Alien True Bugs of Europe (Insecta: Hemiptera: Heteroptera). *Zootaxa*, 1827: 1-44. [DOI](#)
- Rabitsch, W. 2010. True Bugs (Hemiptera, Heteroptera). Chapter 9.1. *BioRisk*, 4: 407-433. [DOI](#)
- Rédei, D. 2016. The identity of the *Brachyplatys* species recently introduced to Panama, with a review of bionomics (Hemiptera: Heteroptera: Plataspidae). *Zootaxa*, 4136: 141-154. [DOI](#)
- Rengifo-Correa, L.A. & González, R. 2011. Lygaeoidea (Hemiptera: Heteroptera) de Parques Nacionales Naturales (PNN) con nuevos registros para Colombia. *Revista Colombiana de Entomología*, 37(2): 331-340.
- Ruberson, J.R.; Takasu, K.; David Buntin, G.; Eger, J.E.; Gardner, W.A.; Greene, J.K.; Jenkins, T.M.; Jones, W.A.; Olson, D.M.; Roberts, P.M.; Suiter, D.R. & Toews, M.D. 2013. From Asian curiosity to eruptive American pest: *Megacopta cribraria* (Hemiptera: Plataspidae) and prospects for its biological control. *Applied Entomology and Zoology*, 48: 3-13. [DOI](#)
- Schaefer, C.W. 1988. The food plants of some "primitive" Pentatomoidea (Hemiptera: Heteroptera). *Phytophaga*, 2: 19-45.
- Schaefer, C.W. & Panizzi, A.R. 2000. *Heteroptera of economic importance*. Boca Raton, CRC Press. 852p.
- Schuh, R.T. & Weirauch, C. 2020. *True bugs of the world (Hemiptera: Heteroptera). Classification and natural history*. 2 ed. Manchester, UK, Siri Scientific Press. 768+32 plates (Monograph Series, 2).
- Scudder, G.G.E. & Footitt, R.G. 2006. Alien true bugs (Hemiptera: Heteroptera) in Canada: composition and adaptations. *The Canadian Entomologist*, 138: 24-51. [DOI](#)
- Simberloff, D.; Martin, J.-L.; Genovesi, P.; Maris, V.; Wardle, D.A.; Aronson, J.; Courchamp, F.; Galil, B.; García-Berthou, E.; Pascal, M.; Pyšek, P.; Sousa, R.; Tabacchi, E. & Vilà, M. 2013. Impacts of biological invasions: what's what and the way forward. *Trends in Ecology & Evolution*, 28: 58-66. [DOI](#)

-
- Souza, G.K.; Pikart, T.G.; Pikart, F.C.; Serrão, J.E.; Wilcken, C.F. & Zanuncio, J.C. 2012. First record of a native Heteropteran preying on the introduced *Eucalyptus* pest, *Thaumastocoris peregrinus* (Hemiptera: Thaumastocoridae), in Brazil. *Florida Entomologist*, 95(2): 517-520. [DOI](#)
- Stork, N.E. 2018. How many species of insects and other terrestrial arthropods are there on earth? *Annual Review of Entomology*, 63: 31-45. [DOI](#)
- van der Heyden, T. & Faúndez, E. 2020. First records of *Leptoglossus occidentalis* Heidemann (Heteroptera: Coreidae: Coreinae) in Brazil and South Africa. *Boletín del Museo Nacional de Historia Natural del Paraguay*, 24: 28-30.
- Weirauch, C.; Alvarez, C. & Zhang, G. 2012. *Zelus renardii* and *Z. tetracanthus* (Hemiptera: Reduviidae): biological attributes and the potential for dispersal in two assassin bug species. *Florida Entomologist*, 95(3): 641-649. [DOI](#)
- Zhang, G.; Hart, E. & Weirauch, C. 2016. A taxonomic monograph of the assassin bug genus *Zelus* Fabricius (Hemiptera: Reduviidae): 71 species based on 10,000 specimens. *Biodiversity Data Journal*, 4: e8150. [DOI](#)