

HERPETOFAUNA OF THE QUATERNARY SAND DUNES OF THE
MIDDLE RIO SÃO FRANCISCO: BAHIA: BRAZIL. VII. *TYPHLOPS*
AMOIPIRA SP. NOV., A POSSIBLE RELATIVE OF *TYPHLOPS*
YONENAGAE (SERPENTES, TYPHLOPIDAE)

MIGUEL TREFAUT RODRIGUES¹
FLORA ACUNÃ JUNCA²

ABSTRACT

A new species of a small typhlopoid snake is described from Ibiraba, in the sand- dune area of the left bank of Rio São Francisco, State of Bahia, Brazil. Typhlops amoipira sp. nov. is a small, light brown, and slightly pigmented Typhlops characterized by an incomplete nasal suture, 18 scale rows around the body (SAB), and 212-242 dorsal scales. The geographic and morphologically closer species, Typhlops yonenagae (18 SAB, 259-291 dorsals) lives in the same area, in the sands of the opposite side of the river.

KEYWORDS: Serpentes, Typhlopidae, *Typhlops amoipira*, Brazil, Caatingas, Taxonomy.

RESUMO

Descreve-se uma nova espécie de Typhlops de Ibiraba, no campo de dunas da margem esquerda do Rio São Francisco, Bahia, Brasil. Typhlops

¹ Universidade de São Paulo, Instituto de Biociências, Departamento de Zoologia, Caixa Postal 11.461, CEP 05422-970, São Paulo, SP, Brasil.

² Universidade Estadual de Feira de Santana, Departamento de Ciências Biológicas, Av. Universitária, s/n, Campus Universitário, CEP 44031-060, Feira de Santana, BA, Brasil.

amoipira sp. nov. é um pequeno tiflopídeo castanho claro, pouco pigmentado, caracterizado por apresentar 18 fileiras de escamas ao redor do corpo e 212 a 242 dorsais. A espécie geográfica e morfológicamente mais próxima, *Typhlops yonenagae*, ocorre na mesma área, nas areias da margem oposta do rio e, embora também tenha 18 fileiras de escamas ao redor do corpo, apresenta 259 a 291 escamas dorsais.

PALAVRAS-CHAVE: Serpentes, Typhlopidae, *Typhlops amoipira*, Brazil, Caatingas, Taxonomia.

INTRODUCTION

The sand dune area crossed by the middle Rio São Francisco in the Brazilian domain of semiarid Caatingas, State of Bahia, has emerged in the last years as an unparalleled case of an extraordinary newly discovered squamate fauna. The uniqueness of this example of “new age’s discovery” (*sensu* Donoghue and Alverson, 2000), derives from the study at a smaller spacial scale of the presumably well-known Caatingas domain. Most squamates living within the 800.000 km² area covered by the dry forest caatingas, were considered widespread in the domain and the whole fauna relatively well known (Vanzolini, 1974). Overall caatinga faunal predictability was considered high and after intensive field studies (summary in Vitt, 1995, and references cited) a comprehensive book on reptiles from the area was published (Vanzolini *et al.*, 1980). Occasionally, one or two new species were described after intensive surveys across large areas. These new species, most of them cryptic or very similar to previously known relatives, were only recognized after detailed comparisons with material from museum shelves (see Rodrigues, 1986 for an example).

Work in the dune region of Rio São Francisco have shown a remarkable contrast to the picture of the caatinga squamate fauna as a well-known and highly-predictable faunistic component. From a sandy area with about 7.000 km² surrounding the banks and adjacent areas of Rio São Francisco, four new genera and a total of 20 new species of squamates (32% of the fauna, most endemic) were described (Rodrigues, 1984a,b,c, 1986, 1991a,b,c,d, 1993a,b, 1996; Rodrigues *et al.*, 1988; Vanzolini, 1991a,b). New species of other vertebrates have been also described from there (Rocha, 1995, a new rodent; Lencioni-Neto, 1994, a new bird).

Some of the new genera and most of the new species revealed adaptations to fossoriality or to life in sand unsuspected to exist among South American squamates or rodents (Rocha, 1995; Rodrigues, 1996). Furthermore, some endemic squamates likely form vicariant allopatric sister species pairs occupy-

ing adjacent and opposite sand areas along the river (Rodrigues, 1996). As among these species pairs are psamophilic lizards, snakes and amphisbaenians represented, this fauna became the focus of ecological, biogeographical, and integrative evolutionary studies (Martins, 1995, 1997; Rodrigues, 1995, 1996; Renous *et al.*, 1995, 1999; Yonenaga-Yassuda *et al.*, 1996; Rocha, 1998; Pellegrino *et al.*, 1999, 2001; Passoni *et al.*, 2000; Benozzatti and Rodrigues, 2002).

Surveys continue in the area and are revealing the existence of additional undescribed lizards, snakes, amphisbaenians, and frogs. Most of the invertebrate fauna yet remains largely unexplored.

We report here on a new species of a fossorial snake, genus *Typhlops*, obtained in a recent trip. The only species of the genus previously known from the area is the endemic *Typhlops yonenagae* Rodrigues, 1991, described from Santo Inácio, in the sands of the right bank of the Rio São Francisco. The new species, obtained in the sands of the opposite side of the river, almost in front of Santo Inácio, differs clearly from *Typhlops yonenagae* but share several features with the latter as to lead us to conclude that it could form another example of vicariant species pair for the area (see Rodrigues, 1996 for other examples of species pairs of squamates in the area).

SPECIES DESCRIPTION

***Typhlops amoipira* sp. nov.**

Holotype: MZUSP 12298, Ibiraba (10°48'S, 42°50'W): state of Bahia: Brazil, collected by Flora Acuña Juncá on 5 March 2000; field number 200170.

Paratypes: MZUSP 12299-12303, 6-9.iii.2000; other data as for the holotype.

Etymology: A homage to the extinct Amoipira indians, a Tupi tribe that lived in the area in the XVI century and probably went extinct by the end of the XVII century (Souza, 1942; Niemendadju, 1946). By coincidence, *amoipira*, in Tupi, also means "the relative from the other side" (Souza, 1942).

Diagnosis: A small, slightly pigmented *Typhlops* with incomplete nasal suture, 18 scales around the body (without reduction), and 212-242 dorsals.

Description of the holotype: (Figs 1-3). Rostral broad, longer than wide, extending about two thirds on dorsal face of head, with a pronounced narrowing at nostril level; narrowest at labial level and touching labial border. Posterior margin of rostral rounded, reaching the level of the anterior margin of eye. Nasal long, convex anteriorly, concave posteriorly, imbricating and contacting




Figure 1. *Typhlops amoipira*, sp. nov.: lateral, ventral, and dorsal views of head (MZUSP 12298, holotype)



Figure 2. *Typhlops amoipira* (MZUSP 12299, paratype) from Ibiraba, State of Bahia, Brazil.



Figure 3. *Typhlops amoipira* (MZUSP 12299, paratype) from Ibiraba, State of Bahia, Brazil.

rostral, preocular, first and second supralabials, and frontal. Nasal wider than frontal and wider at nostril level, narrower posteriorly. Nostril in the posterior part of nasal, at the level of inferior part of ocular. Nasal suture incomplete, occupying the inferior third of nasal and coincident with suture between first and second labials. Frontal as long as large, cicloid, contacting rostral and preventing midline contact between nasals. Following frontal two imbricate and cicloid scales, the first identical to frontal in size and shape, the second wider. A pair of longer than wide distinctive supraoculars, diagonally oriented, minimally separated by frontal at midline. Supraocular covers the superior part of ocular; anterior part of supraocular imbricates under part of frontal, nasal, and preocular. Preocular large, wider at nostril level, contacting second and third supralabials, covering anterior part of ocular and supraocular, and reaching the

Table 1. Measurements (mm) of the type series of *Typhlops amoipira*, sp. nov. (* = holotype)

MZUSP number	Total length	Tail length	Diameter at midbody	Head width	Eye diameter
12298*	208	4	7.5	5.2	2.0
12299	147	3	6.1	3.0	1.8
12300	142	3	5.2	3.2	1.8
12301	141	3	5.1	3.5	1.8
12302	73	2	3.2	2.5	1.6
12303	155	4	6.3	3.5	2.0

eye. Preocular indenting deeply the suture between second and third supralabials; ventrally reaching about the same level of preocular. Ocular larger than preocular, in contact with third supralabial with the eye in the superior two thirds of scale. Posterior to ocular, ventrally, a small, higher than wide temporal; dorsally, in larger contact, two identical and successively imbricate wider than long "parietals". The first "parietal" contacts supraocular. Three supralabials, first smallest, third the largest. Third supralabial followed posteriorly by a larger scale which contacts the base of ocular and temporal. Three infralabials.

Dorsals smooth, cicloidal, slightly larger than long, in 18 longitudinal rows, without reduction; 212-242 transversal rows between rostral and tip of tail; 7-12 subcaudals.

Ground color creamy brown, few pigmented. In the anterior dorsal part of body and tail, a fine darker brown reticulum, particularly concentrated in the central part of dorsal head scales. In juveniles, an inconspicuous longitudinal mid-dorsal, one scale wide, brown line extends from posterior part of head to the first third of body; the line disappears progressively towards tail. Snout, imbricate part of dorsal head scales, labials, and the rest of body, immaculate.

Measurements of the type series are shown in Table 1.

DISCUSSION

Typhlops is a large and poorly known genus of pantropical fossorial snakes for which there is no available phylogenetically oriented studies of relationships. For this reason, we based our comparisons to South American species. Data on scale-count variation and maximum snout-vent length was taken from specimens examined at the MZUSP collection, and complemented by data from Dixon and Hendricks (1979) and Rodrigues (1991d).

Typhlops amoipira shares the presence of an incomplete nasal suture with five other South American species of *Typhlops*: *T. reticulatus*, *T. brongersmianus*, *T. minuisquamus*, *T. paucisquamus*, and *T. yonenagae*. *Typhlops lehneri*, from northern Venezuela, the only other known South Ameri-

has a complete nasal suture as the other Caribbean and Central-American congeners. *Typhlops brongersmianus* and *T. reticulatus* are larger species (maximum length, respectively, 320 mm and 522 mm – against a maximum of 200 mm in *T. amoipira*) and have 20 scales without reduction around midbody. *Typhlops brongersmianus* is a widespread species in South America, occurring in almost all habitats; it is largely sympatric with *T. reticulatus* in Amazonia, which is also known from an isolated population in northeastern Brazil. *Typhlops minuisquamus* is a large, sharply bicolored species (maximum size 361 mm) restricted to Amazonia, and has 18 SAB with scale row reduction (20 to 14, 18 to 14 or 18 to 12). *Typhlops paucisquamus* is restricted to the Atlantic forest of northeastern Brasil. Although *T. paucisquamus*, *T. yonenagae*, and *T. amoipira* have similar sizes (maximum of 200 mm) and 18 SAB, the former has the lowest dorsal counts in the genus (169-182). The only differences between *T. yonenagae* and *T. amoipira* are the total number of dorsal scales (212-242 in *T. amoipira*; 259-291 in *T. yonenagae*) and color pattern. Specimens of *T. yonenagae* are creamy brown, with a conspicuous middorsal line along body extending from posterior part of head to tail. Specimens of *T. amoipira* are less pigmented and the line, mostly inconspicuous, is present only in the anterior third of body in juveniles.

Even considering the absence of strong evidence, morphological similarity, ecological preferences, and the geographic distribution of *T. amoipira* and *T. yonenagae* make their close relationship very probable. If this possibility is confirmed, it will be interesting to know if they diverged contemporaneously with the other pairs of vicariant species in the area.

The type series of *Typhlops amoipira* was obtained in a grid of 100 pitfall traps consisting of 10x10 parallel lines of 30 liter buckets, each one with three 1.5 m drift fences. Distance between parallel lines was 7.5 m. The grid was placed in a sandy and flat area bordering a large sand dune, close to the river. Total effort was equivalent to 600 pitfalls/day, for a sampled area of about 5000 m². The six specimens were taken from buckets early in the morning (see Rodrigues, 1991a, 1996, for a detailed description of the area). The presence of a juvenile in the sample obtained suggests that egg laying most likely occurred during the dry season which extends from November to March in this region (see Rodrigues, 1996).

Material of *Typhlops yonenagae* examined: MZUSP 10086, 10087, 12497-12506, Santo Inácio, Bahia, Brazil.

ACKNOWLEDGEMENTS

We thank Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP), Conselho Nacional de Desenvolvimento Científico e Tecnológico

(CNPq) and Universidade Estadual de Feira de Santana for support. For their invaluable help in the field, we thank Felipa and the people of Icatú (Ibiraba officially), and Pedro L. B. Rocha. We also thank Luciana Lobo for the drawing, and Hussam Zaher and Andrew Check for reading the manuscript.

REFERENCES

- Benozzati, M.L. & M.T. Rodrigues. 2003. Mitochondrial DNA phylogeny of a Brazilian group of eyelid-less gymnophthalmid lizards, and the origin of the unisexual *Gymnophthalmus underwoodi* from Roraima (Brazil). *J. Herp.* (in press).
- Dixon, J.R. & F.S. Hendricks. 1979. The wormsnakes (family Typhlopidae) of the neotropics, exclusive of the Antilles. *Zool. Verhand.* 173:39pp.
- Donoghue, M.J. & W.S. Alverson. 2000. A new age of discovery. *Ann. Missouri Bot. Gard.* 87:110-126.
- Lencioni-Neto, F. 1994. Une nouvelle espèce de *Chordeiles* (Aves, Caprimulgidae) de Bahia (Brésil). *Alauda* 62(4):241-245.
- Nimuendajú, K. 1946. Mapa das tribos indígenas do Brasil, in (J. H. Steward, edit.) *Handbook of South American Indians*, Vol. 1. Bureau of American Ethnology. Bull. Smiths. Inst. 143.
- Martins, J.M. 1995. Allozyme variation and expression in lizards of the *Tropidurus nanuzae* species group (Iguania: Tropiduridae). *Copeia* 1995:665-675.
- Martins, J.M. 1997. Estudo alozímico de um grupo de lagartos gimnofthalmídeos (Squamata: Gymnophthalmidae). Unpublished PhD Thesis, Universidade de São Paulo, Brazil.
- Passoni, J.C.; M.L. Benozzati & M.T. Rodrigues. 2000. Mitochondrial DNA polymorphism and heteroplasmy in populations of three species of *Tropidurus* of the *nanuzae* group (Squamata, Tropiduridae). *Genet. Mol. Biol.* 23(2):351-356.
- Pellegrino, K.C.M., M.T. Rodrigues & Y. Yonenaga-Yassuda. 1999. Chromosomal polymorphisms due to supernumerary chromosomes and pericentric inversions in the eyelid-less microteiid lizard *Nothobachia ablephara* (Squamata, Gymnophthalmidae). *Chromosome Research* 7(4):247-254.
- Pellegrino, K.C.M., M.T. Rodrigues, Y. Yonenaga-Yassuda & J. 2001. A molecular perspective on the evolution of microteiid lizards (Squamata, Gymnophthalmidae), and a new classification for the family. *Biol. Jour. Lin. Soc.* 74:315-338.
- Renous, S.; E. Höfling & J.P. Gasc. 1995. Analysis of the locomotion pattern of two microteiid lizards with reduced limbs, *Calyptommatus leirolepis* and *Nothobachia ablephara* (Gymnophthalmidae). *Zoology* 99(1995):21-38.
- Renous, S., E. Höfling & J.P. Gasc. 1999. On the rhythmical coupling of the axial and appendicular systems in small terrestrial lizards (Sauria: Gymnophthalmidae). *Zoology* 102(1999):31-49.
- Rocha, P.L.B. 1995. *Proechimys yonenagae*, a new species of spiny rat (Rodentia: Echimyidae) from the fossil sand dunes in the Brazilian Caatinga. *Mammalia* 59(4):537-549.
- Rocha, P.L.B. 1998. Uso e partição de recursos por uma comunidade de lagartos das dunas interiores do Rio São Francisco (Squamata). Unpublished PhD Thesis, Universidade de São Paulo, Brazil.
- Rodrigues, M.T. 1984a. *Nothobachia ablephara*: novo gênero e espécie do nordeste do Brasil (Sauria: Teiidae). *Papéis Avulsos Zool., S. Paulo* 35(28):361-366.
- Rodrigues, M.T. 1984b. Uma nova espécie brasileira de *Tropidurus* com crista dorsal (Sauria, Iguanidae). *Papéis Avulsos Zool., S. Paulo* 35(16):169-175.
- Rodrigues, M.T. 1984c. Sobre *Platynotus* Wagler, 1890, pré-ocupado, substituído por *Tapinurus* Amaral, 1933, com a descrição de uma nova espécie (Sauria, Iguanidae). *Papéis Avulsos Zool., S. Paulo* 35(29):367-373.

- Rodrigues, M.T. 1986. Um novo *Tropidurus* com crista dorsal do Brasil com comentários sobre suas relações, distribuição e origem (Sauria; Iguanidae). *Papéis Avulsos Zool.*, S. Paulo 36(17):171-179.
- Rodrigues, M.T. 1991a. Herpetofauna das dunas interiores do rio São Francisco: Bahia: Brasil. I. Introdução à área e descrição de um novo gênero de microteídeos (*Calyptommatius*) com notas sobre sua ecologia, distribuição e especiação (Sauria, Teiidae). *Papéis Avulsos Zool.*, S. Paulo 37(19):285-320.
- Rodrigues, M.T. 1991b. Herpetofauna das dunas interiores do rio São Francisco: Bahia: Brasil. II. *Psilophthalmus*: um novo gênero de microteídeos sem pálpebra (Sauria, Teiidae). *Papéis Avulsos Zool.*, S. Paulo 37(20):321-327.
- Rodrigues, M.T. 1991c. Herpetofauna das dunas interiores do rio São Francisco: Bahia: Brasil. III. *Procellosaurinus*: um novo gênero de microteídeo sem pálpebra, com a redefinição do gênero *Gymnophthalmus* (Sauria, Teiidae). *Papéis Avulsos Zool.*, S. Paulo 37(21):329-342.
- Rodrigues, M.T. 1991d. Herpetofauna das dunas interiores do rio São Francisco: Bahia: Brasil. IV. Uma nova espécie de *Typhlops* (Ophidia, Typhlopidae). *Papéis Avulsos Zool.*, S. Paulo 37(22):343-346.
- Rodrigues, M.T. 1993a. Herpetofauna das dunas interiores do rio São Francisco: Bahia: Brasil. V. Duas novas espécies de *Apostolepis* (Serpentes, Colubridae). *Mem. Inst. Butantan* 54(2):53-59 (1992).
- Rodrigues, M.T. 1993b. Herpetofauna of palaeoquaternary sand dunes of the middle São Francisco river: Bahia: Brazil. VI. Two new species of *Phimophis* (Serpentes: Colubridae) with notes on the origin of psammophilic adaptations. *Papéis Avulsos Zool.*, S. Paulo 38(11):187-198.
- Rodrigues, M.T. 1995. Filogenia e história geográfica de uma radiação de lagartos microteídeos (Sauria, Teiioidea, Gymnophthalmidae). Unpublished Thesis (Livre- docência), Universidade de São Paulo, Brazil.
- Rodrigues, M.T. 1996. Lizards, snakes and amphisbaenians from the Quaternary sand dunes of the middle Rio São Francisco: Bahia: Brazil. *Jour. Herpetol.* 30(4):513-523.
- Rodrigues, M.T., S. Kasahara & Y. Yonenaga-Yassuda. 1988. *Tropidurus psammonastes*: uma nova espécie do grupo *torquatus* com notas sobre seu cariótipo e distribuição (Sauria, Iguanidae). *Papéis Avulsos Zool.*, S. Paulo 36(26):307-313.
- Souza, G.S. 1942. *Notícia do Brasil*. 7th. Ed., 2 Vols. Biblioteca Histórica Brasileira XVI. Livraria Martins Fontes Editora, S. Paulo.
- Vanzolini, P.E. 1974. Ecological and geographical distribution of lizards in Pernambuco, northeastern Brazil (Sauria). *Papéis Avulsos Zool.*, S. Paulo 28(4):61-90.
- Vanzolini, P.E. 1991a. Two new small species of *Amphisbaena* from fossil dune field of the middle Rio São Francisco, State of Bahia, Brasil (Reptilia, Amphisbaenia). *Papéis Avulsos Zool.*, S. Paulo 37(17):259-276.
- Vanzolini, P.E. 1991b. Two further new species of *Amphisbaena* from the semi-arid northeast of Brasil (Reptilia, Amphisbaenia). *Papéis Avulsos Zool.*, S. Paulo 37(23):247-361.
- Vanzolini, P.E., A.M.M. Ramos-Costa & L.J. Vitt. 1980. Répteis das Caatingas. Academia Brasileira de Ciências, Rio de Janeiro.
- Vitt, L.J. 1995. The ecology of tropical lizards in the caatinga of northeast Brazil. *Occ. Pap. Oklahoma. Mus. Nat. Hist.* 1:1-29.
- Yonenaga-Yassuda, Y., L. Mori, T.H. Chu & M.T. Rodrigues. 1996. Chromosomal banding patterns in the eyelid-less microteiid radiation: *Procellosaurinus* and *Vanzosaura* (Squamata, Gymnophthalmidae). *Cytogenetics and Cell Genetics* 74:203-210.



Papéis Avulsos de Zoologia

EDITORIAL COMMITTEE

Editor-in-Chief. Hussam Zaher, Serviço de Vertebrados, Museu de Zoologia, Universidade de São Paulo, Caixa Postal 42594, 04299-970, São Paulo, SP, Brazil. E-mail: hzaher@ib.usp.br.

Associated Editors. Mario C. C. de Pinna (Universidade de São Paulo, Brazil), Antonio C. Marques (Universidade de São Paulo, Brazil), Sergio A. Vanin (Universidade de São Paulo, Brazil).

Editorial Board. Adriano Kury (Museu Nacional do Rio de Janeiro, Brasil), Aziz N. Ab-Saber (Universidade de São Paulo, Brazil), Carlos Roberto F. Brandão, (Universidade de São Paulo, Brazil), Christian de Muizon (Museum National d'Histoire Naturelle, France), Darrel Frost (American Museum of Natural History, U.S.A.), Gerardo Lamas (Museu Javier Prado de Lima, Peru), H. R. Heyer (National Museum of Natural History, U.S.A.), James Carpenter (American Museum of Natural History, U.S.A.), James Patton (University of Berkeley, U.S.A.), John Maisey (American Museum of Natural History, U.S.A.), Marcos Raposos (Museu Nacional do Rio de Janeiro, Brazil), Marcos Tavares (Universidade Santa Ursula, Brazil), Mario de Vivo (Universidade de São Paulo, Brazil), Miguel T. U. Rodrigues (Universidade de São Paulo, Brazil), Naércio Menezes (Universidade de São Paulo, Brazil), Nelson Papavero (Museu Paraense Emilio Goeldi, Brazil), Olivier Rieppel (Field Museum of Natural History, U.S.A.), Paulo E. Vanzolini (Universidade de São Paulo, Brazil), Paulo Young (Museu Nacional do Rio de Janeiro, Brazil), Ralf Holzentahl (University of Minesotta, U.S.A.), Randahl Schuh (American Museum of Natural History, U.S.A.), Ricardo Macedo Correa e Castro (Universidade de São Paulo, Brasil), Richard Prum (University of Kansas, U.S.A.), Richard Vari (National Museum of Natural History, U.S.A.), Rudiger Bieler (Field Museum of Natural History, U.S.A.), Ubirajara Martins (Universidade de São Paulo, Brasil), Walter A. P. Boeger (Universidade Federal do Paraná, Brasil).

INSTRUCTIONS FOR AUTHORS

General Information. *Papéis Avulsos de Zoologia (PAZ)* covers primarily the fields of Zoology, publishing original contributions in systematics, paleontology, evolutionary biology, ecology, taxonomy, anatomy, behavior, functional morphology, molecular biology, ontogeny, faunistic studies and biogeography. *PAZ* also encourages submission of theoretical and empirical studies that explore principles and methods of systematics.

All contributions must follow the International Code of Zoological Nomenclature and relevant specimens should be properly curated and deposited in a recognized public or private, non-profit institution. Tissue samples should be referred to their voucher specimens and all nucleotide sequence data (aligned as well as unaligned) should be submitted to Genbank (<http://www.ncbi.nlm.nih.gov/genbank/>) or EMBL (<http://www.ebi.ac.uk/>).

Peer Review. All submissions to *PAZ* are subject to review by at least two referees and the Editor-in-Chief. Three legible copies (including photocopies of original illustrations) and original illustrations must be submitted; all authors will be notified of submission date. Authors may suggest potential reviewers. Communications regarding acceptance or rejection of manuscripts are made through correspondence with the first or corresponding author only. Once a manuscript is accepted providing changes suggested by the reviewers, the author is requested to return a revised version incorporating those changes (or a detailed explanation of why reviewer's suggestions were not followed) in four weeks upon receiving the communication by the editor. Revised manuscripts must be submitted as both hard copy and electronic file (3.5" disk, Zip Drive, or CD-Rom with text in Microsoft Word format).

Proofs. Page-proofs with the revised version will be sent to the first or corresponding author. Page-proofs must be returned to the editor in two weeks, preferably within 48 hours. Failure to return the proof promptly may be interpreted as approval with no changes and/or may delay publication. Only necessary corrections in proof will be permitted. Once page proof is sent to the author, further alterations and/or significant additions of text are permitted only at the author's expense or in the form of a brief appendix ("note added in proof").

Submission of Manuscripts. Manuscripts should be sent to the Editor-in-Chief (H. Zaher, Museu de Zoologia da USP, Caixa Postal 42694, CEP 04299-970, São Paulo, SP, Brasil). Manuscripts are considered on the understanding that they have not been published or will not appear elsewhere in substantially the same or abbreviated form. The criteria for acceptance of articles are quality and relevance of research, clarity of text, and compliance with the guidelines for manuscript preparation.

Manuscripts should be written preferentially in English, but texts in Portuguese or Spanish will also be considered. Studies with a broad coverage are encouraged to be submitted in English. All manuscripts should include abstracts in Portuguese and English regardless of the original language.

Authors are requested to pay attention to the instructions concerning the preparation of the manuscripts. Close adherence to the guidelines will expedite processing of the manuscript, whereas manuscripts deviating from the required form will be returned for revision prior to review.

Papéis Avulsos de Zoologia

Manuscript Form. Manuscripts should not exceed 100 pages of double-spaced typescript on 21 by 29.7 cm (A4 format) or 21.5 by 28 cm (letter format) paper, with wide margins. The pages of the manuscript should be numbered consecutively.

The text of articles should be arranged in the following order: title page, abstract, body of text, literature cited, tables, appendices, and figure captions. Each of these sections should begin on a new page. All typescript pages must be double-spaced.

- (1) *Title page:* This should include the title, author(s) name(s), institutions, and key words in English as well as in the language of the manuscript, and a short running title in English. The title should be concise and, where appropriate, should include mention of families and/or higher taxa. Names of new taxa should not be included in titles.
- (2) *Abstract:* All papers should have an abstract in English, regardless of the original language. The abstract is of great importance as it may be reproduced elsewhere. It should be in a form intelligible if published alone in conjunction with the title and should summarize the main facts, ideas, and conclusions of the article. Indicative abstracts are strongly discouraged. Include all new taxonomic names for referencing purposes. Abbreviations should be avoided. It should not include references. Abstracts should not exceed 350 words.
- (3) *Body of text:* The main body of the text should include the following sections: Introduction, Materials and Methods, Results, Discussion and Acknowledgments at end. Primary headings in the text should be in capital letters and centered; the following text should begin on the next line, indented. Secondary headings should be in capital and lowercase letters and flush left; the following text should begin on the next line, indented. Tertiary headings should be in capital and lower case letters, in italics and indented; the following text should be on the same line and separated from the heading by a hyphen.
- (4) *Literature cited:* Citations in the text should be given as: Silva (1998)..., Silva (1998: 14-20)..., Silva (1998: figs. 1, 2)..., Silva (1998a, b)..., Silva and Oliveira (1998)..., (Silva, 1998)..., (Rangel, 1890; Silva and Oliveira, 1998a, b; Adams, 2000)..., (Silva, pers. comm.)..., (Silva et al., 1998), the latter when the paper has three or more authors. The reference need not be cited when author and date are given only as authority for a taxonomic name. The literature section should be arranged strictly alphabetically and given in the following format:

Journal Article – Silva, H. R., H. Oliveira & S. Rangel. Year. Article title. Journal name, 00:000-000. Names of journals must be spelled out in full.

Books – Silva, H. R. Year. Book title. Publisher, Place, 000 pp.

Articles in Books – Silva, H. R. Year. Article title; pp. 000-000. In: H. Oliveira & S. Rangel (Eds.), Book Title. Publisher, Place.

Articles in Larger Works – Silva, H. R. Year. Article title; pp. 000-000. In: H. Oliveira & S. Rangel (Eds.), Title of Larger Work. Serial Publication 00. Publisher, Place.

Dissertations and Theses – Silva, H. R. Year. Dissertation title. Ph.D. dissertation, University, Place, 000 pp.

Tables. All tables must be numbered in the same sequence in which they appear in the text. Authors are encouraged to indicate where the tables should be placed in the text. They should be comprehensible without reference to the text. Tables should be formatted with horizontal, not vertical, rules. In the text, tables should be referred as Table 1, Tables 2 and 3, Tables 2-6. Use "TABLE" in the table heading.

Illustrations. Figures should be numbered consecutively, in the same sequence they appear in the text. Separate illustrations of a composite figure should be identified by capital letters and referred in the text as so (fig. 1A). Where possible, letters should be placed in the lower right corner of each illustration of a composite figure. Hand-written lettering on illustrations are unacceptable. Illustrations should be mounted on stout, white cardboard. Figures should be mounted in order to minimize blank areas between separate illustrations. High quality color or black and white photographs, and computer generated figures are preferable. Authors are encouraged to indicate where the figures should be placed in the text. Use "(Fig(s).)" and "Figure(s)" for referring to figures in the text, but "FIGURE(S)" in the figure captions and "(fig(s).)" when referring to figures in another paper.

For other details of manuscript preparation of format, consult the CBE Style Manual, available from the Council of Science Editors (<http://www.councilscienceeditors.org/pubs>).