

Papéis Avulsos de Zoologia

Museu de Zoologia da Universidade de São Paulo

Volume 57(11):137-139, 2017

www.mz.usp.br/publicacoes
www.revistas.usp.br/paz

ISSN impresso: 0031-1049
ISSN on-line: 1807-0205

FIRST RECORD OF A BREEDING COLONY OF MASKED BOOBY (*SULA DACTYLATRA* LESSON, 1831; SULIDAE) IN THE MAIN ISLAND OF THE ARCHIPELAGO OF FERNANDO DE NORONHA (PERNAMBUCO, BRAZIL)

ARIANE CAMPOS GOUVÊA^{1,2,4}
THAYNÁ MELLO³

ABSTRACT

*Fernando de Noronha Archipelago is one of the places with the highest richness of seabirds in Brazil; however, little information about the breeding biology of many species is available. Here we report a breeding colony of the Masked Booby (*Sula dactylatra*) in the main island of Fernando de Noronha, and present new data about the natural history and breeding biology of this species.*

KEY-WORDS: *Sula dactylatra*; Fernando de Noronha; Oceanic islands; Suliformes.

Fernando de Noronha is located in the South Equatorial Atlantic (03°51'S; 32°24'W), and the archipelago belongs to Pernambuco State. The Fernando de Noronha archipelago is the top of a huge submarine mountain with its base located about 4.000 m below sea level. It covers an area of 26 km² distributed in 21 islands and islets. The larger of the islands, Fernando de Noronha, covers 17 km² and is the only inhabited (IBAMA, 1990; Silva, 2008; Castro, 2009).

Fernando de Noronha holds the greatest richness of seabirds in the Brazil, being an important area of breeding and feeding for at least 11 species of seabirds (Schulz-Neto, 2004; Silva, 2008). Most studies of birds at Fernando de Noronha were on the occurrence of species or general inventories (Oren, 1982,

1984; Nacinovic & Teixeira, 1989; Antas *et al.*, 1990; Antas, 1991; Schulz-Neto, 1995, 2004) and, despite the richness of seabirds found in the archipelago, little information is available about biology, ecology, movements and other aspects of the natural history for most of the species, even for the most common and abundant found there.

The Masked Booby (*Sula dactylatra* Lesson, 1831) is the largest species of booby, measuring about 90 cm long and with wingspan of more than 150 cm. They have yellow bills and yellow legs, which get more strongly colored during the breeding season. They are strictly marine and pelagic, preferring deeper areas in relation to other species of boobies (Sick, 1997; Silva & Campos, 2006). They nest on flat areas of undergrowth and the nests are simple open spaces in

¹ Universidade de São Paulo (USP), Instituto de Biociências (IB). Rua do Matão, Travessa 14, nº 101, Butantã, CEP 05508-090, São Paulo, SP, Brasil.

² Universidade de São Paulo (USP), Museu de Zoologia (MZ). Avenida Nazaré, 481, Ipiranga, CEP 04263-000, São Paulo, SP, Brasil.

³ Instituto Chico Mendes de Conservação da Biodiversidade, Parna Marinho de Fernando de Noronha (PARNAMAR).

Alameda do Bodró, s/n, Boldró, CEP 53990-000, Fernando de Noronha, PE, Brasil. E-mail: thaynajm@gmail.com.

⁴ E-mail: arigouvea@gmail.com.

vegetation covered with excreta, eventually filled with small stones or shells (Schreiber & Hensley, 1976; Sick, 1997). Incubation lasts about 45 days and in about 120 days the chicks are flying. After this time, the young birds remain under the care of parents for about 150 days. Most adults persist in their breeding colony, but young and not reproductive adults disperse widely (Del Hoyo *et al.*, 1992; Sick, 1997). In Brazil, colonies of this species are found in Abrolhos, Atol das Rocas and Fernando de Noronha, breeding also on the island of Trindade, with occasional records in coastal areas from northeast to south of the Brazil (Del Hoyo *et al.*, 1992; Fonseca-Neto, 2004; Schulz-Neto, 2004; Silva & Campos, 2006; Hughes *et al.*, 2011).

The Masked Booby breeds only in the secondary islands of In Fernando de Noronha Archipelago. The main breeding colony is located on the Meio Island (300 nests). Other colonies were recorded in southwest of Rata Island (10 nests), Pontal da Macaxeira (200 nests) and Pontal Sul (10 nests), Ovos Island (70 nests) and Rasa Island (Antas *et al.*, 1990; Antas, 1991; Schulz-Neto, 2004; Silva, 2008). According to Antas (1991) due to human impacts, including the introduction of exotic predators, seabirds that breed on the main island would be only those that make their nests on trees or on cliffs, where access by predators would be more difficult. Thus, species of birds like the Masked Booby, which nests exclusively on the ground, would breed only in the secondary islands of the archipelago, where predator access would be more difficult (Antas, 1991).

However, there is a small, unrecorded breeding colony (Figure 1) located on the main island of Fernando de Noronha Archipelago, on the end of the Capim-açu trail (03°52'49.38"S; 32°27'29.47"W) (Figure 2). We found at least six breeding pair in



FIGURE 1: Breeding colony of *Sula dactylatra* located at the end of the Capim-açu trail, main island of Fernando de Noronha archipelago, September 2015. Photo: Deborah Gutierrez, 2015.




FIGURE 2: Map of Fernando de Noronha Archipelago, highlighting the location of the breeding colony of *Sula dactylatra*, located at the end of the Capim-açu trail (03°52'49.38"S; 32°27'29.47"W). Unscaled. Source: © 2016 google maps.

September 2015 and August 2016, thus representing a new breeding locality for this species. It is possible that this colony of the Masked Booby is a new occupation, once it had not been previously recorded by other researchers, despite being located near a trail of National Park. However, despite being a possible new nesting area, it is already threatened. Throughout the history of human occupation of the archipelago there were many types of anthropogenic pressures on birdlife, especially on the main island, and some of these problems caused by man persist until today, as in the case of the presence of introduced predators such as domestic cats (*Felis catus*), many already in feral state, brown rats (*Rattus norvegicus*), black rats (*Rattus rattus*), mice (*Mus musculus*) and lizards (*Salvator merianae*) (Soto, 2009). Currently these exotic species are found on the main island in alarming numbers and are considered potential predators of eggs, nestlings and even adults.

In September 2015, traces of *Felis catus* predation to *Sula dactylatra* nestlings in the Capim-açu colony were recorded, thus representing a real threat to the conservation of several migratory species, residents and even endemic birds. In this way, management actions and eradication of invaders species are urgent and essential for the conservation of this and other species of avifauna of the Fernando de Noronha Archipelago.

ACKNOWLEDGMENTS

We thank Jihad Daou and the ICMBIO from National Marine Park of Fernando de Noronha, especially Deborah Gutierrez and all volunteer staff.

REFERENCES

- ANTAS, P.T.Z. 1991. Status and conservation of seabirds breeding in Brazilian Waters. *ICBP Technical Publication*, 11:141-158.
- ANTAS, P.T.Z.; FILIPINI, A. & AZEVEDO-JUNIOR, S.M. 1990. Anilhamento de aves Oceânicas e/ou migratórias no Arquipélago de Fernando de Noronha em 1987 e 1988. In: Encontro Nacional de Anilhadores de Aves, 4^o. *Anais*. Recife, Universidade Federal Rural de Pernambuco, 1988. p. 13-17.
- CASTRO, J.W.A. 2009. Geologia Ambiental das Ilhas oceânicas de Trindade e Fernando de Noronha, Brasil. In: Mohr, L.V. *Ilhas Oceânicas Brasileiras: da Pesquisa ao Manejo*. Brasília, MMA/ICMBIO. v. 2, p. 35-51.
- DEL HOYO, J.; ELLIOTT, A. & SARGATAL, J. 1992. Family Sulidae (Ganets and Boobies). In: *Handbook of the Birds of the World*. Barcelona, Lynx Edicions. v. 1, p. 312-325.
- FONSECA-NETO, F.P. 2004. Aves marinhas da Ilha Trindade. In: Branco, J.O. (Org.). *Aves marinhas insulares brasileiras: bioecologia e conservação*. Itajaí, Editora da UNIVALI. p. 119-146.
- HUGHES, J.B.; MARNTIN, G.R. & REYNOLDS, J. 2011. The use of Google Earth™ satellite imagery to detect the nests of masked boobies *Sula dactylatra*. *Wildlife Biology*, 17(2):210-216.
- INSTITUTO BRASILEIRO DO MEIO AMBIENTE E DOS RECURSOS NATURAIS RENOVÁVEIS – IBAMA. 1990. *Plano de Manejo do Parque Nacional Marinho de Fernando de Noronha*. Brasília, Fundação Pró-Natureza – FUNATURA.
- NACINOVIC, J.B. & TEIXEIRA, D.M. 1989. As aves de Fernando de Noronha: uma lista sistemática anotada. *Revista Brasileira de Biologia*, 49(3):709-729.
- OREN, D.C. 1982. A avifauna do Arquipélago de Fernando de Noronha. *Boletim do Museu Paraense Emílio Goeldi. Nova Série Zoologia*, 118:1-22.
- OREN, D.C. 1984. Resultados de uma nova expedição ao Arquipélago de Fernando de Noronha. *Boletim do Museu Paraense Emílio Goeldi. Nova Série Zoologia*, 1(1):19-84.
- SCHREIBER, R.W. & HENSLEY, D.A. 1976. The diets of *Sula dactylatra*, *Sula sula*, and *Fregata minor* on Christmas Island, Pacific Ocean. *Pacific Science*, 30:241-248.
- SCHULZ-NETO, A. 1995. Observando Aves no Parque Nacional Marinho de Fernando de Noronha: *guia de campo*. Brasília, Instituto Brasileiro dos Recursos Naturais Renováveis. 33p.
- SCHULZ-NETO, A. 2004. Aves insulares do arquipélago de Fernando de Noronha. In: Olinto, B. (Org.). *Aves marinhas e insulares brasileiras: bioecologia e conservação*. Itajaí, SC, Editora da UNIVALI. p. 147-168.
- SICK, H. 1997. Atobás: Família Sulidae (4). In: Sick, H. *Ornitologia brasileira*. Rio de Janeiro, Editora Nova Fronteira. p. 190-191.
- SILVA, R.S. 2008. *Aves de Fernando de Noronha*. São Paulo, Avis Brasilis. p. 50-51; 70-75.
- SILVA, R.S. & CAMPOS, F.C. 2006. Registros do atobá-mascarado (*Sula dactylatra*) no Estado de São Paulo. *Revista Brasileira de Ornitologia*, 14(3):283-284.
- SOTO, J.M.R. 2009. Ações Antrópicas Negativas nas Ilhas Oceânicas Brasileiras. *Ilhas Oceânicas Brasileiras: da Pesquisa ao Manejo*. Brasília, MMA/ICMBIO. v. 2, p. 331-350.

Aceito em: 14/12/2016

Publicado em: 16/03/2017

Editor Responsável: Luís Fábio Silveira